DRUG FACTS

Active ingredient(s)	*Purpose*

Use(s)

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Do not use

Do not use if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are

Ask a doctor of pharmacist before use if you are taking a prescription drug

When using this product

When using this product you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating or cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

If pregnant or breast-feeding

Keep out of reach of children

Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed unless advised by your doctor**
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
- adults and children 17 years of age and older:
- fill to top of white section in cap which is marked to indicate the correct dose (17 g)
- stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
- use once a day
- use no more than 7 days
- children 16 years of age or under: ask a doctor
- **do not take more than directed unless advised by your doctor**
- adults and children 17 years of age and older:
- stir and dissolve one packet of powder (17 g) in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
- use once a day
- use no more than 7 days
- children 16 years of age or under: ask a doctor

Other information

Drug Facts

Do not use if you are allergic to polyethylene glycol

- store at 20°-25°C (68°-77°F)
- tamper-evident: do not use if foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken
- store at 20°-25°C (68°-77°F)
- tamper-evident: do not use if foil is open or broken

Distributed by MSD Consumer Care, Inc.,
PO Box 377, Memphis, TN 38151 USA, a subsidiary of
Merck & Co., Inc., Whitehouse Station, NJ USA.

Inactive ingredients

none