DRUG FACTS

Purpose Active ingredient(s)

Use(s)

For the temporary relief of occasional minor irritation, pain, sore mouth and sore throat.

Sore throad warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or give to children under 3 years of age.

Do not use

Ask a doctor before use if you have

Ask a doctor or pharmacist before use if you are

When using this product

do not use more than directed

Stop use and ask a doctor if

- sore mouth symptoms do not improve in 7 days irritation, pain or redness persists or worsens
- swelling, rash or fever develops

If pregnant or breast-feeding

Keep out of reach of children

If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

Adults and children 3 years of age and older:

- apply to the affected area (one spray)
- allow to remain in place for at least 15 seconds, then spit out use every 2 hours or as directed by a doctor or dentist. Children under 12 years of age should be supervised in the use of this product.

Children under 3 years of age: consult a doctor or dentist

Other information

store at 20-25 $^{\rm O}$ C (68-77 $^{\rm O}$ F)

Inactive ingredients

FD&C red #40, flavor, glycerin, purified water, saccharin sodium