# DRUG FACTS

Active ingredient(s) Purpose

## Use(s)

- for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.
- · ask your doctor about other uses for safety coated 81 mg aspirin

# Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- · facial swelling
- shock
- · asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- · are age 60 or older
- · have had stomach ulcers or bleeding problems
- · take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- · take more or for a longer time than directed

#### Do not use

if you are allergic to aspirin or any other pain reliever/fever reducer.

#### Ask a doctor before use if you have

- · stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- · you have high blood pressure, heart disease, liver cirrhosis or kidney disease
- · you have asthma
- · you are taking a diuretic

### Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

# When using this product

#### Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- · have bloody or black stools
- vomit blood
- have stomach pain that does not get better
- allergic reaction occurs. Seek medical help right away.
- new symptoms occur
- pain gets worse or lasts more than 10 days
- redness or swelling is present
  ringing in the ears or loss of hearing occurs

# If pregnant or breast-feeding

## Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away

# Directions

- · drink a full glass of water with each dose
- · adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use

## Other information

store at controlled room temperature 15°-30°C (59°-86°F)

# Inactive ingredients

anhydrous lactose\*\*, black iron oxide\*\*, brown iron oxide\*\*, croscarmellose sodium\*\*, D&C yellow #10, FD&C yellow #6\*\*, hypromellose\*\*, methacrylic acid copolymer, microcrystalline cellulose, mineral oil\*\*, polysorbate 80\*\*, potassium hydroxide\*\*, progelatinized starch\*\*, propylene glycol\*\*, purified water\*\*, shellac\*\*, silicon dioxide, simethicone\*\*, sodium bicarbonate\*\*, sodium hydroxide\*\*, sodium lauryl sulfate\*\*, stearic acid\*\*, talc, titanium dioxide, triethyl citrate, yellow iron oxide\*\*

\*\*contains one or more of these ingredients