ROBITUSSIN SEVERE MULTI-SYMPTOM COUGH COLD FLU- acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride liquid Haleon US Holdings LLC

Drug Facts

Active ingredients (in each 20 ml)

Acetaminophen, USP 650 mg
Dextromethorphan HBr, USP 20 mg
Guaifenesin, USP 400 mg
Phenylephrine HCl, USP 10 mg

Purposes

Pain reliever/Fever reducer

Cough suppressant

Expectorant

Nasal decongestant

Uses

- temporarily relieves these symptoms occurring with a cold or flu:
 - cough due to minor throat and bronchial irritation
 - nasal congestion
 - sinus congestion and pressure
 - minor aches and pains
 - sore throat
 - headache
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes and make coughs more productive

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 doses in any 24-hour period, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert:acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters

rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen ask a doctor or pharmacist.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking any other oral nasal decongestant or stimulant
- taking any other pain reliever/fever reducer

When using this product do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children, even if you do not notice any signs or symptoms.

Directions

- do not take more than 6 doses in any 24-hour period
- do not exceed recommended dosage. Taking more than the recommended dose (overdose) may cause serious liver damage.
- measure only with dosing cup provided
- keep dosing cup with product
- ml = milliliter
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and	20 ml every 4 hours
over	
children under 12 years	do not use

Other information

- each 20 ml contains:sodium 14 mg
- store at 20-25°C (68-77°F)

Inactive ingredients

anhydrous citric acid, edetate disodium, FD&C red no. 40, glycerin, menthol, natural and artificial flavors, polyethylene glycol, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose, triacetin, xanthan gum

Questions or comments?

call weekdays from 9 AM to 5 PM EST at 1-800-762-4675

Made in Canada

For most recent product information, visit www.robitussin.com

Distributed by: Pfizer, Madison, NJ 07940 USA

PRINCIPAL DISPLAY PANEL

ADULT

Robitussin ®

MAXIMUM STRENGTH

SEVERE Multi-Symptom Cough Cold + Flu

ACETAMINOPHEN (Pain Reliever/Fever Reducer)
DEXTROMETHORPHAN HBr (Cough Suppressant)
GUAIFENESIN (Expectorant)
PHENYLEPHRINE HCI (Nasal Decongestant)

- ✓ Cough, Sore Throat
- ✓ Body Aches, Fever
- ✓ Nasal Congestion

✓ Chest Congestion

Non-Drowsy
POWERFUL
multi-symptom relief

CF MAX

For Ages 12 & Over 4 FL OZ (118 ml)



ROBITUSSIN SEVERE MULTI-SYMPTOM COUGH COLD FLU

acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0031-8751
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	650 mg in 20 mL	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL	
		400	

GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	in 20 mL
	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL

Inactive Ingredients		
Ingredient Name	Strength	
EDETATE DISODIUM (UNII: 7FLD91C86K)		
FD&C RED NO. 40 (UNII: WZB9127XOA)		
GLYCERIN (UNII: PDC6A3C0OX)		
MENTHOL, UNSPECIFIED FORM (UNII: L7T10EIP3A)		
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)		
PROPYL GALLATE (UNII: 8D4SNN7V92)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
WATER (UNII: 059QF0KO0R)		
SODIUM BENZOATE (UNII: OJ245FE5EU)		
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)		
SORBITOL (UNII: 506T60A25R)		
SUCRALOSE (UNII: 96K6UQ3ZD4)		
TRIACETIN (UNII: XHX3C3X673)		
XANTHAN GUM (UNII: TTV12P4NEE)		
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)		

Product Characteristics			
Color	red	Score	
Shape		Size	
Flavor	RASPBERRY, CHOCOLATE, MINT	Imprint Code	
Contains			

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0031- 8751-12	1 in 1 CARTON	05/01/2015	
1		118 mL in 1 BOTTLE; Type 1: Convenience Kit of Co-Package		
2	NDC:0031- 8751-18	1 in 1 CARTON	05/01/2015	
2		237 mL in 1 BOTTLE; Type 1: Convenience Kit of Co-Package		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/01/2015	

Labeler - Haleon US Holdings LLC (079944263)

Revised: 4/2024 Haleon US Holdings LLC