

VIDA MIA PAIN RELIEF- aspirin, citric acid monohydrate, and sodium bicarbonate tablet, effervescent

Navarro Discount Pharmacies, LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Vida Mia Pain Relief

Drug Facts

Active ingredients (in each effervescent tablet)	Purpose
Aspirin 325 mg (NSAID)*	Pain reliever
Citric acid 1000 mg	Antacid
Sodium bicarbonate (heat-treated) 1916 mg	Antacid

* nonsteroidal anti-inflammatory drug.

Uses

for the temporary relief of

- heartburn, acid indigestion and sour stomach when accompanied by headache or body aches and pains
- upset stomach with headache from overindulgence in food or drink
- headache, body aches, and pain alone

Warnings

Reye's syndrome

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness

Allergy alert

Aspirin may cause a severe allergic reaction which may include

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning

This product contains an **NSAID**, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

Do not use if

- you have ever had an allergic reaction to any other pain reliever/fever reducer

- you are allergic to aspirin

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

Ask a doctor or pharmacist before use if

- you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.
- you have diabetes, gout or arthritis

When using this product do not exceed the recommended dosage

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away
- symptoms get worse or last more than 10 days
- redness or swelling is present
- ringing in the ears or loss of hearing occurs
- new symptoms occur
- you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better

If pregnant or breast-feeding, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- completely dissolve two tablets in 4 ounces of water before taking

Adults and children 12 years and over	2 tablets every 4 hours do not exceed 8 tablets in 24 hours
60 years and older	2 tablets every 4 hours do not exceed 4 tablets in 24 hours

Other Information

- **each tablet contains:** sodium 540 mg
- protect from excessive heat
- in water, this tablet contains principally the antacid sodium citrate and the analgesic sodium acetylsalicylate

Inactive Ingredients

none

Questions or Comments?

call 1-800-632-2993 M-F 8AM – 4:30PM MST

Distributed By:
MAGELLAN
DISTRIBUTION SOLUTIONS
Medley, Florida 33178

PRINCIPAL DISPLAY PANEL - 36 Tablet Packet Carton

**Compare to
Alka-Seltzer®**

Vida

Mia

Effervescent

Pain

Relief

Aspirin (NSAID) / Pain Reliever, Citric Acid /
Antacid, Sodium Bicarbonate / Antacid.*

**nonsteroidal anti-inflammatory drug*

Fast Relief of:

- **Heartburn • Acid Indigestion**
- **Sour Stomach with Headache**
- **Body Aches and Pains**

36

Tablets



mi salud

Tablets sealed in printed packets for your protection. Do not use if torn or punctured.

PARA INFORMACIÓN EN ESPAÑOL LEA EL INSERTO

Alivio para el Dolor

Pain Relief

Effervescent/Efervescente

Aspirin (NSAID)* / Pain Reliever, Citric Acid / Antacid, Sodium Bicarbonate / Antacid.

**nonsteroidal anti-inflammatory drug*

Aspirina (NSAID = medicamento antiinflamatorio sin esteroides) / Alivia el dolor, Ácido Cítrico/Antiácido, Bicarbonato de Sodio/Antiácido

FAST RELIEF / ALIVIO RÁPIDO

Compare to/Compara con Alka-Seltzer®



mi salud

PARA INFORMACIÓN EN ESPAÑOL LEA EL INSERTO

Effervescent/Efervescente

Pain Relief

Alivio para el Dolor

Aspirin (NSAID)* / Pain Reliever, Citric Acid / Antacid, Sodium Bicarbonate / Antacid.

**nonsteroidal anti-inflammatory drug*

Aspirina (NSAID = medicamento antiinflamatorio sin esteroides) / Alivia el dolor, Ácido Cítrico/Antiácido, Bicarbonato de Sodio/Antiácido

Fast Relief of:

- Heartburn • Acid Indigestion
- Sour Stomach with Headache
- Body Aches and Pains

Alivio Rápido de:

- Ardor Estomacal • Indigestión Ácida
- Acidez Estomacal con Dolor de Cabeza
- Dolores del Cuerpo

36 Tablets Tabletas

BOXD-140 R-0



8 47717 03491 7



mi salud

Pain Relief

Alivio para el Dolor

Place two (2) tablets in 4 oz. of water. Dissolve tablets completely.



You do not have to drink any of the residue that may be on the bottom of the glass. The medicines are in the water.

Take this product any time - morning, noon or night - when you need relief from heartburn, acid indigestion, sour stomach with headache or body aches and pains.



Distributed By:

MAGELAN
DISTRIBUTION SOLUTIONS

Medley, Florida 33178



MADE IN USA



Drug Facts

Active ingredients (in each effervescent tablet)	Purpose
Aspirin 325 mg (NSAID)*	Pain reliever
Citric acid 1000 mg	Antacid
Sodium bicarbonate (heat-treated) 1916 mg	Antacid

*nonsteroidal anti-inflammatory drug.

Uses

- for the temporary relief of
- heartburn, acid indigestion and sour stomach when accompanied by headache or body aches and pains
 - upset stomach with headache from overindulgence in food or drink
 - headache, body aches, and pain alone

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug

Do not use if ■ you have ever had an allergic reaction to any other pain reliever/fever reducer
■ you are allergic to aspirin

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

Ask a doctor or pharmacist before use if

- you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.
- you have diabetes, gout or arthritis

When using this product do not exceed the recommended dosage ▶

Drug Facts (continued)

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away
- symptoms get worse or last more than 10 days
- redness or swelling is present
- ringing in the ears or loss of hearing occurs
- new symptoms occur
- you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better

If pregnant or breast-feeding, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- completely dissolve two tablets in 4 ounces of water before taking

Adults and children 12 years and over	2 tablets every 4 hours do not exceed 8 tablets in 24 hours
60 years and older	2 tablets every 4 hours do not exceed 4 tablets in 24 hours

Other Information

- each tablet contains: sodium 540mg
- protect from excessive heat
- in water, this tablet contains principally the antacid sodium citrate and the analgesic sodium acetylsalicylate

Inactive Ingredients none

Questions or Comments?

call 1-800-632-2993 M-F 8AM - 4:30PM MST

*Alka-Seltzer® is a registered trademark of Bayer HealthCare LLC, Morristown, NJ 07962. This product is not distributed by Bayer HealthCare LLC.

VIDA MIA PAIN RELIEF

aspirin, citric acid monohydrate, and sodium bicarbonate tablet, effervescent

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:59970-205
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Aspirin (UNII: R16CO5Y76E) (Aspirin - UNII:R16CO5Y76E)	Aspirin	325 mg
Citric Acid Monohydrate (UNII: 2968PHW8QP) (Anhydrous Citric Acid - UNII:XF417D3PSL)	Anhydrous Citric Acid	1000 mg
Sodium Bicarbonate (UNII: 8MDF5V39QO) (Bicarbonate Ion - UNII:HN1ZRA3Q20)	Sodium Bicarbonate	1916 mg

Product Characteristics

Color	WHITE	Score	2 pieces
Shape	ROUND	Size	25mm
Flavor		Imprint Code	C11
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59970-205-36	18 in 1 CARTON		
1		2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part343	08/13/2012	

Labeler - Navarro Discount Pharmacies, LLC (094930963)

Revised: 9/2014

Navarro Discount Pharmacies, LLC