SALICYLIC ACID- liquid corn and callus remover liquid Topco Associates LLC

Topco Liquid Corn and Callus Remover

Active ingredient

Salicylic acid 17%w/w

Purpose

Corn and callus remover

Uses

- for the removal of corns and calluses
- relieves pain by removing corns and calluses

Warnings

For external use only.

Flammable

- keep away from fire or flame
- cap bottle tightly and store at room temperature away from heat

Do not use

- on irritated skin
- on any area that is infected or reddened
- if you are a diabetic, or if you have poor blood circulation, except under the advice and supervision of a doctor or podiatrist

When using this product

- avoid contact with eyes
- avoid inhaling vapors
- if product gets into the eyes, flush with water for 15 minutes

Stop and ask a doctor if

If discomfort persists, see your doctor or podiatrist

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

• wash affected area and dry thoroughly

- apply one drop at a time with applicator to sufficiently cover each corn/callus; let dry
- use cushion with hole centered over corn or callus to relieve pressure (additional cushions may be purchased separately)
- repeat this procedure once or twice daily as needed for up to 14 days (until corn/callus is removed)

Other information

- store between 20°C to 25°C (68°F to 77°F)
- close tightly to prevent solidification
- store in a cool place

Inactive ingredients

camphor, castor oil, ethanol, ethyl ether, nitrocellulose

Questions?

Call 1-888-423-0139

Principal Display Panel

Topcare health


Liquid Corn & Callus Remover


SALICYLIC ACID 17%


- Safely Removes Corns and Calluses
- Soft Foam Cushions and Protects

3 CUSHIONS

0.33 FL OZ (9.8mL)







Do not use this product on irritated skin, on any area that is infected or reddened. When using this product awoid contact with eyes, if product gets into the eyes, flush with water for 15 minutes. Avoid inhaling vapors. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. Directions: Wash affected area and dry thoroughly, Apoly one drop at a time with applicator to sufficiently cover each compreadus. Let dry. Use cushions with hole centered over corn or callus to releve pressure. Repeat this procedure once or twice daily as needed for up to 14 days (until convical is is removed). Other information: Store in a cool place. Mactive ingredients: camplor, castro oil, ethanol, ethy ether, introcelluses. Oversions? Cell 1-888-423-0139

Keep packaging for complete drug facts information of lifeburies in the conference of the character oil. ethanol, ethy ether. Introcelluses. Oversions of lifeburies are all the conference of the character of the c

Keep packaging for complete drug facts information obstracting the Cooperation of the Coo

SALICYLIC ACID

liquid corn and callus remover liquid

Product Information					
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:36800-005		
Route of Administration	TOPICAL				

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength

SALICYLIC ACID (UNII: O414PZ4LPZ) (SALICYLIC ACID - UNII:O414PZ4LPZ) SALICYLIC ACID

0.17 mg in 9.8 mL

Inactive Ingredients				
Ingredient Name	Strength			
PYROXYLIN (UNII: KYR8BR2X6O)				
CAMPHOR (NATURAL) (UNII: N20HL7Q941)				
CASTOR OIL (UNII: D5340Y2I9G)				

ı	Packaging					
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
	1	NDC:36800- 005-09	9.8 mL in 1 PACKAGE; Type 0: Not a Combination Product	01/03/2012		

Marketing Information					
Application Number or Monograph Citation	Marketing Start Date	Marketing End Date			
1030	01/03/2012				
	Application Number or Monograph Citation	Application Number or Monograph Marketing Start Citation Date			

Labeler - Topco Associates LLC (006935977)

Revised: 2/2024 Topco Associates LLC