GERI-MOX ANTACID ANTIGAS- aluminum hydroxide, magnesium hydroxide, dimethicone suspension Geri-Care Pharmaceuticals, Corp

geri-mox

Active ingredients (in each 5 mL teaspoonful)

Aluminum hydroxide 200 mg (equivalent to dried gel, USP) Magnesium hydroxide 200 mg Simethicone 20mg

Purposes

Antacid Antigas

Uses

relieves

- heartburn
- sour stomach
- acid indigestion
- the symptoms referred to as gas

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are presently taking a prescription drug.

Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last more than 2 weeks **If pregnant or breast-feeding,** ask a health professional before use.

Keep out of reach of children.

Directions

- shake well before use
- adults and children 12 years of age and older: take 2 to 4 teaspoonfuls between meals, at bedtime, or as directed by a doctor
- do not take more than 16 teaspoonfuls in 24 hours or use the maximum dosage for more than 2 weeks
- children under 12 years: ask a doctor

Other information


- each 5 mL teaspoonful contains: magnesium 85 mg, sodium 3 mg
- store at room temperature
- protect from freezing
- keep tightly closed

Inactive ingredients

benzyl alcohol, butylparaben, carboxymethylcellulose sodium, flavor, hypromellose, microcrystalline cellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

Questions or comments? 1-800-540-3765

package Label



Drug Facts

TAMPER EVIDENT: DO NOT USE IF THE BREAKAWAY BAND ON CAP IS BROKEN OR MISSING.

Active ingredients (in each 5 mL teaspoonful)

Purposes

Uses relieves • heartburn • sour stomach • acid indigestion • the symptoms referred to as gas

Warnings

Ask a doctor before use if you have • kidney disease • a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last for more than 2 weeks

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

Directions • shake well before use

- adults and children 12 years of age and older: take 2 to 4 teaspoonfuls between meals, at bedtime, or as directed by a doctor
- do not take more than 16 teaspoonfuls in 24 hour hours or use the maximum dosage for more than 2 weeks
- children under 12 years: ask a doctor

Other information • each 5 mL teaspoonful contains: magnesium 85 mg, sodium 3 mg • store at room temperature • protect from freezing • keep fightly closed

Inactive ingredients benzyl alcohol, butyl paraben, carboxymethyc ellulose sodium, flavor, hypromellose, microcrystalline celulose, propylparaben, purified water, saccharin sodium, sorbitol solution

Questions or comments? 1-800-540-3765

*This product is not manufactured or distributed by the owner of the registere trademark Maalox®.

Distributed by Geri-Care Pharmaceuticals 1295 Towbin Ave. Lakewood, NJ 08701

REV 1222 910-11212-3

ed or distributed by the owner of the registere

GERI-MOX ANTACID ANTIGAS

aluminum hydroxide, magnesium hydroxide, dimethicone suspension

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:57896-639

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDROXIDE	200 mg in 5 mL
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	200 mg in 5 mL
DIMETHICONE (UNII: 92RU3N3Y10) (DIMETHICONE - UNII:92RU3N3Y10)	DIMETHICONE	20 mg in 5 mL

Inactive Ingredients				
Ingredient Name	Strength			
BENZYL ALCOHOL (UNII: LKG8494WBH)				
BUTYLPARABEN (UNII: 3QPI1U3FV8)				
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K6790BS311)				
HYPROMELLOSES (UNII: 3NXW29V3WO)				
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)				
PROPYLPARABEN (UNII: Z8IX2SC10H)				
WATER (UNII: 059QF0KO0R)				
SACCHARIN SODIUM (UNII: SB8ZUX40TY)				
SORBITOL SOLUTION (UNII: 8KW3E207O2)				

Product Characteristics			
Color		Score	
Shape		Size	
Flavor	MINT (mint)	Imprint Code	
Contains			

ı	Packaging				
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
l	1	NDC:57896-639- 12	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	04/01/2011	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	01/01/2000	

Labeler - Geri-Care Pharmaceuticals, Corp (611196254)

Registrant - GCP Laboratories (965480861)

Establishment			
Name	Address	ID/FEI	Business Operations
GCP Laboratories		965480861	manufacture(57896-639) , pack(57896-639)