

**IBUPROFEN- ibuprofen capsule, liquid filled**  
**Shopko Stores Operating Co., LLC.**

-----

**DRUG FACTS**

**Active ingredient**

**Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)\***  
**(present as the free acid and potassium salt)**  
**\*nonsteroidal anti-inflammatory drug**

**Purpose**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - muscular aches
  - minor pain of arthritis
  - toothache
  - backache
  - the common cold
  - menstrual cramps
- temporarily reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease or asthma
- you are taking a diuretic

**Ask a doctor or pharmacist before use if**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - have bloody or black stools
  - vomit blood
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over:
  - take 1 capsule every 4 to 6 hours while symptoms persist
  - if pain or fever does not respond to 1 capsule, 2 capsules may be used
  - do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

**Other information**

- each capsule contains: **potassium 20 mg**
- read all warnings and directions before use. Keep carton.
- store at 20° to 25°C (68° to 77°F)
- avoid excessive heat above 40°C (104°F). Protect from light.

### **Inactive ingredients**

FD&C blue #1, gelatin, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitan, sorbitol

### **Questions or comments?**

Call toll free: **1-877-753-3935 Monday-Friday 9AM-5PM EST**

### **Principal Display Panel**

†COMPARE TO THE ACTIVE INGREDIENT IN ADVIL® LIQUI-GELS®

### **Ibuprofen**

Capsules, 200 mg

Pain Reliever/Fever Reducer **(NSAID)\***

Gluten Free

Softgels\*\*

(\*\*Liquid Filled Capsules)

†This product is not manufactured or distributed by Pfizer Consumer Healthcare, owner of the registered trademark Advil® Liqui-Gels®.

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

**Do not use if tamper-evident seal under bottle cap imprinted with "*SEALED for YOUR PROTECTION*" is broken or missing.**

**Packaged by: PL Developments**


**200 Hicks Street**

**Westbury, NY 11590, USA**

©2012 Shopko Stores

Operating Co., LLC

### **Package Label**



Shopko Ibuprofen Capsules, 200 mg

IBUPROFEN			
ibuprofen capsule, liquid filled			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:370 12-199
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10 QM) (IBUPROFEN - UNII:WK2XYI10 QM)		IBUPROFEN	200 mg
Inactive Ingredients			
Ingredient Name			Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)			
GELATIN (UNII: 2G86QN327L)			
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)			
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)			
WATER (UNII: 059QF0KO0R)			
SORBITAN (UNII: 6O92ICV9RU)			
SORBITOL (UNII: 506T60A25R)			
Product Characteristics			
Color	BLUE	Score	no score
Shape	CAPSULE	Size	19mm

<b>Flavor</b>		<b>Imprint Code</b>	IB200	
<b>Contains</b>				
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:370 12-199-80	1 in 1 CARTON		
1		80 in 1 BOTTLE		
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>		<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANDA	ANDA078682		07/09/2010	

**Labeler** - Shopko Stores Operating Co., LLC. (023252638)

**Registrant** - P and L Development of New York Corporation (800014821)

Revised: 12/2012

Shopko Stores Operating Co., LLC.