WALGREENS CHILDREN GLYCERIN LAXATIVE- glycerin suppository suppository Walgreens

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Active Ingredient

Active Ingredient Purpose (in each suppository)
Glycerin, USP 1.2 g.....Laxative

Uses

- -relieves occasional constipation (irregularity)
- -generally produces bowel movement in 1/4 to 1 hour

Warnings

For rectal use only

Ask a doctor before use if your child has

Ask a doctor before use if your child has

- stomach pain -nausea -vomiting
- a sudden change in bowel habits lasting more than 2 weeks

Stop use and ask a doctor if

Stop use and ask a doctor if your child has rectal bleeding or fails to have a bowel movement after use of a laxative.

These could be signs of a serious condition.

Keep out of reach of children.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

children 2 to under 6 years 1 suppository, or as directed by doctor

children under 2 years ask a doctor

Insert suppository well up into rectum.

Suppository need not melt completely to produce laxative action.

Other information

-keep tightly closed.


-avoid excessive heat.

Inactive ingredients

Inactive ingredients
purified water, sodium hydroxide, stearic acid

Product Label

FRONT



BACK

This label contains important information. Do not discard Other Information - keep tighty doesd has rectal bleeding or falls to have a bower for 15 minutes; it need not met **Inactive Ingredients** purified water, sodium hydroxibe, stearic acid well Into the rectum and retain not exceed 1 suppository daily to produce lesstive action. Do When using this product your child may swallowed, get medical help or contact a carefully Insert 1 suppository movement after use of a laxative. These I-800-WALGREENS (1-800-926-4733) Stop use and ask doctor if your child or as directed by a doctor STOP PEELING HERE could be signs of a serious condition. have rectal discomfort or a burning Keep out of reach of children. If Ruestions or comments? Polson Control Center right away. Orug Facts (continued) ask a doctor avoid excessive heat Directions 2 to under 6 years of Sensation children children under 2

Directions

children 2 to under 6 years of age retain for 15 minutes; it need not

carefully insert 1 suppository well into the rectum and melt to produce

laxative action. Do not exceed 1 suppository daily or as directed by a doctor.

children under 2

ask a doctor

Do not use

Do not use laxative products for longer than 1 week unless directed by a doctor

When using this product your child

When using this product your child may have rectal discomfort or a burning sensation.

WALGREENS CHILDREN GLYCERIN LAXATIVE

glycerin suppository suppository

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-0440

Route of Administration RECTAL

Active Ingredient/Active Moiety

ı	Ingredient Name	Basis of Strength	Strength
l	GLYCERIN (UNII: PDC6A3C0OX) (GLYCERIN - UNII:PDC6A3C0OX)	GLYCERIN	83 g in 100 g

Inactive Ingredients

Ingredient Name	Strength			
STEARIC ACID (UNII: 4ELV7Z65AP)	8.5 g in 100 g			
SO DIUM HYDRO XIDE (UNII: 55X0 4QC32I)	1.2 g in 100 g			
WATER (UNII: 059QF0KO0R)	7.3 g in 100 g			

Product Characteristics

Color	white (CLEAR)	Score	
Shape	BULLET	Size	32mm
Flavor		Imprint Code	
Contains			

Pac	kag	ın
-----	-----	----

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0440-12	62.5 g in 1 JAR		

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph not final	part334	10/06/2014		

Labeler - Walgreens (008965063)

Establishment				
Name	Address	ID/FEI	Business Operations	
Unipack, Inc.		009248480	manufacture(0363-0440)	

Revised: 10/2014 Walgreens