

**IBUPROFEN, CASEYS 4GOOD- ibuprofen tablets tablet, film coated
Lil' Drug Store Products, Inc.**

Ibuprofen 50ct, Casey's 4good

Drug facts

Drug facts

Active ingredient (in each tablet)

Active ingredient (in each tablet)

Ibuprofen USP, 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Purpose

Pain reliever/fever reducer

Uses

Uses

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- minor pain of arthritis
- toothache
- backache
- the common cold
- menstrual cramps
- temporarily reduces fever

Warnings

Warnings

Allergy alert

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- have bloody or black stools
- feel faint
- vomit blood
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
- chest pain
- slurred speech
- leg swelling
- trouble breathing
- weakness in one part or side of the body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and older:
- take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

Other information

- store between 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature]
- avoid excessive heat above 40°C (104°F)

Inactive ingredients

Inactive ingredients

colloidal silicon dioxide, iron oxide red, maize starch, polyethylene glycol, povidone k-30, pregelatinized starch, sodium starch glycolate, stearic acid, talc, titanium dioxide

Questions or comments?

Questions or comments? Call toll-free **1-877-507-6516** (M-F 8AM-4:30PM CST)

Ibuprofen 50ct, Casey's 4good ™

SEE NEW WARNINGS

Ibuprofen

Tablets USP, 200 mg (NSAID)*

Pain Reliever/Fever Reducer

Compare to the
active ingredient in
*Advil Tablets***

[tablet image]

actual

size

[casey's logo]

50 COATED TABLETS



IBUPROFEN, CASEYS 4GOOD

ibuprofen tablets tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:66715-6307
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
TALC (UNII: 7SEV7J4R1U)	
STARCH, CORN (UNII: O8232NY3SJ)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

Product Characteristics

Color	brown	Score	no score
Shape	ROUND	Size	9mm
Flavor		Imprint Code	G2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-6307-4	1 in 1 CARTON	08/01/2023	
1		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079174	08/01/2023	

Labeler - Lil' Drug Store Products, Inc. (093103646)

Revised: 9/2023

Lil' Drug Store Products, Inc.