

**HEADACHE RELIEF EXTRA STRENGTH- acetaminophen, aspirin, caffeine tablet,
film coated**

Meijer Distribution Inc

Meijer 44-334B

Active ingredients (in each caplet)

Acetaminophen 250 mg

Aspirin 250 mg (NSAID)*

Caffeine 65 mg

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Pain reliever

Pain reliever aid

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - arthritis
 - a cold
 - muscular aches
 - toothache
 - premenstrual and menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- shock
- facial swelling
- hives
- asthma (wheezing)

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters

- rash

If a skin reaction occurs, stop use and seek medical help right away.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Do not use

- if you have ever had an allergic reaction to acetaminophen, aspirin, or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout, or arthritis
- taking any other drug or are under a doctor's care for any serious condition

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools

- vomit blood
- have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- redness or swelling is present
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets every 6 hours. Do not take more than 8 caplets in 24 hours.
- children under 12 years: ask a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, crospovidone, hypromellose, microcrystalline cellulose, povidone, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?

1-800-426-9391

Principal Display Panel

NDC 41250-334-12

Meijer®

Compare to Excedrin® Extra Strength active ingredients†

EXTRA STRENGTH

headache relief

Acetaminophen † Aspirin (**NSAID**) † Caffeine

Pain Reliever † Pain Reliever Aid

100 Caplets

Actual


Size

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

†This product is not manufactured or distributed by GSK Consumer Healthcare SARL, owner of the registered trademark Excedrin® Extra Strength.
50844 REV1121E33412

**DIST. BY MEIJER
DISTRIBUTION, INC.
GRAND RAPIDS, MI 49544
www.meijer.com**

**OUR QUALITY
GUARANTEE**
WWW.MEIJER.COM/SATISFACTION



HEADACHE RELIEF EXTRA STRENGTH

acetaminophen, aspirin, caffeine tablet, film coated

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:41250-334

Route of Administration

ORAL

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

Inactive Ingredients	
Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics			
Color	white	Score	no score
Shape	OVAL	Size	17mm
Flavor		Imprint Code	44;334
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-334-15	1 in 1 CARTON	03/13/2008	
1		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:41250-334-12	1 in 1 CARTON	03/13/2008	
2		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:41250-334-98	2 in 1 CARTON	03/13/2008	
3		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
4	NDC:41250-334-08	1 in 1 CARTON	03/13/2008	12/01/2023
4		24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information			
Marketing	Application Number or Monograph	Marketing Start	Marketing End

Category	Citation	Date	Date
OTC Monograph Drug	M013	03/13/2008	

Labeler - Meijer Distribution Inc (006959555)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(41250-334)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(41250-334)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(41250-334)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(41250-334)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(41250-334)