


BENZONATATE- benzonatate capsule
Liberty Pharmaceuticals, Inc.

BENZONATATE - INTERGEL

DESCRIPTION

Benzonatate, a non-narcotic oral antitussive agent, is 2, 5, 8, 11, 14, 17, 20, 23, 26-nonaoxaocctacosan-28-yl p-(butylamino) benzoate; with a molecular weight of 603.7.



Each soft gelatin capsule, for oral administration, contains 100 mg or 200 mg of benzonatate USP.

Benzonatate Capsules, USP also contain the following inactive ingredients: D&C Yellow # 10, gelatin, glycerin, purified water, methylparaben, propylparaben and titanium dioxide.

CLINICAL PHARMACOLOGY

Benzonatate acts peripherally by anesthetizing the stretch receptors located in the respiratory passages, lungs, and pleura by dampening their activity and thereby reducing the cough reflex at its source. It begins to act within 15 to 20 minutes and its effect lasts for 3 to 8 hours. Benzonatate has no inhibitory effect on the respiratory center in recommended dosage.

INDICATIONS AND USAGE

Benzonatate is indicated for the symptomatic relief of cough.

CONTRAINDICATIONS

Hypersensitivity to benzonatate or related compounds.

WARNINGS

Severe hypersensitivity reactions (including bronchospasm, laryngospasm and cardiovascular collapse) have been reported which are possibly related to local anesthesia from sucking or chewing the capsule instead of swallowing it. Severe reactions have required intervention with vasopressor agents and supportive measures.

Isolated instances of bizarre behavior, including mental confusion and visual hallucinations, have also

been reported in patients taking benzonatate in combination with other prescribed drugs.

PRECAUTIONS

Benzonatate is chemically related to anesthetic agents of the para-aminobenzoic acid class (e.g. procaine; tetracaine) and has been associated with adverse CNS effects possibly related to a prior sensitivity to related agents or interaction with concomitant medication.

Information for patients :

Release of benzonatate from the capsule in the mouth can produce a temporary local anesthesia of the oral mucosa and choking could occur. Therefore, the capsules should be swallowed without chewing.

Usage in Pregnancy:

Pregnancy Category C. Animal reproduction studies have not been conducted with benzonatate. It is also not known whether benzonatate can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Benzonatate should be given to a pregnant woman only if clearly needed.

Nursing Mothers :

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk caution should be exercised when benzonatate is administered to a nursing woman.

Carcinogenesis, mutagenesis, impairment of fertility:

Carcinogenicity, mutagenicity, and reproduction studies have not been conducted with benzonatate.

Pediatric Use:

Safety and effectiveness in children below the age of 10 have not been established.

ADVERSE REACTIONS

Potential Adverse Reactions to benzonatate may include:

Hypersensitivity reactions including bronchospasm, laryngospasm, cardiovascular collapse possibly related to local anesthesia from chewing or sucking the capsule.

CNS: sedation; headache; dizziness; mental confusion; visual hallucinations.

GI: constipation; nausea; GI upset.

Dermatologic: pruritus; skin eruptions.

Other: nasal congestion; sensation of burning in the eyes; vague "chilly" sensation; numbness of the chest; hypersensitivity.

Rare instances of deliberate or accidental overdose have resulted in death.

OVERDOSAGE

Overdose may result in death.

The drug is chemically related to tetracaine and other topical anesthetics and shares various aspects of their pharmacology and toxicology. Drugs of this type are generally well absorbed after ingestion.

Signs and Symptoms:

If capsules are chewed or dissolved in the mouth, oropharyngeal anesthesia will develop rapidly. CNS stimulation may cause restlessness and tremors which may proceed to clonic convulsions followed by profound CNS depression.

Treatment:

Evacuate gastric contents and administer copious amounts of activated charcoal slurry. Even in the conscious patient, cough and gag reflexes may be so depressed as to necessitate special attention to protection against aspiration of gastric contents and orally administered materials. Convulsions should be treated with a short-acting barbiturate given intravenously and carefully titrated for the smallest effective dosage. Intensive support of respiration and cardiovascular - renal function is an essential feature of the treatment of severe intoxication from overdose.

Do not use CNS stimulants.

DOSAGE AND ADMINISTRATION

Adults and Children over 10 years of age: Usual dose is one 100 mg or 200 mg capsule t.i.d. as required. If necessary, up to 600 mg daily may be given.

HOW SUPPLIED

Benzonatate Capsules USP, 100 mg: Yellow soft gelatin capsules, imprinted "ASC" on one side and "105" on the other side, available in bottles of 10's (NDC 00440-1185-10).

Store at controlled room temperature 20° - 25°C (68° - 77°F) [See USP].

API-01, 5/07

Mfd by
Intergel Division of IVC Industries, Inc.
Irvington, NJ 07111

Repackaged By:
Aidarex Pharmaceuticals, LLC
Corona, CA 92880

RX ONLY	LIBERTY Pharmaceuticals Inc.	PACKAGED BY: AIDAREX PHARMACEUTICALS LLC CORONA, CA 92880
	NDC: 00440-1185-10	
	BENZONATATE	
	GENERIC FOR BRAND NAME: TESSALON PERLES	
	100mg	10 CAPS
	EACH CAPSULE CONTAINS THE FOLLOWING ACTIVE INGREDIENTS BENZONATATE..... 100mg	
	MFG: INTERGEL DIV. OF IVC INDUSTRIES, INC., IRVINGTON, NJ 07111	CAUTION: KEEP OUT OF REACH OF CHILDREN. STORE AT CONTROLLED ROOM TEMPERATURE 15-30 C (59-86) F SEE PACKAGE INSERT.
		DOCTOR _____ DATE _____
		PATIENT: _____
		YELLOW OVAL CAPSULE W/ ASC ON ONE SIDE & 105 ON REVERSE
		Take _____ Cap(s) Every _____ Hour(s)
		_____ Time(s) a day

BENZONATATE

benzonatate capsule

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0440-1185(NDC:67877-105)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZONATATE (UNII: 5P4DHS6ENR) (BENZONATATE - UNII:5P4DHS6ENR)	BENZONATATE	100 mg

Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
WATER (UNII: 059QF0KO0R)	
METHYL PARABEN (UNII: A2I8C7HI9T)	
PROPYL PARABEN (UNII: Z8IX2SC1OH)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	YELLOW	Score	no score
Shape	CAPSULE	Size	19mm
Flavor		Imprint Code	ASC;105
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0440-1185-10	10 in 1 BOTTLE, PLASTIC		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA040627	07/25/2007	

Labeler - Liberty Pharmaceuticals, Inc. (012568840)