

STOMACH LIL DRUG STORE PRODUCTS- bismuth subsalicylate liquid
Lil' Drug Store Products, Inc

Lil' Drug Store® **Stomach**

Drug Facts

Active ingredient (in each 30 mL dose)

Bismuth subsalicylate 525 mg

Purpose

Upset stomach reliever/Antidiarrheal

Uses

relieves

- diarrhea
- travelers' diarrhea
- upset stomach due to overindulgence in food and drink, including
 - heartburn
 - gas
 - indigestion
 - nausea
 - fullness
 - belching

Warnings

Reye's syndrome

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert

Contains salicylate. Do not take if you are:

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use if you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for

- diabetes
- gout
- arthritis
- anticoagulation (thinning the blood)

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- mL = milliliter
- shake well before using
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- adults and children 12 years and over:
 - 30 mL (1 dose) every ½ hour or 60 mL (2 doses) every hour as needed for diarrhea/travelers' diarrhea
 - 30 mL (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
 - do not exceed 8 doses (240 mL) in 24 hours
 - use until diarrhea stops but not more than 2 days
- children under 12 years of age: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

- **each 30 mL contains:** potassium 8 mg, sodium 6 mg
- **each 30 mL contains:** salicylate 206 mg
- low sodium
- keep tightly closed
- protect from freezing
- avoid excessive heat (over 104°F or 40°C)

Inactive ingredients

benzoic acid, D&C red #22, D&C red #28, flavor, glycerin, purified water, sucralose, xanthan gum

Questions?

Call toll free **1-877-507-6516** (M-F 8AM-4:30PM CST)

PRINCIPAL DISPLAY PANEL - 237 mL Bottle Label

QUALITY
GUARANTEED

Compare to the Active
Ingredient in Pepto-Bismol®*

Stomach

Bismuth Subsalicylate 525 mg
Upset Stomach Reliever/Antidiarrheal

Relieves

- Nausea • Heartburn • Indigestion
- Upset Stomach • Diarrhea

8 FL OZ
(237 mL)

Lil'
Drug Store®

<p>Drug Facts (continued)</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)</p> <p>Directions • mL = milliliter • shake well before using</p> <ul style="list-style-type: none">• measure only with dosing cup provided. Do not use any other dosing device.• keep dosing cup with product• adults and children 12 years and over:<ul style="list-style-type: none">• 30 mL (1 dose) every ½ hour or 60 mL (2 doses) every hour as needed for diarrhea/travelers' diarrhea• 30 mL (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)• do not exceed 8 doses (240 mL) in 24 hours• use until diarrhea stops but not more than 2 days• children under 12 years of age: ask a doctor• drink plenty of clear fluids to help prevent dehydration caused by diarrhea <p>Other Information • each 30 mL contains: potassium 8 mg, sodium 6 mg</p> <ul style="list-style-type: none">• each 30 mL contains: salicylate 206 mg• low sodium • keep tightly closed• protect from freezing • avoid excessive heat (over 104°F or 40°C) <p>Inactive ingredients benzoic acid, D&C red #22, D&C red #28, flavor, glycerin, purified water, sucralose, xanthan gum</p> <p>Questions? Call toll free 1-877-507-6516 (M-F 8AM-4:30PM CST)</p>	<p>Compare to the Active Ingredient in Pepto-Bismol®*</p> <p>Stomach</p> <p>Bismuth Subsalicylate 525 mg Upset Stomach Reliever/Antidiarrheal</p> <p>Relieves</p> <ul style="list-style-type: none">• Nausea • Heartburn • Indigestion• Upset Stomach • Diarrhea <p>8 FL OZ (237 mL)</p> 	<p>TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.</p> <p>Drug Facts</p> <table border="1"><thead><tr><th>Active Ingredient (In each 30 mL dose)</th><th>Purpose</th></tr></thead><tbody><tr><td>Bismuth subsalicylate 525 mg...</td><td>Upset stomach reliever/Antidiarrheal</td></tr></tbody></table> <p>Uses relieves • diarrhea • travelers' diarrhea</p> <ul style="list-style-type: none">• upset stomach due to overindulgence in food and drink, including• heartburn • gas • indigestion • nausea • fullness • belching <p>Warnings</p> <p>Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.</p> <p>Allergy alert: Contains salicylate. Do not take if you are:</p> <ul style="list-style-type: none">• allergic to salicylates (including aspirin)• taking other salicylate products <p>Do not use if you have</p> <ul style="list-style-type: none">• an ulcer • a bleeding problem • bloody or black stool <p>Ask a doctor before use if you have • fever • mucus in the stool</p> <p>Ask a doctor or pharmacist before use if you are taking any drug for</p> <ul style="list-style-type: none">• diabetes • gout • arthritis• anticoagulation (thinning the blood) <p>When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur.</p> <p>Stop use and ask a doctor if • symptoms get worse or last more than 2 days • ringing in the ears or loss of hearing occurs • diarrhea lasts more than 2 days</p>	Active Ingredient (In each 30 mL dose)	Purpose	Bismuth subsalicylate 525 mg...	Upset stomach reliever/Antidiarrheal
Active Ingredient (In each 30 mL dose)	Purpose					
Bismuth subsalicylate 525 mg...	Upset stomach reliever/Antidiarrheal					

*This product is not manufactured or distributed by The Procter & Gamble Company. Pepto-Bismol® is a registered trademark of The Procter & Gamble Company.



Product manufactured for:
Lil' Drug Store Products, Inc.
9300 Earhart Lane SW
Cedar Rapids, IA 52404
88564-US-12-19
PLD-A492A LB007266

STOMACH LIL DRUG STORE PRODUCTS

bismuth subsalicylate liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:66715-6855
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD18YE)	BISMUTH SUBSALICYLATE	525 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
BENZOIC ACID (UNII: 8SKN0B0MIM)	
D&C RED NO. 22 (UNII: 1678RKX8RT)	
D&C RED NO. 28 (UNII: 7671P0Y5NH)	
GLYCERIN (UNII: PDC6A3C0OX)	
WATER (UNII: 059QF0K00R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-6855-4	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/01/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	12/01/2014	

Labeler - Lil' Drug Store Products, Inc (093103646)

Revised: 12/2023

Lil' Drug Store Products, Inc