

BENCOCAINE TOPICAL ANESTHETIC- benzocaine gel

Benco Dental

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Active Ingredients

Benzocaine 200 mg (in each g)

Purpose

Oral Anesthetic

For oral mucosal use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.

Warnings

Warnings

Methemoglobinemia

warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray or blue colored skin (cyanosis)
- headache
- rapid heart rate

- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy Alert

Allergy Alert Do not use on patients with a history of allergies to local anesthetics

such as procaine, butacaine, benzocaine or other "caine" anesthetics.

Do not use

- for more than 7 days unless directed by a physician. If sore mouth symptoms do not improve in 7 days; irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your physician promptly.
- for teething
- in children under 2 years of age

When using this product

Avoid contact with eyes. If contact occurs, flush with water.

Do not exceed recommended dosage

If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

If pregnant or breast feeding,

ask a physician before use.

Keep out of reach of children

Directions

- Apply only amount needed to the oral mucosa to prevent or relieve pain.
- children under 2 years of age: do not use

Other Information

Store at 59 -86°F (15-30°C).
Protect from freezing.

Inactive Ingredients

flavoring, PEG 3350, PEG 400, sodium saccharin.

Questions or Comments?

800.GO.BENCO or benco.com

Distributed by
Benco Dental

800.GO.BENCO • benco.com
295 CenterPoint Blvd., Pittston, PA 18640

BencoCaine™
topical anesthetic gel

20% benzocaine
GLUTEN FREE

DYE-FREE

1 fl. oz. (30ml)

OPEN HERE
Made In U.S.A.

directed by a physician. If sore mouth symptoms do not improve in 7 days; irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your physician promptly.

- for teething
- in children under 2 years of age

When using this product
Avoid contact with eyes. If contact occurs, flush with water.

Do not exceed recommended dosage. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

If pregnant or breast feeding, ask a physician before use.

Keep out of reach of children.

Directions

- Apply only amount needed to the oral mucosa to prevent or relieve pain.
- children under 2 years of age: do not use

Other information
Store at 59° - 86° F (15 - 30° C). Protect from freezing.

Inactive ingredients
flavoring, PEG 3350, PEG 400, sodium saccharin.

Questions or comments?
800.GO.BENCO or benco.com

99091791 Rev 10/2020

Drug Facts

Active ingredients
Benzocaine 200 mg (in each g)

Purpose
Oral Anesthetic

Use
For oral mucosal use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.

Warnings
Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy Alert: Do not use on patients with a history of allergies to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

Do not use

- for more than 7 days unless

BENCOCAINE TOPICAL ANESTHETIC

benzocaine gel

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:66975-341
Route of Administration	DENTAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	200 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	

Product Characteristics

Color	white (DYE-FREE)	Score	
Shape		Size	
Flavor	BUBBLE GUM	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66975-341-30	30 g in 1 JAR; Type 0: Not a Combination Product	10/01/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part356	10/01/2020	

Labeler - Benco Dental (015108087)

Revised: 9/2023

Benco Dental