

**POTASSIUM CHLORIDE IN LACTATED RINGERS AND DEXTROSE- dextrose monohydrate, sodium chloride, sodium lactate, potassium chloride, calcium chloride injection, solution
Baxter Healthcare Corporation**

**Potassium Chloride in Lactated Ringer’s and 5% Dextrose Injection, USP in Plastic Container
VIAFLEX Plus Container**


DESCRIPTION

Potassium Chloride in Lactated Ringer’s and 5% Dextrose Injection, USP is a sterile, nonpyrogenic solution for fluid and electrolyte replenishment and caloric supply in a single dose container for intravenous administration. It contains no antimicrobial agents. Composition, osmolarity, pH, ionic concentration and caloric content are shown below:

Table 1.

Potassium Chloride in Lactated Ringer’s and 5% Dextrose Injection, USP	Size (mL)	Composition (g/L)					*Osmolarity (mOsmol/L) (calc.)
		†Dextrose Hydrous, USP	Sodium Chloride, USP (NaCl)	Sodium Lactate, (C ₃ H ₅ NaO ₃)	Potassium Chloride, USP (KCl)	Calcium Chloride, USP (CaCl ₂ •2H ₂ O)	
mEq Potassium added							
20 mEq	1000	50	6	3.1	1.79	0.2	565

* Normal physiologic osmolarity range is approximately 280 to 310 mOsmol/L.
† The chemical structure for Dextrose Hydrous, USP is shown below.



Dextrose is derived from corn.

Dextrose is derived from corn.

Table 2.

Potassium Chloride in Lactated Ringer’s and 5% Dextrose Injection, USP	pH	Ionic Concentration (mEq/L)					Caloric Content (kcal/L)
		Sodium	Potassium	Calcium	Chloride	Lactate	
mEq Potassium added							

20 mEq	5.0 (3.5 to 6.5)	130	24	3	129	28	170
--------	---------------------	-----	-----------	---	-----	----	-----

The VIAFLEX Plus plastic container is fabricated from a specially formulated polyvinyl chloride (PL 146 Plastic). VIAFLEX Plus on the container indicates the presence of a drug additive in a drug vehicle. The VIAFLEX Plus plastic container system utilizes the same container as the VIAFLEX plastic container system. The amount of water that can permeate from inside the container into the overwrap is insufficient to affect the solution significantly. Solutions in contact with the plastic container can leach out certain of its chemical components in very small amounts within the expiration period, e.g., di-2-ethylhexyl phthalate (DEHP), up to 5 parts per million. However, the safety of the plastic has been confirmed in tests in animals according to USP biological tests for plastic containers as well as by tissue culture toxicity studies.

CLINICAL PHARMACOLOGY

Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP have value as a source of water, electrolytes, and calories. It is capable of inducing diuresis depending on the clinical condition of the patient.

Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP produce a metabolic alkalinizing effect. Lactate ions are metabolized ultimately to carbon dioxide and water, which requires the consumption of hydrogen cations.

INDICATIONS AND USAGE

Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP are indicated as a source of water, electrolytes, and calories or as alkalinizing agents.

CONTRAINDICATIONS

Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP is contraindicated in:

- Newborns (\leq 28 days of age), receiving concomitant treatment with ceftriaxone, even if separate infusion lines are used due to the risk of fatal ceftriaxone-calcium salt precipitation in the neonate's bloodstream.
- In patients older than 28 days, including adults, administered ceftriaxone simultaneously through the same infusion line (e.g., via Y-connector). If the same infusion line is used for sequential administration, the line must be thoroughly flushed between infusions with a compatible fluid.
- Patients with a known hypersensitivity to sodium lactate (see **WARNINGS**).
- Patients with hyperkalemia (see **WARNINGS**).

WARNINGS

Hypersensitivity Reactions

Hypersensitivity and infusion reactions have been reported with Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP (see **ADVERSE REACTIONS**). Stop the infusion immediately if any signs or symptoms of a suspected hypersensitivity reaction develop. Appropriate therapeutic countermeasures must be instituted as clinically indicated.

Electrolyte Imbalances

Fluid Overload

Depending on the volume and rate of infusion, the intravenous administration of Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP can cause electrolyte disturbances such as overhydration and congested states, including pulmonary congestion and edema. Avoid Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP in patients with or at risk for fluid and/or solute overloading. If use cannot be avoided, monitor fluid balance, electrolyte concentrations and acid base

balance, as needed and especially during prolonged use.

Hyperglycemia and Hyperosmolar Hyperglycemic State

Administration of solutions containing dextrose and lactate in patients with impaired glucose tolerance or diabetes mellitus may worsen hyperglycemia (see **PRECAUTIONS, Pediatric Use**). Administration of dextrose at a rate exceeding the patient's utilization rate may lead to hyperglycemia, coma, and death.

Hyperglycemia is associated with an increase in serum osmolality, resulting in osmotic diuresis, dehydration and electrolyte losses.

Patients with underlying central nervous system disease and renal impairment who receive dextrose infusions, may be at greater risk of developing hyperosmolar hyperglycemic state.

Monitor blood glucose concentrations and treat hyperglycemia to maintain concentrations within normal limits while administering Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP. Insulin may be administered or adjusted to maintain optimal blood glucose concentrations.

Hyperkalemia

Potassium-containing solutions, including Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP may increase the risk of hyperkalemia.

Patients at increased risk of developing hyperkalemia include those:

- With conditions predisposing to hyperkalemia and/or associated with increased sensitivity to potassium, such as patients with severe renal impairment, acute dehydration, extensive tissue injury or burns, certain cardiac disorders such as congestive heart failure.
- Treated concurrently or recently with agents or products that cause or increase the risk of hyperkalemia (see **PRECAUTIONS**).

Avoid use of Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP in patients with, or at risk for, hyperkalemia. If use cannot be avoided, monitor serum potassium concentrations.

To avoid life threatening hyperkalemia, do not administer Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP as an intravenous push (i.e., intravenous injection manually with a syringe connected to the intravenous access) without a quantitative infusion device.

Alkalosis

Because lactate is metabolized to bicarbonate, administration of Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP may result in, or worsen, metabolic alkalosis. Avoid Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP in patients with alkalosis or at risk for alkalosis.

Hyponatremia

Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP may cause hyponatremia. Glucose intravenous infusions are usually isotonic solutions. In the body, however, glucose containing fluids can become extremely physiologically hypotonic due to rapid glucose metabolism. Monitoring of serum sodium is particularly important for hypotonic fluids.

Hyponatremia can lead to acute hyponatremic encephalopathy characterized by headache, nausea, seizures, lethargy, and vomiting. Patients with brain edema are at particular risk of severe, irreversible and life-threatening brain injury.

The risk of hospital-acquired hyponatremia is increased in patients with cardiac or pulmonary failure, and in patients with non-osmotic vasopressin release (including SIADH) treated with high volume of hypotonic Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP.

The risk for hyponatremia is increased in pediatric patients, elderly patients, postoperative patients, those with psychogenic polydipsia, and in patients treated with medications that increase the risk of hyponatremia (such as diuretics, certain antiepileptic and psychotropic medications). See **PRECAUTIONS, Drug Interactions and Pediatric Use**.

Patients at increased risk for developing complications of hyponatremia such as hyponatremic encephalopathy, include pediatric patients, women (in particular, premenopausal women), patients with hypoxemia, and patients with underlying central nervous system disease. Avoid Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP in patients with or at risk for hyponatremia. If use cannot be avoided, monitor serum sodium concentrations.

Rapid correction of hyponatremia is potentially dangerous with risk of serious neurologic complications. Brain adaptations reducing risk of cerebral edema make the brain vulnerable to injury when chronic hyponatremia is too rapidly corrected, which is known as osmotic demyelination syndrome (ODS). To avoid complications, monitor serum sodium and chloride concentrations, fluid status, acid-base balance, and signs of neurologic complications.

PRECAUTIONS

Patients with Renal Impairment

Administration of Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP in patients with or at risk of severe renal impairment, may result in hyperkalemia and/or fluid overload (see **WARNINGS**). Avoid Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP in patients with severe renal impairment or conditions that may cause sodium and/or potassium retention, fluid overload, or edema. If use cannot be avoided, monitor patients with severe renal impairment for development of these adverse reactions.

Patients with Hepatic Impairment

In patients with severe hepatic impairment, lactate metabolism may be impaired and Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP may not produce alkalization. Consider when monitoring serum lactate levels.

Hypercalcemia

Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP contains calcium salts and may cause hypercalcemia. Avoid administration of Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP in patients with hypercalcemia or conditions predisposing to hypercalcemia; and in patients with calcium renal calculi or history of such calculi.

Vein Thrombosis

Use a peripheral vein if the final dextrose concentration is 5% or less and the osmolality is approximately less than 900 mOsm/L. Consider central vein administration of hypertonic solutions with osmolality of 900 mOsmol or more to avoid venous irritation, including phlebitis. Monitoring of Serum Lactate Levels

Administration of Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP may result in an iatrogenic increase in serum lactate levels and interfere with interpretation of serum lactate levels in patients with severe metabolic acidosis including lactic acidosis.

Drug Interactions

Ceftriaxone

For information on interaction with ceftriaxone – see **CONTRAINDICATIONS**

Other Products that Cause Hyperkalemia

Administration of Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP to patients treated concurrently or recently with products that are associated with hyperkalemia increases the risk of severe and potentially fatal hyperkalemia, in particular in the presence of other risk factors for hyperkalemia.

Avoid use of Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP to patients receiving such products (e.g., potassium sparing diuretics, ACE inhibitors, angiotensin II receptor antagonists, or the immunosuppressants tacrolimus and

cyclosporine). If use cannot be avoided, monitor serum potassium concentrations.

Other Products that Affect Fluid and/or Electrolyte Balance

Administration of Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP to patients treated concomitantly with drugs associated with sodium and fluid retention may increase the risk of hypernatremia and volume overload. Avoid use of Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP in patients receiving such products, such as corticosteroids or corticotropin. If use cannot be avoided, monitor serum electrolytes, fluid balance and acid-base balance.

Other Drugs that Increase the Risk of Hyponatremia

Administration of Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP to patients treated concomitantly with medications associated with hyponatremia may increase the risk of developing hyponatremia.

Avoid use of Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP in patients receiving products, such as diuretics, and certain antiepileptic and psychotropic medications. Drugs that increase the vasopressin effect reduce renal electrolyte free water excretion and may also increase the risk of hyponatremia following treatment with intravenous fluids. If use cannot be avoided, monitor serum sodium concentrations.

Lithium

Renal sodium and lithium clearance may be increased during administration of Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP. Avoid use of Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP in patients receiving lithium. If use cannot be avoided, monitor serum lithium concentrations during concomitant use.

Digoxin

Administration of calcium may increase the effects of digitalis and lead to serious or fatal cardiac arrhythmia. In patients treated with digoxin, consider reducing the volume, and/or rate of administration of Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP.

Other Drugs that Increase the Risk of Hypercalcemia

Avoid Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP in patients treated with thiazide diuretics or vitamin D, as these can increase the risk of hypercalcemia.

Drugs with pH Dependent Renal Elimination

Due to the alkalinizing action of lactate (formation of bicarbonate), Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP may interfere with the elimination of drugs with pH dependent renal elimination. Renal clearance of acidic drugs may be increased. Renal clearance of alkaline drugs may be decreased.

Pregnancy

There are no adequate, well-controlled studies with Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP in pregnant women and animal reproduction studies have not been conducted with this drug. Therefore, it is not known whether Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP can cause fetal harm when administered to a pregnant woman. Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP should be given during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Labor and Delivery

Intrapartum maternal intravenous infusion of glucose-containing solutions may produce maternal hyperglycemia with subsequent fetal hyperglycemia and fetal metabolic acidosis as well as rebound hypoglycemia in the neonate. Fetal hyperglycemia can result in increased fetal insulin levels which may result in neonatal hypoglycemia following delivery. Consider the potential risks and benefits for each specific patient before administering Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP.

Nursing Mothers

It is not known whether this drug is present in human milk. Because many drugs are present in human milk, caution should be exercised when Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP is administered to a nursing mother.

Pediatric Use

The use of Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP in pediatric patients is based on clinical practice (see DOSAGE AND ADMINISTRATION).

Neonates, especially premature infants with low birth weight, are at increased risk of developing hypo- or hyperglycemia and therefore need close monitoring during treatment with intravenous glucose solutions to ensure adequate glycemic control in order to avoid potential long-term adverse effects.

Closely monitor plasma electrolyte concentrations in pediatric patients who may have impaired ability to regulate fluids and electrolytes. In very low birth weight infants, excessive or rapid administration of Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP may result in increased serum osmolality and risk of intracerebral hemorrhage.

Children (including neonates and older children) are at increased risk of developing hyponatremia as well as for developing hyponatremic encephalopathy.

Administration of a lactate-containing intravenous solution to infants should take into account that the liver and kidneys are still maturing during the first year of life, which also affects the biotransformation and renal excretion of lactate.

Geriatric Use

Geriatric patients are at increased risk of developing electrolyte imbalances. Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP is known to be substantially excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Therefore, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy. Consider monitoring renal function in elderly patients.

ADVERSE REACTIONS

Post-Marketing Adverse Reactions

The following adverse reactions associated with the use of Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP were identified in clinical trials or postmarketing reports. Because postmarketing reactions were reported voluntarily from a population of uncertain size, it is not always possible to estimate their frequency, reliably, or to establish a causal relationship to drug exposure.

Hypersensitivity and infusion reactions: angioedema, chest pain, chest discomfort, decreased heart rate, tachycardia, blood pressure decreased, respiratory distress, bronchospasm, dyspnea, cough, urticaria, rash, pruritus, erythema, flushing, throat irritation, paresthesias, oral hypoesthesia, dyspepsia, nausea, anxiety, pyrexia and headache.

General Disorders and Administration Site Conditions: phlebitis, extravasation, infusion site inflammation, infusion site swelling, infusion site rash, infusion site pruritus, infusion site erythema, infusion site pain, infusion site burning, infusion site anesthesia (numbness)

Metabolism and Nutrition Disorders: hyperkalemia, hypervolemia, hyponatremia

Nervous System Disorders: hyponatremic encephalopathy.

OVERDOSAGE

Excessive administration of Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP can cause:

- hyperkalemia and hypernatremia, especially in patients with severe renal

- impairment.
- fluid overload (which can lead to pulmonary and/or peripheral edema).
- hyperglycemia, hyperosmolality and osmotic diuresis, dehydration and electrolyte loss.
- metabolic alkalosis with or without hypokalemia.
- loss of bicarbonate with an acidifying effect.
- hypercalcemia.

See **WARNINGS** and **ADVERSE REACTIONS**.

When assessing an overdose, any additives in the solution must also be considered. The effects of an overdose may require immediate medical attention and treatment. Interventions include discontinuation of Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP administration, dose reduction, and other measures as indicated for the specific clinical constellation (e.g., monitoring of fluid balance, electrolyte concentrations and acid base balance).

DOSAGE AND ADMINISTRATION

Important Administration Instructions

- Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP is intended for intravenous administration using sterile equipment.
- Do not connect flexible plastic containers in series in order to avoid air embolism due to possible residual air contained in the primary container.
- Set the vent to the closed position on a vented intravenous administration set to prevent air embolism.
- Use a dedicated line without any connections to avoid air embolism.
- Do not pressurize intravenous solutions contained in flexible plastic containers to increase flow rates in order to avoid air embolism due to incomplete evacuation of residual air in the container.
- Prior to infusion, visually inspect the solution for particulate matter and discoloration. The solution should be clear and there should be no precipitates. Do not administer unless solution is clear, and container is undamaged.
- Do not administer Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP simultaneously with citrate anticoagulated/preserved blood through the same administration set because of the likelihood of coagulation precipitated by the calcium content in Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP.

Dosing Information

The choice of product, dosage, volume, rate, and duration of administration is dependent upon the age, weight and clinical condition of the patient and concomitant therapy, and administration should be determined by a physician experienced in intravenous fluid therapy.

Introduction of Additives

Evaluate all additions to the plastic container for compatibility and stability of the resulting preparation. Consult with a pharmacist, if available.

Additives may be incompatible. Complete information is not available. Those additives known to be incompatible should not be used. Consult with pharmacist, if available. If, in the informed judgment of the physician, it is deemed advisable to introduce additives, use aseptic technique. Mix thoroughly when additives have been introduced. Do not store solutions containing additives.

HOW SUPPLIED

Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP in VIAFLEX Plus plastic containers is available as shown below:

Code	Size (mL)	NDC	Product Name
2B2224	1000	0338-0811-04	20 mEq/L Potassium

Exposure of pharmaceutical products to heat should be minimized. Avoid excessive heat. It is recommended the product be stored at room temperature (25°C); brief exposure up to 40°C does not adversely affect the product.

DIRECTIONS FOR USE OF VIAFLEX PLUS PLASTIC CONTAINER

For Information on Risk of Air Embolism - see **DOSAGE AND ADMINISTRATION**

To Open

Tear overwrap down side at slit and remove solution container. Visually inspect the container. If the outlet port protector is damaged, detached, or not present, discard container as solution path sterility may be impaired. Some opacity of the plastic due to moisture absorption during the sterilization process may be observed. This is normal and does not affect the solution quality or safety. The opacity will diminish gradually. Check for minute leaks by squeezing inner bag firmly. If leaks are found, discard solution as sterility may be impaired. If supplemental medication is desired, follow directions below.

Preparation for Administration

1. Suspend container from eyelet support.
2. Remove protector from outlet port at bottom of container.
3. Attach administration set. Refer to complete directions accompanying set.

To Add Medication

To add medication before solution administration

1. Prepare medication site.
2. Using syringe with 19 to 22 gauge needle, puncture resealable medication port and inject.
3. Mix solution and medication thoroughly. For high density medication such as potassium chloride, squeeze ports while ports are upright and mix thoroughly.

To add medication during solution administration

1. Close clamp on the set.
2. Prepare medication site.
3. Using syringe with 19 to 22 gauge needle, puncture resealable medication port and inject.
4. Remove container from IV pole and/or turn to an upright position.
5. Evacuate both ports by squeezing them while container is in the upright position.
6. Mix solution and medication thoroughly.
7. Return container to in use position and continue administration.

Baxter Healthcare Corporation

Deerfield, IL 60015 USA

Printed in USA

07-19-00-1629

Rev. October 2019

Baxter and Viaflex are trademarks of Baxter International Inc.

PACKAGE LABEL - PRINCIPAL DISPLAY PANEL

LOT

EXP



2B2224
NDC 0338-0811-04
DIN 00786314

1

20 mEq Potassium Chloride

2

3

4

(20 mEq/L) Potassium Chloride in
Lactated Ringer's and 5% Dextrose
Injection USP

5

1000 mL

EACH 100 mL CONTAINS 5 g DEXTROSE HYDROUS USP 600 mg SODIUM CHLORIDE USP 310 mg SODIUM LACTATE 179 mg POTASSIUM CHLORIDE USP 20 mg CALCIUM CHLORIDE USP pH 5.0 (3.5 TO 6.5) mEq/L SODIUM 130 POTASSIUM 24 CALCIUM 3 CHLORIDE 129 LACTATE 28 HYPERTONIC OSMOLARITY 565 mOsmol/L (CALC) STERILE NONPYROGENIC SINGLE DOSE CONTAINER **NOT FOR USE IN THE TREATMENT OF LACTIC ACIDOSIS** ADDITIVES MAY BE INCOMPATIBLE CONSULT WITH PHARMACIST IF AVAILABLE WHEN INTRODUCING ADDITIVES USE ASEPTIC TECHNIQUE MIX THOROUGHLY DO NOT STORE DOSAGE INTRAVENOUSLY AS DIRECTED BY A PHYSICIAN SEE DIRECTIONS CAUTIONS SQUEEZE AND INSPECT INNER BAG WHICH MAINTAINS PRODUCT STERILITY DISCARD IF LEAKS ARE FOUND MUST NOT BE USED IN SERIES CONNECTIONS DO NOT ADMINISTER SIMULTANEOUSLY WITH BLOOD DO NOT USE UNLESS SOLUTION IS CLEAR Rx ONLY STORE UNIT IN MOISTURE BARRIER OVERWRAP AT ROOM TEMPERATURE (25°C/77°F) UNTIL READY TO USE AVOID EXCESSIVE HEAT SEE INSERT

6

7

8

VIAFLEX PLUS CONTAINER PL 146 PLASTIC
BAXTER VIAFLEX AND PL 146 ARE TRADEMARKS OF
BAXTER INTERNATIONAL INC
FOR PRODUCT INFORMATION 1-800-933-0303

Baxter

BAXTER HEALTHCARE CORPORATION
DEERFIELD IL 60015 USA
MADE IN USA

DISTRIBUTED IN CANADA BY
BAXTER CORPORATION
TORONTO ONTARIO CANADA

9

LOT

EXP

2B2224
NDC 0338-0811-04
DIN 00786314

20 mEq

Potassium Chloride

(20 mEq/L) Potassium Chloride in
Lactated Ringer's and 5% Dextrose
Injection USP

1000 mL

EACH 100 mL CONTAINS 5 g DEXTROSE HYDROUS USP 600 mg SODIUM CHLORIDE USP 310 mg SODIUM LACTATE 179 mg POTASSIUM CHLORIDE USP 20 mg CALCIUM CHLORIDE USP pH 5.0 (3.5 TO 6.5) mEq/L SODIUM 130 POTASSIUM 24

CALCIUM 3 CHLORIDE 129 LACTATE 28 HYPERTONIC
 OSMOLARITY 565 mOsmo/L (CALC) STERILE NONPYROGENIC
 SINGLE DOSE CONTAINER **NOT FOR USE IN THE TREATMENT OF
 LACTIC ACIDOSIS** ADDITIVES MAY BE INCOMPATIBLE CONSULT WITH
 PHARMACIST IF AVAILABLE WHEN INTRODUCING ADDITIVES USE
 ASEPTIC TECHNIQUE MIX THOROUGHLY DO NOT STORE DOSAGE
 INTRAVENOUSLY AS DIRECTED BY A PHYSICIAN SEE DIRECTIONS
 CAUTIONS SQUEEZE AND INSPECT INNER BAG WHICH MAINTAINS
 PRODUCT STERILITY DISCARD IF LEAKS ARE FOUND MUST NOT BE
 USED IN SERIES CONNECTIONS DO NOT ADMINISTER
 SIMULTANEOUSLY WITH BLOOD DO NOT USE UNLESS SOLUTION IS
 CLEAR Rx ONLY STORE UNIT IN MOISTURE BARRIER OVERWRAP AT
 ROOM TEMPERATURE (25°C/77°F) UNTIL READY TO USE AVOID
 EXCESSIVE HEAT SEE INSERT

VIAFLEX PLUS CONTAINER

PL 146 PLASTIC

BAXTER VIAFLEX AND PL 146 ARE TRADEMARKS OF
 BAXTER INTERNATIONAL INC

FOR PRODUCT INFORMATION 1-800-933-0303

Baxter

BAXTER HEALTHCARE CORPORATION
 DEERFIELD IL 60015 USA

MADE IN USA

DISTRIBUTED IN CANADA BY
BAXTER CORPORATION
 TORONTO ONTARIO CANADA

POTASSIUM CHLORIDE IN LACTATED RINGERS AND DEXTROSE

dextrose monohydrate, sodium chloride, sodium lactate, potassium chloride, calcium chloride
 injection, solution

Product Information				
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0338-0811	
Route of Administration	INTRAVENOUS			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
DEXTROSE MONOHYDRATE (UNII: LX22YL083G) (ANHYDROUS DEXTROSE - UNII:5SLOG7ROOK)		DEXTROSE MONOHYDRATE	5 g in 100 mL	
SODIUM CHLORIDE (UNII: 451V47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37, CHLORIDE ION - UNII:Q32ZN48698)		SODIUM CHLORIDE	600 mg in 100 mL	
SODIUM LACTATE (UNII: TU7HW0W0QT) (SODIUM CATION - UNII:LYR4M0NH37, LACTIC ACID - UNII:33X04XA5AT)		SODIUM LACTATE	310 mg in 100 mL	
POTASSIUM CHLORIDE (UNII: 660YQ98I10) (POTASSIUM CATION - UNII:295O53K152, CHLORIDE ION - UNII:Q32ZN48698)		POTASSIUM CHLORIDE	179 mg in 100 mL	
CALCIUM CHLORIDE (UNII: M4I0D6VV5M) (CALCIUM CATION - UNII:2M83C4R6ZB, CHLORIDE ION - UNII:Q32ZN48698)		CALCIUM CHLORIDE	20 mg in 100 mL	
Inactive Ingredients				
Ingredient Name		Strength		
WATER (UNII: 059QF0KO0R)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-0811-	1000 mL in 1 BAG; Type 0: Not a Combination	04/05/1995	

04	Product	04/05/1985	
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA019367	04/05/1985	

Labeler - Baxter Healthcare Corporation (005083209)

Establishment			
Name	Address	ID/FEI	Business Operations
Baxter Healthcare Corporation		059140764	ANALYSIS(0338-0811) , MANUFACTURE(0338-0811) , LABEL(0338-0811) , PACK(0338-0811) , STERILIZE(0338-0811) , API MANUFACTURE(0338-0811)

Establishment			
Name	Address	ID/FEI	Business Operations
Baxter Healthcare Corporation		194684502	ANALYSIS(0338-0811)

Establishment			
Name	Address	ID/FEI	Business Operations
Baxter Healthcare Corporation		189326168	ANALYSIS(0338-0811) , MANUFACTURE(0338-0811) , LABEL(0338-0811) , PACK(0338-0811) , STERILIZE(0338-0811)

Revised: 10/2019

Baxter Healthcare Corporation