

**MEDI-FIRST IBUPROFEN- ibuprofen tablet, coated**  
**MEDI-FIRST PLUS IBUPROFEN- ibuprofen tablet, coated**  
**MEDIQUE IPRIN- ibuprofen tablet, coated**  
**DOVER ADDAPRIN- ibuprofen tablet, coated**  
**OTIS CLAPP ULTRAPRIN- ibuprofen tablet, coated**  
**Unifirst First Aid Corporation**

-----

## **UniFirst First Aid Ibuprofen**

### ***Drug Facts***

#### ***Active ingredient***

Ibuprofen 200 mg (NSAID)

\*nonsteroidal antiinflammatory drug

#### ***Purpose***

Pain reliever/fever reducer

#### ***Uses***

Temporarily relieves minor aches and pains associated with

■ headache ■ toothache ■ backache ■ menstrual cramps

■ common cold ■ muscular aches ■ minor arthritis pain

Temporarily reduces fever.

#### ***Warnings***

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

■ hives ■ skin reddening ■ asthma (wheezing) ■ facial swelling ■ rash ■ shock ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

■ are age 60 or older

■ have had stomach ulcers or bleeding problems

■ take a blood thinning (anticoagulant) or steroid drug

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

■ have 3 or more alcoholic drinks every day while using this product

■ take more or for a longer time than directed

**Heart attack or stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more

than directed or for longer than directed.

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- you have problems or serious side effects from taking pain relievers or fever reducers
- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- redness or swelling is present in the painful area
- any new or unexpected symptoms occur

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

***Directions***

- **do not use more than directed**
- **the smallest effective dose should be used**
- do not take longer than 10 days, unless directed by a doctor (see Warnings)

**Adults and children:(12 years and older)**

Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used.

Do not exceed 6 tablets in 24 hours, unless directed by a doctor.

**Children under 12 years:**

Ask a doctor

***Other information***

- read all product information before using
- store at 68-77°F (20-25°C)
- avoid excessive heat 104°F (above 40°C )
- tamper evident sealed packets
- do not use any opened or torn packets

***Inactive ingredients***

carnauba wax\*, corn starch, hypromellose\*, iron oxide red, lactose\*, magnesium stearate\*, microcrystalline cellulose\*, polydextrose\*, polyethylene glycol, povidone (K30)\*, silicon dioxide, sodium starch glycolate, stearic acid, talc\*, titanium dioxide

*\*may contain*

***Questions or comments?*** 1-800-634-7680

**Medi-First Ibuprofen Label**

Medi-First®

Ibuprofen 200 mg

100 tablets (50 x 2)

Pain Reliever/Fever Reducer

Aches, Fever • Ibuprofen (NSAID) 200 mg

Pull to Open

Compare active ingredient to:

Advil®

Registered Trademark of Pfizer Consumer Healthcare

This Package is for Households without Young Children.

### Drug Facts

**Active ingredient (in each tablet)** Purpose  
 Ibuprofen 200 mg (NSAID) Pain reliever/fever reducer  
 "nonsteroidal anti-inflammatory drug"

**Uses**  
 Temporarily relieves minor aches and pains associated with  
 headaches toothache backache menstrual cramps  
 common cold muscular aches minor arthritis pain  
 temporarily reduces fever

**Warnings**  
 Allergic alert: Ibuprofen may cause a severe allergic reaction in people allergic to aspirin. Symptoms may include:  
 hives skin redness/itching asthma (wheezing)  
 facial swelling rash stomach cramps  
 If an allergic reaction occurs, stop use and seek medical help right away.  
 Research behind ibuprofen: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
 are age 60 or older  
 have had stomach ulcers or bleeding problems  
 take a blood thinner (anticoagulant) or osteoporosis medicine  
 take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)



**100 Tablets (50 x 2)**

### Drug Facts (continued)

have 3 or more alcoholic drinks every day while using this product  
 take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use:**  
 If you have ever had an allergic reaction to any other pain reliever/fever reducer  
 right before or after heart surgery

**Ask a doctor before if you:**  
 have problems or serious side effects from taking pain relievers  
 are taking aspirin  
 stomach bleeding warning applies to you  
 you have a history of stomach problems such as heartburn  
 you have high blood pressure, heart disease, liver or kidney disease, asthma, or have a stroke  
 you are taking a diuretic

**Ask a doctor or pharmacist before use if you:**  
 are taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin  
 under a doctor's care for any serious condition  
 taking any other drug (continued on opposite panel)

### Drug Facts (continued)

When using with or without food, if stomach upset occurs

**Stop use and ask a doctor if:**  
 you experience any of the following signs of stomach bleeding:  
 feel faint or vomit blood  
 have bloody or black stools  
 have stomach pain that does not get better  
 you have symptoms of heart problems or stroke  
 chest pain trouble breathing  
 weakness or lightheadedness in one part of body  
 blurred speech loss hearing

**Children under 12 years:**  
 pain lasts more or lasts for more than 10 days  
 fever gets worse or lasts for more than 3 days  
 redness or swelling is present in the painful area  
 any new or unexpected symptoms occur

**If pregnant or breast-feeding:** Ask a health professional before use. It is especially important not to use ibuprofen at 30 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.



### Drug Facts (continued)

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
 do not use more than directed  
 the smallest effective dose should be used  
 do not take longer than 10 days, unless directed by a doctor (see Warnings)

**Adults and children:** Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.

**Children under 12 years:**  
 Ask a doctor

**Other information**  
 Use all product information before using  
 store at 68-77°F (20-25°C)  
 avoid excessive heat (do not store above 70°C)  
 tamper evident unit-dose packets  
 do not use any opened or torn packets (continued on opposite panel)

**Retain carton for complete product information**

**Tamper Evident Unit Dose Packets**

**Retain carton for complete product information**

## Medi-First Plus Ibuprofen Label

Medi-First® Plus

# Ibuprofen

Ibuprofen 200 mg (NSAID)

100 Tablets (50 x 2's)

## Pull To Open

This Package is for Households without Young Children.

## Pain Reliever/Fever Reducer

Compare active ingredient to:

Advil®

Registered Trademark of Pfizer Consumer Healthcare

## Tamper Evident Unit Dose Packets

### Drug Facts

**Active ingredient (in each tablet)** **Purpose**  
 ibuprofen 200 mg (NSAID) Pain reliever/fever reducer  
 nonsteroidal anti-inflammatory drug

**Uses**  
 Temporarily relieves minor aches and pains associated with:  
 • headache • toothache • backache • menstrual cramps  
 • common cold • muscular aches • minor arthritis pain  
 Temporarily reduces fever.

**Warnings**  
**Allergy alert:** ibuprofen may cause a severe allergic reaction, especially in people taking aspirin. Symptoms may include:  
 • hives • skin redness • asthma (wheezing)  
 • facial swelling • rash • shock • blisters  
 If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
 • are age 60 or older  
 • have had stomach ulcers or bleeding problems  
 • take a blood thinning (anticoagulant) or steroid drug  
 • take other stomach pain prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

40 Tablets  
(50 x 2's)

MEDI FIRST  
*Plus*

# Ibuprofen

**Ibuprofen 200 mg (NSAID)**

*Push To Open*

60 Tablets  
(30 x 2's)

**THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.**

**Pain Reliever/Fever Reducer**

### Drug Facts (continued)

• have 3 or more alcoholic drinks every day while using this product  
 • take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These risks can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use:**  
 • if you have ever had an allergic reaction to any other pain reliever/fever reducer  
 • right before or after heart surgery

**Ask a doctor before use if:**  
 • you have had problems or serious side effects from taking pain relievers or fever reducers  
 • you are on blood-thinning or aspirin therapy  
 • you have a history of stomach problems such as heartburn  
 • you have high blood pressure, heart disease, liver, stomach, or kidney problems, asthma or have taken a diuretic  
 • you are taking a diuretic

**Ask a doctor or pharmacist before use if you are:**  
 • taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin  
 • under a doctor's care for any serious condition  
 • taking any other drug (continued on opposite panel)

### Drug Facts (continued)

**When using this product:**  
 • take with food or a full stomach upon occurs

**Stop use and ask a doctor if:**  
 • you experience any of the following signs of stomach bleeding:  
   • black, tarry, or vomit blood  
   • have bloody or black stools  
   • have stomach pain that does not get better  
 • you have symptoms of heart problems or stroke  
   • chest pain • trouble breathing  
   • weakness in one part or side of body  
   • slurred speech • tingling swelling  
   • lower gums worse or lasts for more than 10 days  
   • fever, pain worse or lasts for more than 3 days  
   • redness or swelling is present in the painful area  
   • you have new or unexpected symptoms

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

40 Tablets  
(50 x 2's)

MEDI FIRST  
*Plus*

# Ibuprofen

**Ibuprofen 200 mg (NSAID)**

*Push To Open*

60 Tablets  
(30 x 2's)

**THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.**

**Pain Reliever/Fever Reducer**

### Drug Facts (continued)

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
 • do not use more than directed  
 • the smallest effective dose should be used  
 • do not take longer than 10 days, unless directed by a doctor (see Warnings)

**Adults and children (12 years and older)**  
 Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours. Adults, unless directed by a doctor.

**Children under 12 years:**  
 Ask a doctor

**Other information**  
 • read all product information before using  
 • use with food or a full stomach  
 • avoid excessive heat 104°F (above 40°C)  
 • larger infant sealed packets  
 • do not use any opened or torn packets  
 (continued on opposite panel)

### Drug Facts (continued)

**Inactive ingredients**  
 carboxylic acid, calcium stearate, hypromellose, iron oxide red, lactose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, polyethylene glycol 400, sodium chloride, sodium starch glycolate, stearic acid, talc, titanium dioxide

**Questions or comments?** 1-800-634-7680

40 Tablets  
(50 x 2's)

MEDI FIRST  
*Plus*

# Ibuprofen

**Ibuprofen 200 mg (NSAID)**

*Push To Open*

60 Tablets  
(30 x 2's)

**THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.**

**Pain Reliever/Fever Reducer**

### Drug Facts (continued)

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
 • do not use more than directed  
 • the smallest effective dose should be used  
 • do not take longer than 10 days, unless directed by a doctor (see Warnings)

**Adults and children (12 years and older)**  
 Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours. Adults, unless directed by a doctor.

**Children under 12 years:**  
 Ask a doctor

**Other information**  
 • read all product information before using  
 • use with food or a full stomach  
 • avoid excessive heat 104°F (above 40°C)  
 • larger infant sealed packets  
 • do not use any opened or torn packets  
 (continued on opposite panel)

### Drug Facts (continued)

**Inactive ingredients**  
 carboxylic acid, calcium stearate, hypromellose, iron oxide red, lactose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, polyethylene glycol 400, sodium chloride, sodium starch glycolate, stearic acid, talc, titanium dioxide

**Questions or comments?** 1-800-634-7680

40 Tablets  
(50 x 2's)

MEDI FIRST  
*Plus*

# Ibuprofen

**Ibuprofen 200 mg (NSAID)**

*Push To Open*

60 Tablets  
(30 x 2's)

**THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.**

**Pain Reliever/Fever Reducer**

### Drug Facts (continued)

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
 • do not use more than directed  
 • the smallest effective dose should be used  
 • do not take longer than 10 days, unless directed by a doctor (see Warnings)

**Adults and children (12 years and older)**  
 Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours. Adults, unless directed by a doctor.

**Children under 12 years:**  
 Ask a doctor

**Other information**  
 • read all product information before using  
 • use with food or a full stomach  
 • avoid excessive heat 104°F (above 40°C)  
 • larger infant sealed packets  
 • do not use any opened or torn packets  
 (continued on opposite panel)

### Drug Facts (continued)

**Inactive ingredients**  
 carboxylic acid, calcium stearate, hypromellose, iron oxide red, lactose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, polyethylene glycol 400, sodium chloride, sodium starch glycolate, stearic acid, talc, titanium dioxide

**Questions or comments?** 1-800-634-7680

40 Tablets  
(50 x 2's)

MEDI FIRST  
*Plus*

# Ibuprofen

**Ibuprofen 200 mg (NSAID)**

*Push To Open*

60 Tablets  
(30 x 2's)

**THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.**

**Pain Reliever/Fever Reducer**

### Drug Facts (continued)

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
 • do not use more than directed  
 • the smallest effective dose should be used  
 • do not take longer than 10 days, unless directed by a doctor (see Warnings)

**Adults and children (12 years and older)**  
 Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours. Adults, unless directed by a doctor.

**Children under 12 years:**  
 Ask a doctor

**Other information**  
 • read all product information before using  
 • use with food or a full stomach  
 • avoid excessive heat 104°F (above 40°C)  
 • larger infant sealed packets  
 • do not use any opened or torn packets  
 (continued on opposite panel)

### Drug Facts (continued)

**Inactive**

## Medique Iprin Label

Medique®

I-Prin

Ibuprofen 200 mg

Anti-Inflammatory (NSAID)

This Package is for Households without Young Children.

Pain Reliever/Fever Reducer • Ibuprofen 200 mg

24 Tablets (12 x 2)

Tamper Evident Unit Dose Packets.



Dover Addaprin Label

Dover Addaprin™

Ibuprofen 200 mg Tablets (NSAID)

Pain Reliever-Fever Reducer

This Package is for Households Without Young Children.

Dover Pharmaceutical

Products of the highest quality and effectiveness


Tamper Evident

Sealed Packets

Unit Dose Packs

500 Tablets

(250 Packets of 2)



Otis Clapp Ultraprin Label

500 Tablets (250 Packets of 2)

<div><div>MEDI-FIRST IBUPROFEN</div><div>ibuprofen tablet, coated</div></div>			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-718
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)		IBUPROFEN	200 mg
Inactive Ingredients			
Ingredient Name			Strength
STARCH, CORN (UNII: O8232NY3SJ)			
POVIDONE K30 (UNII: U725QWY32X)			

<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	

### Product Characteristics

<b>Color</b>	red (Reddish Brown)	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	G;2
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-718-13	250 in 1 BOX	01/26/2017	
1	NDC:47682-718-99	2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-718-48	125 in 1 BOX	01/26/2017	
2		2 in 1 PACKET; Type 0: Not a Combination Product		
3	NDC:47682-718-33	50 in 1 BOX	01/26/2017	
3		2 in 1 PACKET; Type 0: Not a Combination Product		
4	NDC:47682-718-30	4 in 1 BOX	01/26/2017	
4		2 in 1 PACKET; Type 0: Not a Combination Product		
5	NDC:47682-718-50	25 in 1 BOX	04/16/2019	
5		2 in 1 PACKET; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079174	01/26/2017	

## MEDI-FIRST PLUS IBUPROFEN

ibuprofen tablet, coated

### Product Information



<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:47682-709
<b>Route of Administration</b>	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
FERRIC OXIDE RED (UNII: 1K09F3G675)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
POVIDONE K30 (UNII: U725QWY32X)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
STARCH, CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	

Product Characteristics

<b>Color</b>	red (Reddish Brown)	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	G;2
<b>Contains</b>			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-709-48	125 in 1 BOX	01/26/2017	
1		2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-709-33	50 in 1 BOX	01/26/2017	
2		2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079174	01/26/2017	



ibuprofen tablet, coated

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:47682-700
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>IBUPROFEN</b> (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	

## Product Characteristics

<b>Color</b>	red (Reddish Brown)	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	G;2
<b>Contains</b>			

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:47682-700-69	3 in 1 BOX	01/26/2017	
1	NDC:47682-700-99	2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-700-64	12 in 1 BOX	01/26/2017	
2		2 in 1 PACKET; Type 0: Not a Combination Product		
3	NDC:47682-700-47	100 in 1 BOX	01/26/2017	
3		2 in 1 PACKET; Type 0: Not a Combination Product		
4	NDC:47682-700-13	250 in 1 BOX	01/26/2017	
4		2 in 1 PACKET; Type 0: Not a Combination		

+	Product		
<b>Marketing Information</b>			
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANDA	ANDA079174	01/26/2017	

<b>DOVER ADDAPRIN</b>			
ibuprofen tablet, coated			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:47682-714
<b>Route of Administration</b>	ORAL		
<b>Active Ingredient/Active Moiety</b>			
<b>Ingredient Name</b>		<b>Basis of Strength</b>	<b>Strength</b>
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)		IBUPROFEN	200 mg
<b>Inactive Ingredients</b>			
<b>Ingredient Name</b>			<b>Strength</b>
STARCH, CORN (UNII: O8232NY3SJ)			
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)			
FERRIC OXIDE RED (UNII: 1K09F3G675)			
STEARIC ACID (UNII: 4ELV7Z65AP)			
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
POVIDONE K30 (UNII: U725QWY32X)			
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)			
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)			
TALC (UNII: 7SEV7J4R1U)			
<b>Product Characteristics</b>			
<b>Color</b>	red (Reddish Brown)	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	G;2
<b>Contains</b>			
<b>Packaging</b>			
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>
			<b>Marketing End Date</b>
1	NDC:47682-714-13	250 in 1 BOX	01/26/2017

1	NDC:47682-714-99	2 in 1 PACKET; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
ANDA	ANDA079174		01/26/2017	

OTIS CLAPP ULTRAPRIN				
ibuprofen tablet, coated				
Product Information				
Product Type		HUMAN OTC DRUG	Item Code (Source)	
Route of Administration		ORAL	NDC:47682-702	
Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)			IBUPROFEN	200 mg
Inactive Ingredients				
Ingredient Name				Strength
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
TALC (UNII: 7SEV7J4R1U)				
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
STEARIC ACID (UNII: 4ELV7Z65AP)				
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)				
POVIDONE K30 (UNII: U725QWY32X)				
STARCH, CORN (UNII: O8232NY3SJ)				
FERRIC OXIDE RED (UNII: 1K09F3G675)				
Product Characteristics				
Color	red (Reddish Brown)		Score	no score
Shape	ROUND		Size	10mm
Flavor			Imprint Code	G;2
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	NDC:47682-702			

1	NDC:47682-702-13	250 in 1 BOX	02/01/2017	04/03/2017
1	NDC:47682-702-99	2 in 1 PACKET; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
ANDA	ANDA079174		11/17/2014	04/03/2017

**Labeler** - Unifirst First Aid Corporation (832947092)