

**BAYER CHEWABLE - ASPIRIN REGIMEN LOW DOSE ASPIRIN CHERRY  
FLAVORED- aspirin tablet, chewable  
Bayer HealthCare LLC.**

-----  
**Bayer Chewable - Aspirin Regimen**

***Drug Facts***

Aspirin 81 mg (NSAID) <sup>1</sup>

---

1 nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever

**Directions**

• drink a full glass of water with each dose • adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor • children under 12 years: consult a doctor

**Uses**

- for the temporary relief of minor aches and pains or as recommended by your doctor
- ask your doctor about other uses for Bayer 81 mg Cherry Chewable Aspirin

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives ● facial swelling ● asthma (wheezing) ● shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs

(aspirin, ibuprofen, naproxen, or others)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### **Do not use**

#### **Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- this product for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

### **Ask a doctor or pharmacist**

**Ask a doctor or pharmacist before use if you are** taking a prescription drug for

- gout ● diabetes ● arthritis

### **Stop use**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint ● vomit blood ● have bloody or black stools
  - have stomach pain that does not get better
  - pain gets worse or lasts more than 10 days
  - redness or swelling is present
  - new symptoms occur
  - ringing in the ears or a loss of hearing occurs

## **If pregnant or breast-feeding**

**If pregnant or breast-feeding**, ask a health professional before use . **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

## **Keep out of reach of children**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

## **Directions**

### ***Directions***

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

## **Other information**

### ***Other information***

- save carton for full directions and warnings
- store at room temperature

## **Inactive ingredients**

***Inactive ingredients*** colloidal silicon dioxide, corn starch, D&C red #27 aluminum lake, dextrose, FD&C red #40 aluminum lake, flavor, microcrystalline cellulose, saccharin sodium

## **Questions or comments**

***Questions or comments?*** **1-800-331-4536** (Mon-Fri 9AM – 5PM EST)

**36 count carton**

5/8"

LOT/EXP text is to be preprinted. No restriction on font size, style or color.

LOT & EXP

LOT/EXP


3

F.P.O. 90% UPC TRUNCATED

1284313231

3

Do not use this product if printed safety seal bearing Bayer HealthCare® vendor can is torn or missing.



PS

**Drug Facts (continued)**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - redness or swelling is present
  - new symptoms occur
  - ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 chewable tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

**Drug Facts (continued)**

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- this product for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if ● stomach bleeding warning applies to you ● you have a history of stomach problems, such as heartburn ● you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ● you are taking a diuretic ● you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if ● an allergic reaction occurs. Seek medical help right away.

**Drug Facts (continued)**

**Other information**

- save carton for full directions and warnings
- store at room temperature. Avoid excessive heat above 40°C (104°F).

**Inactive ingredients:** cobalt silicon dioxide, corn starch, D&C red #27 aluminum lake, dextrose, FMC red #40 aluminum lake, flavor, microcrystalline cellulose, saccharin sodium

**Questions? 1-800-331-4558**

Made in Spain  
Bayer HealthCare LLC  
Whisper, IN 47981  
Bayer and the Bayer Cross are registered trademarks of Bayer.

3494  
85987386

**Drug Facts**

**Active ingredient Purpose (in each chewable tablet)**  
Aspirin (ASA) (C17H15O9) Pain reliever, nonsteroidal anti-inflammatory drug

**Uses**

- for the temporary relief of minor aches and pains or as recommended by your doctor
- ask your doctor about other uses for Bayer 81 mg Cherry Chewable Aspirin

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if change in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
  - are age 60 or older


Aspirin is not appropriate for everyone, so be sure to talk to your doctor before you begin an aspirin regimen.

FPO

3494

12 OF 5  
CODE AREA  
REQUIRES 1/4" QUIET ZONE

CLOSE ANGLE WAVE RULE



N 2080-2090-36

ASPIRIN REGIMEN ASPIRIN (NSAID) PAIN RELIEVER


**BAYER**

**CHEWABLE**

**81mg**

**36 TABLETS**

Low Dose Aspirin CHERRY FLAVORED



**BAYER CHEWABLE - ASPIRIN REGIMEN LOW DOSE ASPIRIN CHERRY FLAVORED**

aspirin tablet, chewable

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0280-2090
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

### Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>ALUMINUM OXIDE</b> (UNII: LMI26O6933)	
<b>D&amp;C RED NO. 27</b> (UNII: 2LRS185U6K)	
<b>DEXTROSE</b> (UNII: IY9XDZ35W2)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	

### Product Characteristics

<b>Color</b>	pink	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	5mm
<b>Flavor</b>	CHERRY	<b>Imprint Code</b>	BAYER
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0280-2090-36	1 in 1 CARTON	07/02/2002	
1		36 in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M013	07/02/2002	

**Labeler** - Bayer HealthCare LLC. (112117283)

## Establishment

Name	Address	ID/FEI	Business Operations
Bayer HealthCare LLC Consumer Care		072827066	pack(0280-2090) , manufacture(0280-2090)

Revised: 11/2023

Bayer HealthCare LLC.