

YINCHIAO- acetaminophen and chlorpheniramine maleate tablet, coated
ANHUI DONGSHENGYOUBANG PHARMACEUTICAL CO., LTD.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

DRUG FACTS

Active ingredient (in each tablet)

Acetaminophen 60 mg

Chlorpheniramine maleate 0.667 mg

Purpose

Pain reliever-fever reducer

Antihistamine

Uses

Temporarily relieves the following symptoms associated with the common cold, hay fever, or other upper respiratory allergies:

minor aches and pains

headache

muscular aches

fever

sore throat

sneezing

itching of the nose or throat

runny nose

itchy, watery eyes

Warnings

Do not use

for pain for more than 10 days

for fever for more than 3 days

Ask a doctor before use if the user has

glaucoma

a breathing problem such as emphysema or chronic bronchitis

difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers

When using this product

may cause excitability especially in children

may cause drowsiness (alcohol, sedatives, and tranquilizers may increase the drowsiness effect)

avoid alcoholic beverages

use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if
pain or fever persists or worsen
new symptoms occur
redness or swelling is present
sore throat:
is severe
last for more than 2 days
is accompanied or followed by:
fever
headache
rash
nausea
vomiting

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

adults and children 12 years of age and older: Take 6 tablets every 4 hours, not more than 36 tablets in 24 hours
children 6 to under 12 years of age: Take 3 tablets every 4 hours, not more than 18 tablets in 24 hours
children under 6 years: consult a doctor

Other information

protect from light and excessive heat
keep tightly closed
keep in dry place

Questions or Comments? (888) 221-3496 M-F 9 am to 5 pm

you may also use this number to report serious adverse events associated with the use of this product

Inactive ingredients

Japanese honeysuckle flower, forsythia fruit, balloon flower root, burdock fruit, Chinese licorice root, corn starch, dextrin, FD&C yellow no.5 (tartrazine), FD&C yellow no.6, lophatherum herb, magnesium stearate, mentha herb, prepared soybean seed, schizonepeta herb, sucrose, and talc.

YINCHIAO TABLET, NDC 72030-001-01, Antihistamine, Pain Reliever-Fever Reducer, 120 Tablets

銀翹解毒片 (西藥糖衣片) 120片盒稿
Yinchiao Tablet Antihistamine - Box
01/17/2018



NDC 72030-001-01

A GOOD REMEDY FOR COLDS

SUPERIOR

**YINCHIAO
TABLET**

Antihistamine
Pain Reliever-Fever Reducer



120 TABLETS

MANUFACTURED BY:
安徽東盛友邦製藥有限公司
ANHUI DONGSHENGYOUBANG
PHARMACEUTICAL CO., LTD.
ANHUI PROVINCE, CHINA

Drug Facts 藥品資料

Active Ingredients (in each tablet) Purpose 有效成份 (每片含量) 用途
Acetaminophen 60mg.....Pain Reliever-Fever Reducer
醋氯酚芬 60毫克.....解痛、退熱
Chlorpheniramine Maleate 0.667mg.....Antihistamine
撲爾敏0.667毫克.....抗組織胺

Uses 作用
Temporarily relieves the following symptoms associated with the common cold, hay fever, or other upper respiratory allergies: 暫時緩解傷風、花粉熱或其它上呼吸道感染引致的下列症狀:
■ minor aches and pains 輕度酸痛 ■ headache 頭痛
■ muscular aches 肌肉酸痛 ■ fever 發熱
■ sore throat 喉痛 ■ sneezing 噴嚏
■ itching of the nose or throat 鼻或喉發癢
■ runny nose 流涕 ■ itchy, watery eyes 發癢、流淚

Warnings 告誡
Do not use 在下列情況下請勿服用
■ for pain for more than 10 days 疼痛十天以上
■ for fever for more than 3 days 發熱三天以上
Ask a doctor before use if you have 如有下列症狀，在服用本品前請教醫師
■ glaucoma 青光眼
■ a breathing problem such as emphysema or chronic bronchitis 肺氣腫或慢性氣管炎等呼吸系統症狀
■ difficulty in urination due to enlargement of the prostate gland 前列腺肥大引起排尿困難
Ask a doctor or pharmacist before use if you are 如您目前正在服用下列藥物，在服用本品前請教醫師或藥劑師
■ taking sedatives or tranquilizers 安眠藥或鎮定劑
When using this product 在服用本品期間
■ may cause excitability, especially in children 可能會引起興奮，尤其對兒童。
■ may cause drowsiness (alcohol, sedatives, and tranquilizers may increase the drowsiness effect) 可能會引起倦睡感 (酒精、安眠藥或鎮定劑會加強倦睡感)
■ avoid alcoholic beverages 請避免飲用酒精性飲料。
■ use caution when driving a motor vehicle or operating machinery 駕駛車輛或操作機械器時要小心。
Stop use and ask a doctor if 如有下列症狀請停止服用本品並請醫師
■ pain or fever persist or worsen 疼痛或發熱持續不癒或惡化
■ new symptoms occur 出現新症狀
■ redness or swelling is present 出現紅腫
■ sore throat 喉痛
■ is severe 嚴重 ■ lasts for more than 2 days 兩天以上
■ is accompanied or followed by: 伴有或隨後出現
■ fever 發熱 ■ headache 頭痛
■ rash 疹子 ■ nausea 噁心 ■ vomiting 嘔吐

NDC 72030-001-01

感冒良藥

精製

銀翹

解毒片

治療傷風感冒良藥



120 片

U.S. DISTRIBUTOR:
SOLSTICE MEDICINE COMPANY
美國蘇氏藥業有限公司
215 W. Ann St., Los Angeles, CA 90012
Toll Free: 1-888-221-3496
www.solsticemed.com

Drug Facts (continued) 藥品資料 (續)

Warnings (continued) 告誡 (續)
If pregnant or breast-feeding, ask a health professional before use. 如在懷孕或哺乳期間，在服用本品前請教醫務人員。
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. 將本藥放在兒童不可接近處。如不慎服用過量，應立即尋求醫療護理或與毒物控制中心(Poison Control Center)聯絡。即使您沒有察覺任何跡象或症狀，及時取得醫療護理對兒童和成年人都是非常重要的。

Directions 服用方法

Adults and children 12 years & older 成年人及12歲或以上兒童	Take 6 tablets every 4 hours. Not more than 36 tablets in 24 hours 每4小時服6片，每24小時時段內不得服用超過36片
Children 6 to under 12 years 6歲至12歲以下兒童	Take 3 tablets every 4 hours. Not more than 18 tablets in 24 hours 每4小時服3片，每24小時時段內不得服用超過18片
Children under 6 years 6歲以下兒童	Consult a doctor 請教醫師

Other information 其它注意事項
■ Protect from light and excessive heat 避免光及高熱
■ Keep tightly closed 關閉緊蓋
■ Keep in dry place 儲放於乾燥處

Inactive ingredients 輔助成份
Japanese honeysuckle flower, korymbia fruit, balloon flower root, burdock fruit, Chinese licorice root, corn starch, dextrin, FD & C yellow no.5 (tartrazine), FD & C yellow no.6, lophatherum herb, magnesium stearate, mentha herb, prepared soybean seed, schizonepeta herb, sucrose, and talc.
金銀花、連翹、桔梗、牛蒡子、甘草、玉米澱粉、糊精、FD & C 5號黃食物色素(酒石黃)、FD & C 6號黃食物色素、忍冬藤、薄荷葉、硬脂酸鎂、薄荷、法半夏、茯苓、蔗糖及滑石。

Questions or Comments 問題或建議? (888) 221-3496
M-F, 9 am to 5 pm 週一至週五，上午9時至下午5時
You may also report serious side effects to this phone number.
您也可以使用這個電話號碼報告嚴重的副作用。

Made in China 中國製造

Exp. Date:

Lot No. :



0 49987 01521 8

YINCHIAO

acetaminophen and chlorpheniramine maleate tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72030-001
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	60 mg
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	0.667 mg

Inactive Ingredients

Ingredient Name	Strength
LONICERA JAPONICA FLOWER (UNII: 4465L2WS4Y)	
FORSYTHIA SUSPENS A FRUIT (UNII: P4793M1E55)	
PLATYCODON GRANDIFLORUS ROOT (UNII: 2DF0NS0O2Z)	
ARCTIUM LAPP A FRUIT (UNII: EA541308MV)	
GLYCYRRHIZA URALENSIS WHOLE (UNII: 8XW1DS8UIR)	
STARCH, CORN (UNII: O8232NY3SJ)	
DEXTRIN PALMITATE (CORN; 20000 MW) (UNII: 89B2BSF9I3)	
FD&C YELLOW NO. 5 (UNII: I753WB2F1M)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
LOPHATHERUM GRACILE WHOLE (UNII: 55L A08VLS9)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MINT (UNII: FV98Z8G1TP)	
SOYBEAN (UNII: L7HT8FIZOD)	
SCHIZONEPETA TENUFOLIA SPIKE (UNII: 2FN3BA1MZE)	
SUCROSE (UNII: C151H8M554)	
TALC (UNII: 7SEV7J4R1U)	

Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	YC
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72030-001-01	120 in 1 BOTTLE; Type 0: Not a Combination Product	01/18/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	01/18/2018	

Labeler - ANHUI DONGSHENGYOUBANG PHARMACEUTICAL CO., LTD. (527929527)

Establishment

Name	Address	ID/FEI	Business Operations
ANHUI DONGSHENGYOUBANG PHARMACEUTICAL CO., LTD.		527929527	manufacture(72030-001)

Revised: 1/2018

ANHUI DONGSHENGYOUBANG PHARMACEUTICAL CO., LTD.