

**CLEARLAX- polyethylene glycol 3350 powder, for solution  
H E B**

-----  
**HEB ClearLax Drug Facts**

**Active ingredient (in each dose)**

Polyethylene Glycol 3350, 17 g (cap filled to line)

**Purpose**

Osmotic Laxative

**Use**

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

**Warnings**

**Allergy alert:** Do not use if you are allergic to polyethylene glycol

**Do not use**

if you have kidney disease, except under the advice and supervision of a doctor

**Ask a doctor before use if you have**

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug

**When using this product**

you may have loose, watery, more frequent stools

**Stop use and ask a doctor if**

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

**If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### **Directions**

- **do not take more than directed unless advised by your doctor**
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
- adults and children 17 years of age and older:
  - use once a day
  - fill to top of white section in cap which is marked to indicate the correct dose (17 g)
  - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
  - do not combine with starch-based thickeners used for difficulty swallowing
  - ensure that the powder is fully dissolved before drinking
  - do not drink if there are any clumps
  - do not use more than 7 days
- children 16 years of age or under: ask a doctor

### **Other information**

- store at 20°-25°C (68°-77°F)
- tamper -evident: do not use if printed foil seal under cap is missing, open or broken

### **Inactive ingredients**

natural and artificial orange flavor, sucralose

### **Questions or comments?**

**1-800-719-9260**

### **Package/Label Principal Display Panel**

Compare to MiraLax<sup>®</sup> active ingredient

H•E•B<sup>®</sup>

ClearLax<sup>®</sup>

Polyethylene Glycol 3350 Powder for Solution

Osmotic Laxative

- Relieves Occasional Constipation/Irregularity
- Softens Stool

- SUGAR-FREE

- DYE-FREE

Orange Flavored

Grit Free Powder

NET WT. 17.9 OZ (510 g)

30 ONCE-DAILY DOSE

Compare to MiraLAX®  
active ingredient\*

NDC 59640-797-03



# ClearLax®

Polyethylene Glycol 3350  
Powder for Solution

## Osmotic Laxative

- Relieves Occasional Constipation/  
Irregularity
- Softens Stool
- SUGAR-FREE
- DYE-FREE

Orange Flavored  
Grit Free Powder



**NET WT. 17.9 OZ (510 g)**  
**30 ONCE-DAILY DOSES**

: 7U9D8 1J F1

Polyethylene Glycol 3350 increases  
frequency of bowel movements and softens the stool.

Dissolves in Any Beverage

### Drug Facts

<b>Active ingredient (in each dose)</b>	<b>Purpose</b>
Polyethylene Glycol 3350, 17 g (cap filled to line).....	Osmotic Laxative

**Use** ■ relieves occasional constipation (irregularity)  
■ generally produces a bowel movement in 1 to 3 days

### Warnings

**Allergy alert:** Do not use if you are allergic to polyethylene glycol

Do not use if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have ■ nausea, vomiting or abdominal pain ■ a sudden change in bowel habits that lasts over 2 weeks ■ irritable bowel syndrome

Ask a doctor or pharmacist before use if you are taking a prescription drug

When using this product you may have loose, watery, more frequent stools

Stop use and ask a doctor if ■ you have rectal bleeding or your anus bleeds, or you have abdominal pain that worsens

### Drug Facts (continued)

■ adults and children 17 years of age and older: ■ use once a day ■ fill to top of white section in cap which is marked to indicate the correct dose (17 g) ■ stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink ■ do not combine with starch-based thickeners used for difficulty swallowing ■ ensure that the powder is fully dissolved before drinking ■ do not drink if there are any clumps ■ do not use more than 7 days ■ children 16 years of age or under: ask a doctor

**Other information** ■ store at 20°-25°C (68°-77°F)

■ tamper-evident: do not use if printed foil seal under cap is missing, open or broken

**Inactive ingredients** natural and artificial orange flavor, sucralose

**Questions or comments?**  
1-800-719-9260



your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition. ■ you get diarrhea ■ you need to use a laxative for longer than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions** ■ do not take more than directed unless advised by your doctor ■ the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)

This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of MiraLAX®.

Patent Pending

**GLUTEN FREE**

MADE WITH PRIDE & CARE FOR H-E-B® SAN ANTONIO, TX 78204



29952 - 2210



: 70908 1J 81

## CLEARLAX

polyethylene glycol 3350 powder, for solution

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:59640-797
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>POLYETHYLENE GLYCOL 3350</b> (UNII: G2M7P15E5P) (POLYETHYLENE GLYCOL 3350 - UNII:G2M7P15E5P)	POLYETHYLENE GLYCOL 3350	17 g in 17 g

### Inactive Ingredients

Ingredient Name	Strength
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	

### Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	ORANGE	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59640-797-02	238 g in 1 BOTTLE; Type 0: Not a Combination Product	01/17/2023	
2	NDC:59640-797-03	510 g in 1 BOTTLE; Type 0: Not a Combination Product	01/17/2023	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090685	01/17/2023	

**Labeler -** H E B (007924756)

Revised: 1/2023

H E B