

**MEIJER STOMACH RELIEF REGULAR STRENGTH- bismuth  
subsalicylate suspension  
MEIJER DISTRIBUTION INC**

-----  
**Meijer Stomach Relief Ultra Strength Bismuth Subsalicylate 525 mg Drug  
Facts**

**Active ingredient (in each 15 mL dose)**

Bismuth subsalicylate 525 mg

**Purposes**

Upset stomach reliever and antidiarrheal

**Uses**

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
- heartburn
- indigestion
- nausea
- gas
- belching

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea or vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert: Contains salicylate. Do not take if you are**

- allergic to salicylates (including aspirin)
- taking other salicylate products

**Do not use if you have**

- an ulcer
- a bleeding problem
- bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool

### **Ask a doctor or pharmacist before use if you are**

taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

### **When using this product**

a temporary, but harmless, darkening of the stool and/or tongue may occur

### **Stop use and ask a doctor if**

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### **Directions**

- shake well before use
- only use dose cup provided
- **adults and children 12 years and over:**
- 15 mL (1 dose) every ½ hour or 30 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea
- 15 mL (1 dose) every ½ hour as needed overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (120 mL) in 24 hours
- use until diarrhea stops but not more than 2 days
- **children under 12 years:** ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

### **Other information**

- **each 15 mL contains:** sodium 10 mg
- **salicylate 230 mg**
- low sodium
- sugar free

- store at room temperature
- protect from freezing
- avoid excessive heat (over 104°F or 40°C)

### **Inactive ingredients**

carboxymethylcellulose sodium, D&C Red No. 22, D&C Red No. 28, flavor, microcrystalline cellulose, potassium hydroxide, potassium sorbate, purified water, salicylic acid, simethicone emulsion, sodium benzoate, sucralose, xanthan gum

### **Questions or comments?**

**1-866-467-2748**

**Do not use if imprinted shrink band is missing or broken.**

### **Principal Display Panel**

meijer®

NDC# **41250-211-12**

**Compare To Pepto-Bismol® Ultra Strength active ingredient\***

Stomach Relief

Bismuth Subsalicylate 525 mg

Ultra Strength

Original Flavor

Naturally and Artificially Flavored

Provides soothing relief for:

Nausea, Heartburn, Indigestion, Upset Stomach, Diarrhea

12 FL. OZ. (354 mL)

OUR QUALITY GUARANTEE

WWW.MEIJER.COM/SATISFACTION

**DIST. BY MEIJER DISTRIBUTION, INC.**

**GRAND RAPIDS, MI 49544**

**www.meijer.com**

**\*This product is not manufactured or distributed by The Procter & Gamble Co., owner of the registered trademark Pepto-Bismol®.**

**Drug Facts (continued)**

**Directions** ■ shake well before use ■ only use dose cup provided  
 ■ **adults and children 12 years and over** ■ 15 mL (1 dose) every ½ hour or 30mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea. ■ 15 mL (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)  
 ■ do not exceed 8 doses (120 mL) in 24 hours ■ use until diarrhea stops but not more than 2 days ■ **children under 12 years:** ask a doctor ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Other information**  
 ■ each 15 mL contains: sodium 10 mg  
 ■ salicylate 230 mg ■ low sodium ■ sugar free  
 ■ store at room temperature  
 ■ protect from freezing  
 ■ avoid excessive heat (over 104°F or 40°C)

**Inactive ingredients** carboxymethylcellulose sodium, D&C Red No. 22, D&C Red No. 28, flavor, microcrystalline cellulose, potassium hydroxide, potassium sorbate, purified water, salicylic acid, simethicone emulsion, sodium benzoate, sucralose, xanthan gum

**Questions or comments?** 1-866-467-2748

Compare to Pepto-Bismol®  
Ultra Strength  
active ingredient\*

# stomach relief

Bismuth Subsalicylate 525 mg

**Ultra Strength**

**Original Flavor**  
Naturally and Artificially Flavored

Provides soothing relief for:  
Nausea, Heartburn, Indigestion  
Upset Stomach, Diarrhea

## 12 FL. OZ. (354 mL)

**Do not use if imprinted shrink band is missing or broken**

**Drug Facts**

**Active ingredient** **Purpose**  
*(in each 15 mL dose)*  
 Bismuth subsalicylate 525 mg.....Upset stomach reliever and anti-diarrheal

**Uses** relieves ■ travelers' diarrhea ■ diarrhea ■ upset stomach due to overindulgence in food and drink, including:  
 ■ heartburn ■ indigestion ■ nausea ■ gas ■ belching ■ fullness

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert: Contains salicylate. Do not take if you are**  
 ■ allergic to salicylates (including aspirin)  
 ■ taking other salicylate products

**Do not use if you have** ■ an ulcer ■ a bleeding problem ■ bloody or black stool

**Ask a doctor before use if you have** ■ fever ■ mucus in the stool

**Ask a doctor or pharmacist before use if you are taking any drug for**  
 ■ anticoagulation (thinning the blood) ■ diabetes ■ gout ■ arthritis

**When using this product** a temporary, but harmless, darkening of the stool and/or tongue may occur.

**Stop use and ask a doctor if** ■ symptoms get worse or last more than 2 days ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days

**If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222. ▶

\*This product is not manufactured or distributed by The Procter & Gamble Co., owner of the registered trademark Pepto-Bismol®.

LOT: UNVARNISHED AREA

EXP:

70712MJRLR      PID 402888

7 08820 06145 3

Do not print

<b>MEIJER STOMACH RELIEF REGULAR STRENGTH</b>			
bismuth subsalicylate suspension			
Product Information			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:41250-211
<b>Route of Administration</b>	ORAL		
Active Ingredient/Active Moiety			
<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>	
<b>BISMUTH SUBSALICYLATE</b> (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE, SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	525 mg in 15 mL	
Inactive Ingredients			
<b>Ingredient Name</b>	<b>Strength</b>		
<b>CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED</b> (UNII: K679OBS311)			
<b>D&amp;C RED NO. 22</b> (UNII: 1678RKX8RT)			
<b>D&amp;C RED NO. 28</b> (UNII: 767IP0Y5NH)			
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)			
<b>POTASSIUM HYDROXIDE</b> (UNII: WZH3C48M4T)			
<b>POTASSIUM SORBATE</b> (UNII: 1VPU26JZ4)			
<b>WATER</b> (UNII: 059QF0KO0R)			
<b>SALICYLIC ACID</b> (UNII: O414PZ4LPZ)			
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)			
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)			

**SUCRALOSE** (UNII: 96K6UQ3ZD4)

**XANTHAN GUM** (UNII: TTV12P4NEE)

### Product Characteristics

<b>Color</b>	PINK (viscous)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-211-12	354 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/15/2017	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	03/15/2017	

**Labeler** - MEIJER DISTRIBUTION INC (006959555)

Revised: 11/2023

MEIJER DISTRIBUTION INC