

REGULAR STRENGTH PAIN RELIEF- acetaminophen capsule, liquid filled
Chain Drug Marketing Association Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Regular Strength Pain Relief

Drug Facts

Active ingredient (in each softgel)

Acetaminophen 325 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - backache
 - minor pain of arthritis
 - the common cold
 - toothache
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. The maximum daily dose of this product is 10 softgels (3,250 mg) in 24 hours for adults or 5 softgels (1,625 mg) in 24 hours for children. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if the user has liver disease

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin

Stop use and ask a doctor if:

- pain gets worse or lasts more than 10 days in adults
- pain gets worse or lasts more than 5 days in children under 12 years
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed** (see **Overdose warning**)

adults and children 12 years of age and over	<ul style="list-style-type: none"> • take 2 softgels every 4 to 6 hours while symptoms last • do not take more than 10 softgels in 24 hours • do not use for more than 10 days unless directed by a doctor
children 6-11 years of age	<ul style="list-style-type: none"> • take 1 softgel every 4 to 6 hours while symptoms last • do not take more than 5 softgels in 24 hours • do not use for more than 5 days unless directed by a doctor
children under 6 years of age	ask a doctor

Other information

- store at room temperature 15°-30°C (59°-86°F)

Inactive ingredients

FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution and white edible ink

Questions or comments?

Call: **248-449-9300**

PRINCIPAL DISPLAY PANEL

Regular Strength Pain Relief

Acetaminophen 325mg 20 Liquid Gels

NDC 63868-217-20

*Compare to the active ingredient in **TYLENOL®** Regular Strength

by Chenxi Li

EXP. DATE: **NC**

LOT NO: **3551599616**

6
8



NDC 63868-217-20

QC
QUALITY CHOICE

Regular Strength

Pain Relief

Liquid Gels

Acetaminophen, 325mg
Pain Reliever | Fever Reducer



20 Liquid Gels (Liquid Filled Capsules)

*Compare to the active ingredient in **TYLENOL®** Regular Strength

SATISFACTION GUARANTEED

Distributed by: C.D.™, Inc.®
43157 W. 9 Mile Rd
Novi, MI 48375
www.cdm.com
Questions: 248-449-9300

Made in China
CDMA20-01



Drug Facts (continued)

Directions

- do not take more than directed (see Overdose warning)

adults and children 12 years and over	<ul style="list-style-type: none"> ■ take 2 softgels every 4 to 6 hours while symptoms last ■ do not take more than 10 softgels in 24 hours, unless directed by a doctor ■ do not use for more than 10 days unless directed by a doctor 	
children 6 years to under 12 years	<ul style="list-style-type: none"> ■ take 1 softgel every 4 to 6 hours while symptoms last ■ do not take more than 5 softgels in 24 hours ■ do not use for more than 5 days unless directed by a doctor 	ask a doctor
children under 6 years		

Other information

- Store at 15°C-30°C (59°F-86°F), and avoid excessive heat.

Inactive ingredients
FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol, polydioxane, propylene glycol, purified water, sorbitol, sorbitan solution and white ink

Questions or Comments?
Call toll free: 248-449-9300

31
030310
0400

Drug Facts

KEEP OUTER CARTON FOR COMPLETE WARNING AND PRODUCT INFORMATION

Active ingredient (in each softgel)

Purpose

Acetaminophen 325 mg.....Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - backache
 - minor pain of arthritis
 - the common cold
 - toothache
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning:

This product contains acetaminophen. The maximum daily dose of this product is 10 softgels (3,250 mg) in 24 hours for adults or 5 softgels (1,625 mg) in 24 hours for children. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if the user has liver disease

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days in adults
 - pain gets worse or lasts more than 5 days in children under 12 years
 - fever gets worse or lasts more than 3 days
 - new symptoms occur
 - redness or swelling is present
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

NDC 63868-217-20



Regular Strength

Pain Relief

Liquid Gels

Acetaminophen, 325mg
Pain Reliever | Fever Reducer

20 Liquid Gels
(Liquid Filled Capsules)

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SEAL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Warnings

Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 softgels (3,250 mg) in 24 hours for adults or 5 softgels (1,625 mg) in 24 hours for children. Severe liver damage may occur if:

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours, which is the maximum daily amount

taken with other drugs containing acetaminophen

- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Directions ■ do not take more than directed (see Overdose warning)

adults and children 12 years and over:

- take 2 softgels every 4 to 6 hours while symptoms last
- do not take more than 10 softgels in 24 hours
- do not use for more than 10 days unless directed by a doctor

children 6-11 years:

- take 1 softgel every 4 to 6 hours while symptoms last
- do not take more than 5 softgels in 24 hours
- do not use for more than 5 days unless directed by a doctor
- children under 6 years: ■ ask a doctor

Manufactured by C.D. McVee, Inc. B
43157 W 9th Ave, Ste 300
Newport, MI 48167
www.cdmcvee.com
Customer Service: 1-800-428-6880

Made in China
CDMA19-01

LOT NO:

EXP DATE:

REGULAR STRENGTH PAIN RELIEF

acetaminophen capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63868-217
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	

Product Characteristics

Color	red (clear)	Score	no score
Shape	capsule (oblong)	Size	20mm
Flavor		Imprint Code	PC17
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63868-217-20	1 in 1 CARTON	07/13/2017	
1		20 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	07/13/2017	

Labeler - Chain Drug Marketing Association Inc. (011920774)

Establishment

Name	Address	ID/FEI	Business Operations
Humanwell PuraCap Pharmaceutical (Wuhan) Co., Ltd.		421293287	manufacture(63868-217) , analysis(63868-217)

