

**TYLENOL EXTRA STRENGTH- acetaminophen tablet, film coated**  
**Johnson & Johnson Consumer Inc.**

-----  
**Tylenol® Extra Strength**

***Drug Facts***

**Active ingredient (in each caplet)**

Acetaminophen 500 mg

**Purpose**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - the common cold
  - headache
  - backache
  - minor pain of arthritis
  - toothache
  - muscular aches
  - premenstrual and menstrual cramps
- temporarily reduces fever

**Warnings**

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or

pharmacist.

- if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if you have** liver disease

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

### **Overdose warning**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none"><li>▪ take 2 caplets every 6 hours while symptoms last</li><li>▪ do not take more than 6 caplets in 24 hours, unless directed by a doctor</li><li>▪ do not use for more than 10 days unless directed by a doctor</li></ul>
children under 12 years	ask a doctor

### **Other information**

- store between 20-25°C (68-77°F)
- **do not use if carton is opened. Do not use if foil inner seal imprinted with "TYLENOL" is broken or missing**

### **Inactive ingredients**

crospovidone, FD&C red no. 40, FD&C yellow no. 6, guar gum, hypromellose, magnesium stearate, maltodextrin, medium chain triglycerides, microcrystalline cellulose,

polyvinyl alcohol, povidone, pregelatinized starch, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

**Questions or comments?**

call **1-877-895-3665** (toll-free) or **215-273-8755** (collect)

**PRINCIPAL DISPLAY PANEL**

NDC 50580-472-01

Extra Strength

TYLENOL<sup>®</sup>

FOR ADULTS

Acetaminophen

Pain Reliever

Fever Reducer

EASY TO SWALLOW\*

With Gentleglide<sup>™</sup> Coating Technology

Actual Size

24 Caplets

500 mg each

# TYLENOL EXTRA STRENGTH

acetaminophen tablet, film coated

## Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50580-472
Route of Administration	ORAL		

Important: Read all product information before using. Keep this box for important information.

**Drug Facts**

**Active ingredient (in each caplet)** **Purpose**  
Acetaminophen 500 mg ... Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - the common cold
  - backache
  - toothache
  - premenstrual and menstrual cramps
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness
- skin rash
- blisters

If a skin reaction occurs, stop use and seek medical help right away.

**Drug Facts (continued)**

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if you have liver disease**

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not take more than directed (see overdose warning)

30055837/130992

NDC 50580-472-01

# TYLENOL®

FOR ADULTS

Acetaminophen Pain Reliever  
Fever Reducer

EASY TO  
SWALLOW\*

With *GentleGlide*® Coating Technology



24 Caplets  
500 mg each



Actual Size

\*Easier to Swallow Coating than  
Extra Strength TYLENOL® Caplets,  
among those with a preference

www.tylenol.com  
Pat. www.kenvuepats.com

OPEN HERE →

Distributed by:  
**JOHNSON & JOHNSON CONSUMER INC.**  
McNeil Consumer Healthcare Division  
Fort Washington, PA 19034 USA  
©J&JCI 2023

## Drug Facts (continued)

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if you have liver disease**

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not take more than directed (see overdose warning)

## Drug Facts (continued)

adults and children under 12 years

- take 2 caplets every 6 hours while symptoms last
- do not take more than 6 caplets in 24 hours, unless directed by a doctor
- do not use for more than 10 days unless directed by a doctor

ask a doctor

## Other information

- store between 20-25°C (68-77°F)
- do not use if carton is opened. Do not use if foil inner seal imprinted with "TYLENOL" is broken or missing

## Inactive ingredients

croscollidone, FD&C red no. 40, FD&C yellow no. 6, guar gum, hypromellose, magnesium stearate, maltodextrin, medium chain triglycerides, microcrystalline cellulose, polyvinyl alcohol, povidone, pregelatinized starch, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

## Questions or comments?

call 1-877-895-3665 ( toll-free) or 215-273-8755 ( collect)

LOT  
EXP



Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients	
Ingredient Name	Strength
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
POVIDONE (UNII: FZ989GH94E)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
CROSPVIDONE (UNII: 2S7830E561)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GUAR GUM (UNII: E89I1637KE)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TALC (UNII: 7SEV7J4R1U)	

Product Characteristics			
Color	red	Score	no score
Shape	OVAL	Size	18mm
Flavor		Imprint Code	TYLENOL;500
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50580-472-01	1 in 1 CARTON	03/18/2024	
1		24 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:50580-472-02	1 in 1 CARTON	03/18/2024	
2		50 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:50580-472-03	1 in 1 CARTON	03/18/2024	
3		100 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:50580-472-04	1 in 1 CARTON	03/18/2024	

4

200 in 1 BOTTLE; Type 0: Not a Combination Product

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	03/18/2024	

**Labeler** - Johnson & Johnson Consumer Inc. (878046358)

Revised: 12/2023

Johnson & Johnson Consumer Inc.