# SMART SENSE NAPROXEN SODIUM- naproxen sodium tablet, film coated Kmart Corporation

-----

#### Kmart Corporation Naproxen Sodium Tablets, 220 mg Drug Facts

#### **Active ingredient (in each caplet)**

Naproxen sodium 220 mg

(naproxen 200 mg) (NSAID)\*

\*nonsteroidal anti-inflammatory drug

#### **Purposes**

Pain reliever/fever reducer

#### Uses

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

#### **Warnings**

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

#### Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

#### Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have asthma

#### Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

#### When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

#### Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

#### If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

#### Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

#### **Directions**

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and children 12 years and older	<ul> <li>take 1 caplet every 8 to 12 hours while symptoms last</li> <li>for the first dose you may take 2 caplets within the first hour</li> <li>do not exceed 2 caplets in any 8- to 12-hour period</li> <li>do not exceed 3 caplets in a 24-hour period</li> </ul>
Children under 12 years	• ask a doctor

#### Other information

- each caplet contains: sodium 21 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

#### **Inactive ingredients**

FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

#### Questions or comments?

1-800-719-9260

#### **Principal Display Panel**

Compare to the active ingredient of Aleve® Caplets

STRENGTH TO LAST 12 HOURS

PAIN RELIEVER / FEVER REDUCER

Naproxen Sodium Tablets, 200 mg

Pain Reliever / Fever Reducer (NSAID)

**ACTUAL SIZE** 

\*\*Capsule-Shaped Tablets

100 CAPLETS\*\*



: 36878 FT (3

EXP.

LOT NO.



KEEP CARTON FOR REFERENCE

#### DO NOT USE IF PRINTED FOIL UNDER CAP IS BROKEN OR MISSING

### Drug Facts

Active ingredient (in each caplet) Purposes

Naproxen sodium 220 mg Pain reliever/ (naproxen 200 mg) (NSAID)\*.....fever reducer \*nonsteroidal anti-inflammatory drug

#### Uses

- temporarily relieves minor aches and pains due to:
  - minor pain of arthritis
     muscular aches
  - backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

#### Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin Symptoms may include:

- hives
- facial swelling
- asthma (wheezing) shock
- skin reddening
- rash blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are ace 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

#### Drug Facts (continued)

#### Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

#### Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- vou have asthma

#### Ask a doctor or pharmacist before use if you are


- under a doctor's care for any serious condition
- taking any other drug

#### When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

#### Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- vou have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear



#### **SMART SENSE NAPROXEN SODIUM**

naproxen sodium tablet, film coated

Product Information				
Product Type	HUMAN OTC DRUG LABEL	Item Code (Source)	NDC:49738-368	
Route of Administration	ORAL	DEA Sche dule		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
NAPRO XEN SO DIUM (NAPRO XEN)	NAPROXEN SODIUM	220 mg	

Inactive Ingredients			
Ingredient Name	Strength		
HYPROMELLOSES			
MAGNESIUM STEARATE			
CELLULO SE, MICRO CRYSTALLINE			
POLYETHYLENE GLYCOLS			
POVIDONES			
TALC			
TITANIUM DIO XIDE			

Product Characteristics				
Color	BLUE (Light Blue)	Score	no score	
Shape	CAPSULE (oval)	Size	12mm	
Flavor		Imprint Code	L368	
Contains				

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:49738-368-62	1 in 1 CARTON			
1		24 in 1 BOTTLE			
2	NDC:49738-368-71	1 in 1 CARTON			
2		50 in 1 BOTTLE			
3	NDC:49738-368-78	1 in 1 CARTON			
3		100 in 1 BOTTLE			
4	NDC:49738-368-82	1 in 1 CARTON			
4		200 in 1 BOTTLE			
5	NDC:49738-368-76	1 in 1 CARTON			
5		120 in 1 BOTTLE			

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA074661	10/08/2010		

## Labeler - Kmart Corporation (008965873)

Revised: 4/2014 Kmart Corporation