

**MUCUS RELIEF SEVERE CONGESTION COUGH NIGHT TIME COLD AND FLU MAXIMUM STRENGTH- dextromethorphan hbr, guaifenesin, phenylephrine hci, acetaminophen, diphenhydramine hci, phenylephrine hci Dolgencorp, Inc. (DOLLAR GENERAL & REXALL)**

-----  
**DRUG FACTS**

**Active ingredients for Nighttime (in each 20 mL)**

**Acetaminophen 650 mg**

Diphenhydramine HCl 25 mg

Phenylephrine HCl 10 mg

**Active ingredients for Daytime (in each 20 mL)**

Dextromethorphan HBr 20 mg

Guaifenesin 400 mg

Phenylephrine HCl 10 mg

**Purpose for Nighttime**

**Pain reliever/fever reducer**

Antihistamine/cough suppressant

Nasal decongestant

**Purpose for Daytime**

Cough suppressant

Expectorant

Nasal decongestant

**Uses**

**Nighttime**

- temporarily relieves these common cold and flu symptoms
- cough
- nasal congestion
- minor aches and pains
- sore throat
- headache
- runny nose
- sneezing

- temporarily reduces fever
- controls cough to help you get to sleep

### **Daytime**

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make cough more productive
- temporarily relieves
  - cough due to minor throat and bronchial irritations as may occur with the common cold or inhaled irritants
  - the intensity of coughing
  - the impulse to cough to help you get to sleep
  - nasal congestion due to a cold

### **Warnings**

#### **NIGHTTIME**

**Liver warning:** This product contain acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash.

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### **Do not use**

##### **Nighttime**

- with any drug containing acetaminophen (prescription or nonprescription) . If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other drug containing diphenhydramine, even one used on the skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product
- for children under 12 years of age

##### **Daytime**

- for children under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI)(certain drugs for depression,psychiatric or emotional conditions, or Parkinson's disease), or

for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

#### **Nighttime**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

#### **Daytime**

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

### **Ask a doctor or pharmacist before use if you are**

#### **Nighttime**

- you are taking the blood thinning drug warfarin
- you are taking sedative or tranquilizers

### **When using these products**

#### **Nighttime**

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

#### **Daytime**

- **do not use more than directed**

### **Stop use and ask a doctor if**

#### **Nighttime**

- nervousness, dizziness, or sleeplessness occur

- pain, nasal congestion, or cough gets worse, or lasts more than 7 days
- fever gets worse, or last more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be a signs of a serious condition

### **Daytime**

- nervousness, dizziness or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

### **Nighttime and DayTime**

ask a health professional before use.

### **Keep out of reach of children.**

### **Nighttime**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **DayTime**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

### **Directions**

#### **Nighttime**

- **do not take more than directed (see overdose warning)**
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device
- keep dosing cup with product
- mL = milliliter
- dose as follows or as directed by a doctor
- adults and children 12 years and older: 20 mL every 4 hours while symptoms last
- children under 12 years of age: do not use

#### **Daytime**

- do not take more than 6 doses in any 24-hours period
- measure only with dosing cup provided. Do not use any other dosing device.
- shake well before using
- keep dosing cup with product
- mL = milliliter
- adults and children 12 years of age and older: 20 mL every 4 hours

- children under 12 years of age: do not use

## **Other information**

### **Nighttime**

- **each 20 mL contains:** sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

### **Daytime**

- **each 20 mL contains:** 17 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

## **Inactive ingredients**

### **Nighttime**

citric acid, disodium EDTA, FD&C Blue #1, FD&C red #40, Flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

### **Daytime**

citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

## **Questions or comments?**

### **Nighttime and DayTime**

Call **1-888-309-9030**

## **Principal Display Panel**

Compare to the active ingredients of Maximum Strength Mucinex® Fast-Max® Severe Congestion & Cough & Night Time Cold & Flu\*

### **NIGHTTIME**

#### **Maximum Strength**

#### **Fast Acting Night Time Cold & Flu**

Multi-Symptom Relief

Acetaminophen

DiphenhydramineHCl

Phenylephrine HCl

Pain Reliever/Fever reducer

Antihistamine/Cough Suppressant

Nasal Decongestant

- For ages 12 years and over

**DAYTIME**

Maximum Strength

Fast Acting

Mucus Relief

Severe Congestion & Cough

Multi-Symptom Relief

Dextromethorphan HBr

Guaifenesin

Phenylephrine HCl

Cough Suppressant

Expectorant

Nasal Decongestant

- For ages 12 years and over

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.**

↑ ↑ This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® FAST-MAX® Severe Congestion & Cough and Nighttime Cold & Flu.

DISTRIBUTED BY OLD EAST MAIN CO.

100 MISSION RIDGE

GOODLETTSVILLE, TN 37072

**Product Label**

## DOLLAR GENERAL HEALTH Maximum Strength Mucus Relief Severe Congestion and Cough, Cold Flu

# MUCUS RELIEF SEVERE CONGESTION COUGH NIGHT TIME COLD AND FLU MAXIMUM STRENGTH

dextromethorphan hbr, guaifenesin, phenylephrine hci, acetaminophen, diphenhydramine hci, phenylephrine hci tit

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55910-562
---------------------	----------------	---------------------------	---------------

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-562-12	1 in 1 KIT; Type 0: Not a Combination Product	03/31/2018	

### Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE, PLASTIC	177 mL
Part 2	1 BOTTLE, PLASTIC	177 mL

## Part 1 of 2

# MUCUS RELIEF CONGESTION COUGH MAXIMUM STRENGTH SEVERE CONGESTION AND COUGH

dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride liquid

### Product Information

<b>Item Code (Source)</b>	NDC:55910-537
---------------------------	---------------

Route of Administration ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>EDETATE CALCIUM DISODIUM</b> (UNII: 25IH6R4SGF)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>TRISODIUM CITRATE DIHYDRATE</b> (UNII: B22547B95K)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYL GALLATE</b> (UNII: 8D4SNN7V92)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/31/2018	

### Part 2 of 2

### NIGHTTIME COLD AND FLU MAXIMUM STRENGTH

acetaminophen, diphenhydramine hci, phenylephrine hci liquid

## Product Information

**Item Code (Source)** NDC:55910-460

**Route of Administration** ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 20 mL
<b>DIPHENHYDRAMINE HYDROCHLORIDE</b> (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg in 20 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CITRATE</b> (UNII: 1Q73Q2JULR)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>PROPYL GALLATE</b> (UNII: 8D4SNN7V92)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>EDETATE CALCIUM DISODIUM</b> (UNII: 25IH6R4SGF)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/31/2018	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/31/2018	

**Labeler** - Dolgencorp, Inc. (DOLLAR GENERAL & REXALL) (068331990)

Revised: 10/2023

Dolgencorp, Inc. (DOLLAR GENERAL & REXALL)