

**BLU-KOTE VETERINARY ANTISEPTIC-PROTECTIVE WOUND DRESSING - sodium propionate, gentian violet, acriflavine aerosol**  
**H. W. Naylor Company Inc.**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, [click here](#).*

-----

**BLU-KOTE® Veterinary Antiseptic-Protective Wound Dressing**

**Active Ingredients**

Sodium Propionate 9.1%, Gentian Violet 1%, Acriflavine 0.15%

**Purpose**

For the effective treatment of FUNGUS INFECTIONS, RINGWORM, SURFACE WOUNDS, CUTS, GALLS, HOOFF-FOOT and PAD SORES, CHAFES, ABRASIONS, MOIST OR SCABBY LESIONS, ITCHY FUNGUS ECZEMA and SORES.

**Directions**

**SPRAY IT ON HORSES • DOGS**

Clean and dry affected area thoroughly before spraying.

Scrub and remove scale and scab on Ringworm before first application.

Shake well.

Remove protective cap.

Hold upright.

Point nozzle opening toward lesion and spray from a distance of 4-8 inches.

Only a light application is needed.

Repeat once or twice a day or as indicated until healing occurs.

**Safety**

Caution: For external veterinary use only.

In case of serious burns, deep or puncture wounds, or if redness, irritation or swelling persists or increases, discontinue use and consult veterinarian.

Keep away from eyes, mouth, nostrils, and mucous membranes.

Do not spray in eyes.

Do not inhale.

**DO NOT USE ON CATS.**

**PREVENT DOGS FROM LICKING TREATED AREAS.**

**WARNING**

Contents under pressure.

Do not puncture.

Do not store or use near heat or open flame.

Exposure to temperatures above 130°F may cause bursting.

Never throw container into fire or incinerator.

BLUE-KOTE® STAINS, WET OR DRY.

**NOT FOR USE ON HORSES INTENDED FOR FOOD.**

**FOR ANIMAL USE ONLY.**

**USE ONLY AS DIRECTED.**

**KEEP FROM REACH OF CHILDREN.**

**Other Ingredients**

Water, Urea, Glycerine, Isopropyl Alcohol 32% by volume.

**Principal Display Panel**

*Dr. Naylor*®

**BLU-KOTE**®

**Veterinary Antiseptic - Protective Wound Dressing**

**Germicidal • Fungicidal**

**Net Weight 5 oz. (142 grams)**

**DANGER: EXTREMELY FLAMMABLE**

**H.W. NAYLOR CO., INC.**

Morris, N.Y. 13808

U.S.A.

BLU-KOTE® is a germicidal, fungicidal, wound dressing and healing aid effective against both bacterial and fungus infections most common in skin lesions of domestic animals. Kills Ringworm and fungus infections. Dries up blisters and pox-like scabby sores or lesions.

**“Instant Medication” Quick Drying – Long Acting Deep Penetrating**

Made in U.S.A.

Spec. 5040

## SPRAY IT ON HORSES • DOGS

For the effective treatment of FUNGUS INFECTIONS, RINGWORM, SURFACE WOUNDS, CUTS, GALLS, HOOF-FOOT and PAD SORES, CHAFES, ABRASIONS, MOIST OR SCABBY LESIONS, ITCHY FUNGUS ECZEMA and SORES.

BLU-KOTE® is a germicidal, fungicidal, wound dressing and healing aid effective against both bacterial and fungus infections most common in skin lesions of domestic animals. Kills Ringworm and fungus infections. Dries up blisters and pox-like scabby sores or lesions.

**"Instant Medication"**

**Quick Drying – Long Acting  
Deep Penetrating**

Made in U.S.A.  
Spec. 5040





# BLU-KOTE®

**Veterinary  
Antiseptic – Protective  
Wound Dressing**

**Germicidal • Fungicidal**

Active Ingredients: Sodium Propionate 9.1%,  
Gentian Violet 1%, Acriflavine 0.15%, in a special  
base of Water, Urea, Glycerine, Isopropyl Alcohol  
32% by volume.

**Net Weight 5 oz. (142 grams)**  
**DANGER: EXTREMELY FLAMMABLE**

H.W. NAYLOR CO., INC.  
Morris, N.Y. 13808  
U.S.A.

**Directions:** Clean and dry affected area thoroughly before spraying. Scrub and remove scale and scab on Ringworm before first application. Shake well. Remove protective cap. Hold upright. Point nozzle opening toward lesion and spray from a distance of 4-8 inches. Only a light application is needed. Repeat once or twice a day or as indicated until healing occurs.

**Caution:** For external veterinary use only. In case of serious burns, deep or puncture wounds, or if redness, irritation or swelling persists or increases, discontinue use and consult veterinarian. Keep away from eyes, mouth, nostrils, and mucous membranes. Do not spray in eyes. Do not inhale. **DO NOT USE ON CATS. PREVENT DOGS FROM LICKING TREATED AREAS.**

**Warning:** Contents under pressure. Do not puncture. Do not store or use near heat or open flame. Exposure to temperatures above 130°F may cause bursting. Never throw container into fire or incinerator.

BLUE-KOTE® STAINS, WET OR DRY.

**NOT FOR USE ON  
HORSES INTENDED FOR FOOD.  
FOR ANIMAL USE ONLY.  
KEEP FROM REACH OF CHILDREN.  
USE ONLY AS DIRECTED.**

## BLU-KOTE VETERINARY ANTISEPTIC-PROTECTIVE WOUND DRESSING

sodium propionate, gentian violet, acriflavine aerosol

### Product Information

<b>Product Type</b>	OTC ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:15878-231
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>Sodium Propionate</b> (UNII: DK6 Y9 P42IN) (Propionic Acid - UNII:JHU490RVYR)	Sodium Propionate	90 mg in 1000 mL
<b>Gentian Violet</b> (UNII: J4Z741D6O5) (Gentian Violet Cation - UNII:3GVJ31T6YY)	Gentian Violet	10 mg in 1000 mL
<b>Acriflavine</b> (UNII: 1T3A50395T) (Acriflavine - UNII:1T3A50395T)	Acriflavine	1.7 mg in 1000 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>Water</b> (UNII: 059QF0KO0R)	
<b>Urea</b> (UNII: 8W8T17847W)	
<b>Isopropyl Alcohol</b> (UNII: ND2M416302)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:15878-231-12	148 mL in 1 CAN		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		01/01/1964	

**Labeler** - H. W. Naylor Company Inc. (002080190)

## Establishment

Name	Address	ID/FEI	Business Operations
H. W. Naylor Company Inc.		002080190	API MANUFACTURE, MANUFACTURE

Revised: 1/2013

H. W. Naylor Company Inc.