

EXCEDRIN EXTRA STRENGTH PAIN RELIEVER- acetaminophen, aspirin (nsaid), and caffeine tablet, film coated
Haleon US Holdings LLC

Drug Facts

Active ingredients (in each caplet)

Acetaminophen 250 mg

Aspirin 250 mg (NSAID*)

Caffeine 65 mg

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever

Pain reliever

Pain reliever aid

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - a cold
 - arthritis
 - muscular aches
 - toothache
 - premenstrual and menstrual cramps

Warnings

Reye's syndrome:Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert:Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert:Aspirin may cause a severe allergic reaction which may include:

- hives

- facial swelling
- asthma (wheezing)
- shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Do not use

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking

- a prescription drug for diabetes, gout, or arthritis
- any other drug, or are under a doctor's care for any serious condition

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood

- have bloody or black stools
- have stomach pain that does not get better
- ringing in the ears or loss of hearing occurs
- painful area is red or swollen
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- any new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed**
- drink a full glass of water with each dose
- adults and children 12 years of age and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours
- children under 12 years: ask a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- close cap tightly after use
- read all product information before using. Keep this box for important information.

Inactive ingredients

corn starch, crospovidone, hypromellose, microcrystalline cellulose, povidone, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, stearic acid, titanium dioxide

Questions?

1-800-468-7746

Principal Display Panel

NDC 0067-8203-04

EXCEDRIN

EXTRA STRENGTH

**Acetaminophen, Aspirin (NSAID) and Caffeine
Pain Reliever/Pain Reliever Aid**

300 CAPLETS

TAMPER-EVIDENT BOTTLE

**DO NOT USE IF PRINTED INNER SAFETY SEAL
UNDER CAP IS BROKEN OR MISSING**

Distributed by: **GSK Consumer Healthcare**

Warren, NJ 07059

©2020 GSK group of companies or its licensor.

Trademarks are owned by or licensed to the GSK group of companies.

B-0630-763-17 - Front Carton

NDC 0067-8203-04

EXCEDRIN

EXTRA STRENGTH

Acetaminophen, Aspirin (NSAID) and Caffeine
Pain Reliever/Pain Reliever Aid

300
CAPLETS



EXCEDRIN EXTRA STRENGTH PAIN RELIEVER

acetaminophen, aspirin (nsaid), and caffeine tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0067-8203
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CROSPROVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	white (White)	Score	no score
Shape	OVAL	Size	17mm
Flavor		Imprint Code	ES
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0067-8203-03	1 in 1 CARTON	01/15/2021	
1		200 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:0067-8203-04	1 in 1 CARTON	03/19/2021	
2		300 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:0067-8203-02	1 in 1 CARTON	02/22/2021	
3		100 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	01/15/2021	

Revised: 4/2024

Haleon US Holdings LLC