

**TUSSIN COUGH AND CHEST CONGESTION DM MAX- dextromethorphan hbr,
guaifenesin solution**

Rite Aid Corporation

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Rite Aid Corporation Tussin Cough & Chest Congestion DM MAX Drug Facts

Active ingredients (in each 20 mL)

Dextromethorphan HBr, USP 20 mg

Guaifenesin, USP 400 mg

Purposes

Cough suppressant

Expectorant

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Stop use and ask a doctor if

cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	20 mL every 4 hours
children under 12 years	do not use

Other information

- **each 20 mL contains:** sodium 13 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

acetic acid, anhydrous citric acid, carboxymethylcellulose sodium, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin, menthol, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose, xanthan gum

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

SEE NEW DOSING

Compare to the active ingredients of Robitussin® Maximum Strength Cough + Chest Congestion DM

MAXIMUM STRENGTH

ADULT

tussin

cough & chest

congestion DM

MAX

dextromethorphan HBr, USP 20 mg

guaifenesin, USP 400 mg

cough suppressant

expectorant

NON-DROWSY
 ALCOHOL FREE
 relieves:
 cough
 mucus
 for ages 12 & over
 4 FL OZ (118 mL)



SEE NEW DOSING
RITE AID
*Compare to the active ingredients of Robitussin® Maximum Strength Cough + Chest Congestion DM
PHARMACY **MAXIMUM STRENGTH**

ADULT
tussin
cough & chest
congestion DM
MAX

dextromethorphan HBr, USP 20 mg
guaifenesin, USP 400 mg

cough suppressant
 expectorant

NON-DROWSY
ALCOHOL FREE

relieves:
 cough
 mucus
 for ages 12 & over

4 FL OZ (118 mL)

SEE NEW DOSING
RITE AID
*Compare to the active ingredients of Robitussin® Maximum Strength Cough + Chest Congestion DM
PHARMACY **MAXIMUM STRENGTH**

ADULT
tussin
cough & chest
congestion DM
MAX

dextromethorphan HBr, USP 20 mg
guaifenesin, USP 400 mg

cough suppressant
 expectorant

NON-DROWSY
ALCOHOL FREE

relieves:
 cough
 mucus
 for ages 12 & over

4 FL OZ (118 mL)

MAXIMUM STRENGTH
ADULT
tussin
cough & chest
congestion DM
MAX
RITE AID PHARMACY
 dextromethorphan HBr, USP 20 mg
 guaifenesin, USP 400 mg
 cough suppressant
 expectorant

Drug Facts

Active ingredients (in each 20 mL)
 Dextromethorphan HBr, USP 20 mg.....Cough suppressant
 Guaifenesin, USP 400 mg.....Expectorant

Purposes
 ■ temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
 ■ helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

Warnings
 Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have
 ■ cough that occurs with too much phlegm (mucus)
 ■ cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema
 Stop use and ask a doctor if cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
 ■ do not take more than 6 doses in any 24-hour period
 ■ measure only with dosing cup provided
 ■ keep dosing cup with product
 ■ mL = milliliter
 ■ this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	20 mL every 4 hours
children under 12 years	do not use

Other information
 ■ each 20 mL contains sodium 13 mg
 ■ store at 20-25°C (68-77°F). Do not refrigerate.

Drug Facts (continued)

Inactive ingredients acetic acid, anhydrous citric acid, carboxymethylcellulose sodium, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin, menthol, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose, xanthan gum

Questions or comments? 1-800-719-9260

*This product is not manufactured or distributed by Pfizer, distributor of Robitussin® Maximum Strength Cough + Chest Congestion DM.

DISTRIBUTED BY: RITE AID
 30 HUNTER LANE, CAMP HILL, PA 17011

SATISFACTION GUARANTEE
RITE AID
 IF YOU'RE NOT SATISFIED, WE'LL HAPPILY REFUND YOUR MONEY.

Dosage Cup Included

PARENTS:
 Learn about teen medicine abuse
www.StopMedicineAbuse.org

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING

RA# 999536

 0 11822 83553 4

2B726 83 C1

TUSSIN COUGH AND CHEST CONGESTION DM MAX

dextromethorphan hbr, guaifenesin solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-1200
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
MENTHOL (UNII: L7T10EIP3A)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	RED	Score	
Shape		Size	
Flavor	FRUIT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-1200-1	1 in 1 CARTON	08/02/2018	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:11822-1200-2	1 in 1 CARTON	08/02/2018	
2		237 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	08/02/2018	

Labeler - Rite Aid Corporation (014578892)

Revised: 12/2019

Rite Aid Corporation