

**MENSTRUAL PAIN RELIEF MAXIMUM STRENGTH- acetaminophen, pamabrom,
pyrilamine maleate tablet, film coated
CHAIN DRUG MARKETING ASSOCIATION INC**

Quality Choice 44-679

Active ingredients (in each caplet)

Acetaminophen 500 mg
Pamabrom 25 mg
Pyrilamine maleate 15 mg

Purpose

Pain reliever
Diuretic
Antihistamine

Uses

for the temporary relief of these symptoms associated with menstrual periods:

- headache
- bloating
- cramps
- backache
- muscular aches
- irritability
- water-weight gain

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- difficulty in urination due to enlargement of the prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- liver disease
- glaucoma

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- drowsiness may occur
- avoid alcoholic beverages
- excitability may occur, especially in children
- alcohol, sedatives and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**
- adults and children 12 years and over:
 - take 2 caplets with water every 6 hours as needed
 - do not exceed 6 caplets in a 24 hour period or as directed by a doctor
- children under 12 years: ask a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, croscarmellose sodium, crospovidone, FD&C blue #2 aluminum lake, FD&C red #40 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, shellac wax, silicon dioxide, stearic acid, talc, titanium dioxide

Questions or comments?**1-800-426-9391****Principal Display Panel****QC
QUALITY®
CHOICE**

NDC 63868-961-20

**Compare to the
Active Ingredients in
Maximum Strength
Pamprin® Multi-Symptom***

Maximum Strength

Menstrual Pain Relief**Multi-Symptom****Acetaminophen** - Pain Reliever

Pamabrom - Diuretic

Pyrilamine Maleate - Antihistamine

Relieves: Cramps, Irritability, Headache, Bloating, Backache

20 Caplets

actual size

*This product is not manufactured or distributed by Focus Consumer Healthcare, LLC, owner of the registered trademark Maximum Strength Pamprin® Multi-Symptom.

50844 REV0718A67909

**QC
100%
SATISFACTION**

GUARANTEED

Distributed by C.D.M.A., Inc. ©
43157 W 9 Mile Rd
Novi, MI 48375
www.qualitychoice.com
Questions 248-449-9300

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

No Print / No Varnish
Lot no. & Exp. date



NDC 63868-961-20

Compare to the
Active Ingredients in
Maximum Strength
Pamprin® Multi-Symptom*

Menstrual Pain Relief

Multi-Symptom

Acetaminophen - Pain Reliever
Pamabrom - Diuretic
Pyrilamine Maleate - Antihistamine

Relieves: Cramps, Irritability, Headache, Bloating, Backache



20 Caplets actual size

*This product is not manufactured or distributed by Focus Consumer Healthcare, LLC, owner of the registered trademark Maximum Strength Pamprin® Multi-Symptom.
50844 REV0718A67909

Distributed by C.D.M.A., Inc.®
43157 W 9 Mile Rd
Novi, MI 48375
www.qualitychoice.com
Questions: 248-449-9300

TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts (continued)
Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed
- adults and children 12 years and over:
 - take 2 caplets with water every 6 hours as needed
- do not exceed 6 caplets in a 24 hour period or as directed by a doctor
- children under 12 years: ask a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients corn starch, croscarmellose sodium, crospovidone, FD&C blue #2 aluminum lake, FD&C red #40 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, shellac, wax, silicon dioxide, stearic acid, talc, titanium dioxide

Questions or comments?
1-800-426-9391

Drug Facts (continued)

not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- difficulty in urination due to enlargement of the prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- liver disease
- glaucoma

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- drowsiness may occur
- avoid alcoholic beverages
- excitability may occur, especially in children
- alcohol, sedatives and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

B-0220-679-09
REV0718A67909

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Drug Facts

Active ingredients (in each caplet)

Acetaminophen 500 mg Pain reliever
Pamabrom 25 mg Diuretic
Pyrilamine maleate 15 mg Antihistamine

Uses for the temporary relief of these symptoms associated with menstrual periods:

- headache
- bloating
- cramps
- backache
- muscular aches
- irritability
- water-weight gain

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription), if you are

6 355151 99445 4

44-679

MENSTRUAL PAIN RELIEF MAXIMUM STRENGTH

acetaminophen, pamabrom, pyrilamine maleate tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63868-961
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg
PAMABROM (UNII: UA8U0KJM72) (BROMOTHEOPHYLLINE - UNII:FZG87K1MQ6)	PAMABROM	25 mg
PYRILAMINE MALEATE (UNII: R35D29L3ZA) (PYRILAMINE - UNII:HPE317O9TL)	PYRILAMINE MALEATE	15 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
FD&C BLUE NO. 2 ALUMINUM LAKE (UNII: 4AQJ3LG584)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SHELLAC (UNII: 46N107B71O)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	purple	Score	no score
Shape	OVAL	Size	17mm
Flavor		Imprint Code	44;679
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63868-961-20	1 in 1 CARTON	01/13/2015	
1		20 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	01/13/2015	

Labeler - CHAIN DRUG MARKETING ASSOCIATION INC (011920774)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(63868-961)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	pack(63868-961)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(63868-961)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	pack(63868-961)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(63868-961)

Revised: 4/2024

CHAIN DRUG MARKETING ASSOCIATION INC