

DRAMAMINE FOR KIDS- dimenhydrinate tablet, chewable
Medtech Products Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Dramamine for Kids

Drug Facts

Active ingredient

(in each tablet)

Dimenhydrinate 25 mg

Purpose

Antiemetic

Use

for prevention and treatment of these symptoms associated with motion sickness:

- nausea
- vomiting
- dizziness

Warnings

Do not use for

children under 2 years of age unless directed by a doctor

Ask a doctor before use if the child has

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma

Ask a doctor or pharmacist before use if the child is

taking sedatives or tranquilizers

When using this product

- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

If pregnant or breast-feeding,

ask a doctor before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222)

right away.

Directions

- to prevent motion sickness, the first dose should be taken ½ to 1 hour before starting activity
- to prevent or treat motion sickness, see below:

Children 2 to under 6 years	<ul style="list-style-type: none">• give ½ to 1 chewable tablet every 6-8 hours• do not give more than 3 chewable tablets in 24 hours, or as directed by a doctor
Children 6 to under 12 years	<ul style="list-style-type: none">• give 1 to 2 chewable tablets every 6-8 hours• do not give more than 6 chewable tablets in 24 hours, or as directed by a doctor

Other information

- Phenylketonurics: contains phenylalanine 0.84 mg per tablet
- store at room temperature 68° - 77°F (20° - 25°C)
- **do not use if blister is broken or torn**

Inactive ingredients

aspartame, citric acid, flavor, magnesium stearate, methacrylic acid copolymer, sorbitol

Questions?

1-800-382-7219 Dramamine.com

PRINCIPAL DISPLAY PANEL

#1 Pharmacist Recommended Brand

Ages 2 – 12

Safe, Precise Dose

Children’s Chewable Formula

Dramamine MOTION SICKNESS RELIEF for Kids

8 Chewable Tablets

Dye Free Grape Flavor

#1 Pharmacist Recommended Brand

Ages 2-12
Safe, Precise Dose

Children's Chewable Formula

Dramamine®

MOTION SICKNESS RELIEF
for kids

NEW!

SAFE TRAVELS...

8 Chewable Tablets

Travel Case

Dye Free

Grape Flavor

Important - Keep This Card for Full Product Information

LACTOSE FREE

Dramamine®

MOTION SICKNESS RELIEF
for kids

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Purpose Dimenhydrinate 25 mg..... Antiemetic</p> <p>Use for prevention and treatment of these symptoms associated with motion sickness: ■ nausea ■ vomiting ■ dizziness</p> <p>Warnings Do not use for children under 2 years of age unless directed by a doctor</p> <p>Ask a doctor before use if the child has ■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma</p> <p>Ask a doctor or pharmacist before use if the child is taking sedatives or tranquilizers</p> <p>When using this product ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery</p> <p>If pregnant or breast-feeding, ask a doctor before use.</p> <p>Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. ▶</p>	<p>Drug Facts (continued)</p> <p>Directions ■ to prevent motion sickness, the first dose should be taken ½ to 1 hour before starting activity ■ to prevent or treat motion sickness, see below:</p> <p>Children 2 to under 6 years ■ give ½ to 1 chewable tablet every 6-8 hours ■ do not give more than 3 chewable tablets in 24 hours, or as directed by a doctor</p> <p>Children 6 to under 12 years ■ give 1 to 2 chewable tablets every 6-8 hours ■ do not give more than 6 chewable tablets in 24 hours, or as directed by a doctor</p> <p>Other information ■ Phenylethanolamines: contains phenylalanine 0.84 mg per tablet ■ store at room temperature 68°–77°F (20°–25°C) ■ do not use if blister is broken or torn</p> <p>Inactive ingredients aspartame, citric acid, flavor, magnesium stearate, methacrylic acid copolymer, sorbitol</p> <p>Questions? 1-800-362-7219 Dramamine.com</p>
--	--

©2012 Dist. By: Medtech Products Inc., Irvington, NY 10533
A Prestige Brands Company Made in USA DAO15102

8 31248 00301 6

DRAMAMINE FOR KIDS

dimenhydrinate tablet, chewable

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63029-904
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DIMENHYDRINATE (UNII: JB937PER5C) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIMENHYDRINATE	25 mg

Inactive Ingredients

Ingredient Name	Strength
ASPARTAME (UNII: Z0H242BBR1)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color	WHITE	Score	2 pieces
-------	-------	-------	----------

Shape	ROUND	Size	10 mm
Flavor	GRAPE	Imprint Code	DRA
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63029-904-01	1 in 1 BLISTER PACK		
1		8 in 1 CASE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part336	02/15/2012	

Labeler - Medtech Products Inc. (122715688)

Establishment

Name	Address	ID/FEI	Business Operations
Contract Pharmacal Corporation		057795122	MANUFACTURE(63029-904)

Revised: 4/2012

Medtech Products Inc.