

NAPROXEN SODIUM HEADACHE PAIN- naproxen sodium tablet, coated
NAPROXEN SODIUM BACK AND MUSCLE PAIN- naproxen sodium tablet, coated
Dr.Reddy's Laboratories Inc

**NAPROXEN SODIUM
TABLETS USP, 220 mg**

PAIN RELIEVER / FEVER REDUCER (NSAID)

Drug Facts

Active ingredient (in each tablet/caplet)

Naproxen sodium USP, 220 mg
(naproxen USP, 200 mg) (NSAID)¹

1 nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - menstrual cramps
 - headache
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash

- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks everyday while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- If you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing

- weakness in one part or side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose

Adults and children 12 years and older	▪ take 1 tablet/caplet every 8 to 12 hours while symptoms last ▪ for the first dose you may take 2 tablets/caplets within the first hour ▪ do not exceed 2 tablets/caplets in any 8 to 12 hour period ▪ do not exceed 3 tablets/caplets in a 24-hour period
Children under 12 years	▪ ask a doctor

Other information

- **each tablet/caplet contains:** sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients

FD &C Blue # 2, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, talc, titanium dioxide

Questions or comments?

call toll-free weekdays 9 AM to 8 PM EST at **1-888-375-3784**

Distributed by:

Dr. Reddy's Laboratories, Inc.

Princeton, NJ 08540

Made in India

Rev: 03/2022

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL SECTION

Headache Pain - Caplet Container

Dr.Reddy's

NDC 43598-072-01

HEADACHE PAIN

Naproxen Sodium

Tablets USP, 220 mg (NSAID)

Pain reliever / fever reducer

STRENGTH TO LAST 12 HOURS

for temporary relief of headache pain

100 Caplets

(CAPSULE-SHAPED TABLETS)

TAMPER EVIDENT: Do not use if foil seal under cap printed with "sealed for your protection" is broken or missing. IMPORTANT: Read the directions and warnings before use. See Carton for Full Information

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, or blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use** if you have ever had an allergic reaction to any other pain reliever/fever reducer or right before or after heart surgery. **Ask a doctor before use** if the stomach bleeding warning applies to you, you have a history of stomach problems (such as heartburn), you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke, you are taking a diuretic, you have problems or serious side effects

Peel Here

LOT

EXP

1500XXXXX

Distributed by: Dr. Reddy's Laboratories, Inc.

Princeton, NJ 08540

Made in India

Rev: 03/2022

1500XXXXX

from taking pain relievers or fever reducers. **Ask a doctor or pharmacist before use if you are** under a doctor's care for any serious condition, or taking any other drug. **When using this product** take with food or milk if stomach upset occurs. **Stop use and ask a doctor** if you experience any of the following signs of stomach bleeding (feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better), you have symptoms of heart problems or stroke (chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling), pain gets worse or lasts more than 10 days, fever gets worse or lasts more than 3 days, you have difficulty swallowing, it feels like the pill is stuck in your throat, redness or swelling is present in the painful area, or any new symptoms appear. **If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions Do not take more than directed. The smallest effective dose should be used. Drink a full glass of water with each dose. Adults and children 12 years and older: take 1 caplet every 8 to 12 hours while symptoms last. For the first dose you may take 2 caplets within the first hour. Do not exceed 2 caplets in any 8- to 12-hour period. Do not exceed 3 caplets in a 24-hour period. Children under 12 years: ask a doctor.

Other information Each caplet contains: sodium 20 mg. Store at 20°-25°C (68°-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Questions or comments? call toll-free weekdays 9 AM to 8 PM EST at 1-888-375-3784

Headache Pain - Tablet Container

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- If you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat

Drug Facts (continued)

- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and children 12 years and older	■ take 1 caplet every 8 to 12 hours while symptoms last ■ for the first dose you may take 2 caplets within the first hour ■ do not exceed 2 caplets in any 8- to 12-hour period ■ do not exceed 3 caplets in a 24-hour period
Children under 12 years	■ ask a doctor

Other information

- each caplet contains: sodium 20 mg
- store at 20° - 25°C (68° - 77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients

FD & C Blue # 2, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, talc, titanium dioxide

Questions or comments?
call toll-free weekdays 9 AM to 8 PM EST at 1-888-375-3784

Drug Facts

Active ingredient (in each caplet)
Naproxen sodium USP, 220 mg (naproxen USP, 200 mg) (NSAID) *.....Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - minor pain of arthritis
 - muscular aches
 - backache
 - menstrual cramps
 - toothache
 - the common cold
 - temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Dr.Reddy's

HEADACHE PAIN

Naproxen Sodium
Tablets USP, 220 mg (NSAID)

100 Caplets (CAPSULE-SHAPED TABLETS)

Pain reliever / fever reducer
for temporary relief of headache pain

STRENGTH TO LAST 12 HOURS

TAMPER EVIDENT: Do not use if foil seal under cap printed with "sealed for your protection" is broken or missing.

IMPORTANT: Read the directions and warnings before use. Keep the carton for reference, it contains important information.

* * * This product is not manufactured or distributed by Bayer Healthcare LLC, owner of the registered trademark Aleve®

Distributed by:
Dr. Reddy's Laboratories, Inc.
Princeton, NJ 08540
Made in India

Rev: 03/2022



0 000000 000000 0

LOT

EXP

Dr.Reddy's

NDC 43598-071-01

HEADACHE PAIN

Naproxen Sodium Tablets USP, 220 mg (NSAID)

Pain reliever / fever reducer
STRENGTH TO LAST 12 HOURS
for temporary relief of headache pain

100 Tablets

TAMPER EVIDENT: Do not use if foil seal under cap printed with "sealed for your protection" is broken or missing.
IMPORTANT: Read the directions and warnings before use.
See Carton for Full Information

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, or blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use** if you have ever had an allergic reaction to any other pain reliever/fever reducer or right before or after heart surgery. **Ask a doctor before use if** the stomach bleeding warning applies to you, you have a history of stomach problems (such as heartburn), you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke, you are taking a diuretic, you have problems or serious side effects

Distributed by: Dr. Reddy's Laboratories, Inc.

Princeton, NJ 08540

Made in India

Rev. 03/2022

1500XXXXX

LOT

EXP

Peel Here

from taking pain relievers or fever reducers. **Ask a doctor or pharmacist before use if you are** under a doctor's care for any serious condition, or taking any other drug. **When using this product** take with food or milk if stomach upset occurs. **Stop use and ask a doctor if you** experience any of the following signs of stomach bleeding (feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better), you have symptoms of heart problems or stroke (chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling), pain gets worse or lasts more than 10 days, fever gets worse or lasts more than 3 days, you have difficulty swallowing, it feels like the pill is stuck in your throat, redness or swelling is present in the painful area, or any new symptoms appear. **If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). **Directions:** Do not take more than directed. The smallest effective dose should be used. Drink a full glass of water with each dose. Adults and children 12 years and older: take 1 tablet every 8 to 12 hours while symptoms last. For the first dose you may take 2 tablets within the first hour. Do not exceed 2 tablets in any 8- to 12-hour period. Do not exceed 3 tablets in a 24-hour period. Children under 12 years: ask a doctor. **Other information** Each tablet contains: sodium 20 mg. Store at 20°-25°C (68°-77°F). Avoid high humidity and excessive heat above 40°C (104°F). **Questions or comments?** call toll-free weekdays 9 AM to 8 PM EST at 1-888-375-3784

Headache Pain - Tablet Container Carton

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat

Drug Facts (continued)

- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and children

- 12 years and older
 - take 1 tablet every 8 to 12 hours while symptoms last
 - for the first dose you may take 2 tablets within the first hour
 - do not exceed 2 tablets in any 8- to 12-hour period
 - do not exceed 3 tablets in a 24-hour period

Children under 12 years

- ask a doctor

Other information

- each tablet contains: sodium 20 mg
- store at 20° - 25°C (68° - 77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients

FD & C Blue # 2, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, talc, titanium dioxide

Questions or comments?

call toll-free weekdays 9 AM to 8 PM EST at 1-888-375-3784

TAMPER EVIDENT: Do not use if foil seal under cap printed with "sealed for your protection" is broken or missing.

IMPORTANT: Read the directions and warnings before use. Keep the carton for reference, it contains important information.

**This product is not manufactured or distributed by Bayer Healthcare LLC, owner of the registered trademark Aleve®

NDC 44398-074-01

Compare to Aleve® Headache Pain Tablets active ingredient**

Dr. Reddy's

HEADACHE PAIN

Naproxen Sodium
Tablets USP, 220 mg (NSAID)

Pain reliever / fever reducer
for temporary relief of headache pain

100 Tablets

Drug Facts

Active ingredient (in each tablet)

Naproxen sodium USP, 220 mg
(naproxen USP, 200 mg) (NSAID) *.....Pain reliever/
fever reducer

Purposes

*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - minor pain of arthritis
 - muscular aches
 - backache
 - menstrual cramps
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Distributed by:
Dr. Reddy's Laboratories, Inc.
Princeton, NJ 08540
Made in India

Rev. 03/2022



LOT

EXP

Back and Muscle Pain - Caplet Container

Dr.Reddy's

NDC 43598-074-01

BACK and MUSCLE PAIN

Naproxen Sodium Tablets USP, 220 mg (NSAID)

Pain reliever / fever reducer

STRENGTH TO LAST 12 HOURS

for temporary relief of minor back and
muscle aches and pains

100 Caplets

(CAPSULE-SHAPED TABLETS)

TAMPER EVIDENT: Do not use if foil seal under cap printed with
"sealed for your protection" is broken or missing.

IMPORTANT: Read the directions and warnings before use.

See Carton for Full Information

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, or blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older

■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. **Heart attack and stroke warning:**

NSAIDs except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use** if you have ever had an allergic reaction to any other pain reliever/fever reducer or right before or after heart surgery. **Ask a doctor before use if** the stomach bleeding warning applies to you, you have a history of stomach problems (such as heartburn), you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke, you are taking a diuretic, you have problems or serious side effects

Distributed by: Dr. Reddy's Laboratories, Inc.

Princeton, NJ 08540

Made in India

Rev: 03/2022

1500XXXXX

LOT

EXP

Peel Here

from taking pain relievers or fever reducers. **Ask a doctor or pharmacist before use if you are** under a doctor's care for any serious condition, or taking any other drug. **When using this product** take with food or milk if stomach upset occurs. **Stop use and ask a doctor if you** experience any of the following signs of stomach bleeding (feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better), you have symptoms of heart problems or stroke (chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling), pain gets worse or lasts more than 10 days, fever gets worse or lasts more than 3 days, you have difficulty swallowing, it feels like the pill is stuck in your throat, redness or swelling is present in the painful area, or any new symptoms appear. **If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions Do not take more than directed. The smallest effective dose should be used. Drink a full glass of water with each dose. Adults and children 12 years and older: take 1 caplet every 8 to 12 hours while symptoms last. For the first dose you may take 2 caplets within the first hour. Do not exceed 2 caplets in any 8- to 12-hour period. Do not exceed 3 caplets in a 24-hour period. Children under 12 years: ask a doctor. **Other information** Each caplet contains: sodium 20 mg. Store at 20°-25°C (68°-77°F). Avoid high humidity and excessive heat above 40°C (104°F). **Questions or comments?** call toll-free weekdays 9 AM to 8 PM EST at 1-888-375-3784

Back & Muscle Pain - Caplet Container Carton

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- If you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat

Drug Facts (continued)

- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose

Adults and children 12 years and older

- take 1 caplet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 caplets within the first hour
- do not exceed 2 caplets in any 8- to 12-hour period
- do not exceed 3 caplets in a 24-hour period

Children under 12 years: ■ ask a doctor

Other information

- each caplet contains: sodium 20 mg
- store at 20° - 25°C (68° - 77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients

FD & C Blue #2, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, talc, titanium dioxide

Questions or comments?

call toll-free weekdays 9 AM to 8 PM EST at 1-888-375-3784

TAMPER EVIDENT: Do not use if foil seal under cap printed with "sealed for your protection" is broken or missing.

IMPORTANT: Read the directions and warnings before use. Keep the carton for reference, it contains important information.

*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Aleve®

NDC 43988-074-01

Compare to Aleve® Back and Muscle Pain Caplets active ingredient**



BACK and MUSCLE PAIN

Naproxen Sodium
Tablets USP, 220 mg (NSAID)

Pain reliever / fever reducer
for temporary relief of minor back
and muscle aches and pains

100 Caplets (CAPSULE-SHAPED TABLETS)

STRENGTH
TO LAST 12
HOURS

Drug Facts

Active Ingredient (In each caplet)

Naproxen sodium USP, 220 mg
(naproxen USP, 200 mg) (NSAID) *.....Pain reliever/
fever reducer

Purposes

Uses

- temporarily relieves minor aches and pains due to:
 - backache
 - minor pain of arthritis
 - headache
 - the common cold
 - temporarily reduces fever
 - muscular aches
 - menstrual cramps
 - toothache

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- asthma (wheezing)
- skin reddening
- facial swelling
- shock
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Distributed by:

Dr. Reddy's Laboratories, Inc.
Princeton, NJ 08540
Made in India

Rev: 03/2022



LOT

EXP

Back and Muscle Pain - Tablet Container

Dr.Reddy's

NDC 43598-073-01

BACK and MUSCLE PAIN

Naproxen Sodium Tablets USP, 220 mg (NSAID)

Pain reliever / fever reducer

STRENGTH TO LAST 12 HOURS

for temporary relief of minor back and
muscle aches and pains

100 Tablets

TAMPER EVIDENT: Do not use if foil seal under cap printed with
"sealed for your protection" is broken or missing.

IMPORTANT: Read the directions and warnings before use.

See Carton for Full Information

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, or blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older

■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs except aspirin increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use** if you have ever had an allergic reaction to any other pain reliever/fever reducer or right before or after heart surgery. **Ask a doctor before use** if the stomach bleeding warning applies to you, you have a history of stomach problems (such as heartburn), you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke, you are taking a diuretic, you have problems or serious side effects

Distributed by: Dr. Reddy's Laboratories, Inc.

Princeton, NJ 08540

Made in India

Rev: 03/2022

1500XXXXX

Peel Here

LOT

EXP


from taking pain relievers or fever reducers. **Ask a doctor or pharmacist before use if you are** under a doctor's care for any serious condition, or taking any other drug. **When using this product** take with food or milk if stomach upset occurs. **Stop use and ask a doctor if you** experience any of the following signs of stomach bleeding (feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better), you have symptoms of heart problems or stroke (chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling), pain gets worse or lasts more than 10 days, fever gets worse or lasts more than 3 days, you have difficulty swallowing, it feels like the pill is stuck in your throat, redness or swelling is present in the painful area, or any new symptoms appear. **If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions Do not take more than directed. The smallest effective dose should be used. Drink a full glass of water with each dose. Adults and children 12 years and older: take 1 tablet every 8 to 12 hours while symptoms last. For the first dose you may take 2 tablets within the first hour. Do not exceed 2 tablets in any 8- to 12-hour period. Do not exceed 3 tablets in a 24-hour period. Children under 12 years: ask a doctor.

Other information Each tablet contains: sodium 20 mg. Store at 20°-25°C (68°-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Questions or comments? call toll-free weekdays 9 AM to 8 PM EST at 1-888-375-3784

Back and Muscle Pain - Tablet Container Carton



NAPROXEN SODIUM HEADACHE PAIN

naproxen sodium tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:43598-071
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Naproxen Sodium (UNII: 9TN87S3A3C) (Naproxen - UNII:57Y76R9ATQ)	Naproxen Sodium	220 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
hypromelloses (UNII: 3NXW29V3WO)	
magnesium stearate (UNII: 70097M6I3O)	
cellulose, microcrystalline (UNII: OP1R32D61U)	
Polyethylene Glycol, Unspecified (UNII: 3WJQ0SDW1A)	
povidone (UNII: FZ989GH94E)	
talc (UNII: 7SEV7J4R1U)	
titanium dioxide (UNII: 15FIX9V2JP)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	

Product Characteristics

Color	WHITE	Score	no score
Shape	ROUND	Size	9mm
Flavor		Imprint Code	R;272
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:43598-071-01	1 in 1 CARTON	12/10/2021	
1		100 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075168	12/10/2021	

NAPROXEN SODIUM HEADACHE PAIN

naproxen sodium tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:43598-072
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Naproxen Sodium (UNII: 9TN87S3A3C) (Naproxen - UNII:57Y76R9ATQ)	Naproxen Sodium	220 mg

Inactive Ingredients				
Ingredient Name			Strength	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)				
hypromelloses (UNII: 3NXW29V3WO)				
magnesium stearate (UNII: 70097M6I30)				
cellulose, microcrystalline (UNII: OP1R32D61U)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
povidone (UNII: FZ989GH94E)				
talc (UNII: 7SEV7J4R1U)				
titanium dioxide (UNII: 15FIX9V2JP)				
POLYSORBATE 80 (UNII: 6OZP39ZG8H)				
Product Characteristics				
Color	WHITE	Score	no score	
Shape	CAPSULE	Size	12mm	
Flavor		Imprint Code	R;273	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:43598-072-01	1 in 1 CARTON	12/10/2021	
1		100 in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
ANDA	ANDA075168		12/10/2021	

NAPROXEN SODIUM BACK AND MUSCLE PAIN

naproxen sodium tablet, coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:43598-073
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
Naproxen Sodium (UNII: 9TN87S3A3C) (Naproxen - UNII:57Y76R9ATQ)		Naproxen Sodium	220 mg

Inactive Ingredients				
Ingredient Name			Strength	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)				
hypromelloses (UNII: 3NXW29V3WO)				
magnesium stearate (UNII: 70097M6I30)				
cellulose, microcrystalline (UNII: OP1R32D61U)				
Polyethylene Glycol, Unspecified (UNII: 3WJQ0SDW1A)				
povidone (UNII: FZ989GH94E)				
talc (UNII: 7SEV7J4R1U)				
titanium dioxide (UNII: 15FIX9V2JP)				
POLYSORBATE 80 (UNII: 6OZP39ZG8H)				
Product Characteristics				
Color		WHITE	Score	no score
Shape		ROUND	Size	9mm
Flavor			Imprint Code	R;272
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:43598-073-01	1 in 1 CARTON	12/10/2021	
1		100 in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA		ANDA075168	12/10/2021	

NAPROXEN SODIUM BACK AND MUSCLE PAIN

naproxen sodium tablet, coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:43598-074
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
Naproxen Sodium (UNII: 9TN87S3A3C) (Naproxen - UNII:57Y76R9ATQ)		Naproxen Sodium	220 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
hypromelloses (UNII: 3NXW29V3WO)	
magnesium stearate (UNII: 70097M6I30)	
cellulose, microcrystalline (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
povidone (UNII: FZ989GH94E)	
talc (UNII: 7SEV7J4R1U)	
titanium dioxide (UNII: 15FIX9V2JP)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	

Product Characteristics

Color	WHITE	Score	no score
Shape	CAPSULE	Size	12mm
Flavor		Imprint Code	R;273
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:43598-074-01	1 in 1 CARTON	12/10/2021	
1		100 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075168	12/10/2021	

Labeler - Dr.Reddy's Laboratories Inc (802315887)

Revised: 12/2022

Dr.Reddy's Laboratories Inc