

CHILDRENS SILAPAP- acetaminophen liquid
Lannett Company, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Children's Silapap Liquid

Active Ingredient: Acetaminophen 160 mg (in each 5 mL (TSP))

Purpose: Pain reliever/fever reducer

Uses To reduce fever and for the temporary relief of minor aches and pains due to:

- Headache
- Muscular aches
- Backache
- Minor pain of arthritis
- The common cold
- Toothache
- Premenstrual and menstrual cramps

Warnings

Liver Warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Alcohol warning: If the user consumes 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen or other pain relievers/fever reducers.

Do not use

- with any other product drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if the user is allergic to acetaminophen or any of the inactive ingredients in this product

When using this product

- **do not exceed recommended dose (see overdose warning)**

Stop use and ask a doctor if

- **new symptoms occur**
- redness or swelling is present
- pain gets worse or lasts for more than 5 days
- fever gets worse or lasts for more than 3 days
- These could be signs of a serious condition

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose Warning: In case of overdose, get medical help or contact a Poison Control Center (1800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs of symptoms.

Directions

- do not take more than directed (see overdose warning).
- if needed, repeat dose every 4 hours or as directed by a doctor
- do not give more than 5 doses in 24 hours

children under 2 yrs (under 24 lbs)	ask a doctor
children 2-3 years (24-35 lbs)	1 teaspoonful (TSP)(5 mL)
children 4-5 years (36-47 lbs)	1 1/2 teaspoonfuls (TSP)(7.5 mL)
children 6-8 years (48-59 lbs)	2 teaspoonfuls (TSP)(10 mL)
children 9-10 years (60-71 lbs)	2 1/2 teaspoonfuls (TSP)(12.5 mL)
children 11 years (72-95 lbs)	3 teaspoonfuls (TSP)(15 mL)
adults & children 12 years & older	4 teaspoonfuls (TSP)(20 mL)

Other information

Store at room temperature 20°-25°C (68°-77°F)

Inactive ingredients

citric acid, D&C red no. 33, FD&C red no. 40, cherry flavor, methylparaben, propylene glycol, saccharin sodium, sodium benzoate, and purified water.

Questions

888-974-5279

This product is not manufactured or distributed by McNeil Consumer & Specialty Pharmaceuticals, distributor of Tylenol®.

Manufactured by:

Silarx Pharmaceuticals, Inc
 1033 Stoneleigh Ave.
 Carmel, NY 10512
 USA



See New Warnings Information
 NDC 54838-144-70

Silapap

Acetaminophen Liquid

Acetaminophen 160 mg/5 mL
PAIN RELIEVER, FEVER REDUCER

- Alcohol Free
- Aspirin Free
- Sugar Free

DO NOT USE IF PRINTED
 SAFETY SEAL ON THE BOTTLE
 IS BROKEN OR MISSING

237 mL (8 fl oz)

Drug Facts

Active ingredient
 (in each 5 mL = 1 teaspoon (TSP))
 Acetaminophen 160 mg Pain reliever/fever reducer

Uses
 To reduce fever and for the temporary relief of minor aches and pains due to:
 ■ Headache ■ Muscular aches ■ Backache
 ■ Minor pain of arthritis ■ The common cold ■ Toothache
 ■ Premenstrual and menstrual cramps.

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 5 doses in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Alcohol warning: If the user consumes 3 or more alcoholic drinks every day, ask your doctor whether the user should take acetaminophen or other pain reliever/fever reducers.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If the user is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if the user has liver disease.
Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin.

When using this product ■ do not exceed recommended dose (see overbottle warning).

Stop use and ask a doctor if ■ new symptoms occur ■ redness or swelling is present ■ pain gets worse or lasts for more than 5 days ■ fever gets worse or lasts for more than 3 days ■ These could be signs of a serious condition.

Control # & Exp. Date



N 3 54838-144-70 3



Drug Facts Continued

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose Warning: In case of overdose, get medical help or contact a Poison Control Center (1800-222-1222) right away. Quick medical attention is critical, even if you do not notice any signs of symptoms.

Directions

- do not take more than directed (see overdose warning).
- if needed, repeat dose every 4 hours or as directed by a doctor
- do not give more than 5 doses in 24 hours

children under 2 yrs (Under 24 lbs)	ask a doctor
children 2-3 years (24-35 lbs)	1 teaspoon (TSP) (5 mL)
children 4-5 years (36-47 lbs)	1½ teaspoons (TSP) (7.5 mL)
children 6-8 years (48-59 lbs)	2 teaspoons (TSP) (10 mL)
children 9-10 years (60-71 lbs)	2½ teaspoons (TSP) (12.5 mL)
children 11 years (72-95 lbs)	3 teaspoons (TSP) (15 mL)
adults & children 12 years & older	4 teaspoons (TSP) (20 mL)

Other Information

Store at room temperature 20°-25°C (68°-77°F)

Inactive Ingredients

citric acid, D&C red no. 38, FD&C red no. 40, cherry flavor, methylparaben, propylene glycol, saccharin sodium, sodium benzoate, and purified water.

Questions

Manufactured by:
 Silarx Pharmaceuticals, Inc
 1033 Stoneleigh Ave
 Carmel, NY 10512
 Rev. 06/13

CHILDRENS SILAPAP
 acetaminophen liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:54838-144
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	CHERRY (cherry flavor)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54838-144-40	118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	09/05/1994	05/31/2024
2	NDC:54838-144-70	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	09/05/1994	01/31/2024
3	NDC:54838-144-80	473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	09/05/1994	05/31/2024

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	09/05/1994	05/31/2024

Labeler - Lannett Company, Inc. (002277481)

