

IBUPROFEN- ibuprofen tablet, film coated

Major Pharmaceuticals

Major Pharmaceuticals Ibuprofen Drug Facts

Active ingredient (in each tablet)

Ibuprofen 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- minor pain of arthritis
- toothache
- backache
- the common cold
- menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**

adults and children 12 years and older	<ul style="list-style-type: none"> • take 1 tablet every 4 to 6 hours while symptoms persist • if pain or fever does not respond to 1 tablet, 2 tablets may be used • do not exceed 6 tablets in 24 hours, unless directed by a doctor
children under 12 years	<ul style="list-style-type: none"> • ask a doctor

Other information

- read all warnings and directions before use
- store between 20-25°C (68-77°F)
- avoid high humidity and excessive heat above 40°C (104°F)

Inactive ingredients

colloidal silicon dioxide, corn starch, croscarmellose sodium, hypromellose, iron oxide red, iron oxide yellow, microcrystalline cellulose, polyethylene glycol, polysorbate 80, stearic acid, titanium dioxide

Questions or comments?

1-800-616-2471

Principal Display Panel

MAJOR[®]

SEE REVISED WARNING

Ibuprofen Tablets, 200 mg

PAIN RELIEVER

FEVER REDUCER

(NSAID)

COMPARE TO the active ingredient of ADVIL[®]

Actual Size

1000 COATED TABLETS

<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> you have symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling, pain gets worse or lasts more than 10 days, fever gets worse or lasts more than 3 days, redness or swelling is present in the painful area, any new symptoms appear <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p> <p>Warnings</p> <ul style="list-style-type: none"> Do not use if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer Do not use if you have a history of stomach problems, such as heartburn, liver disease, asthma, or had a stroke Do not use if you are taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin Do not use if you are under a doctor's care for any serious condition Do not use if you are taking any other drug <p>When using this product</p> <ul style="list-style-type: none"> Take with food or milk if stomach upset occurs Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better <p>Other information</p> <ul style="list-style-type: none"> Read all warnings and directions before use Store between 20-25°C (68-77°F) Avoid high humidity and excessive heat above 40°C (104°F) <p>Inactive ingredients: colloidal silicon dioxide, corn starch, croscarmellose sodium, hypromellose, iron oxide red, iron oxide yellow, microcrystalline cellulose, polyethylene glycol, polysorbate 80, stearic acid, titanium dioxide</p> <p>Questions or comments? 1-800-616-2471</p>	<p>Drug Facts (continued)</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p> <p>Directions</p> <ul style="list-style-type: none"> Do not take more than directed The smallest effective dose should be used <p>adults and children 12 years and older</p> <ul style="list-style-type: none"> Take 1 tablet every 4 to 6 hours while symptoms persist If pain or fever does not respond to 1 tablet, 2 tablets may be used Do not exceed 6 tablets in 24 hours, unless directed by a doctor ask a doctor <p>Children under 12 years</p> <p>Other information</p> <ul style="list-style-type: none"> read all warnings and directions before use store between 20-25°C (68-77°F) avoid high humidity and excessive heat above 40°C (104°F) <p>Inactive ingredients: colloidal silicon dioxide, corn starch, croscarmellose sodium, hypromellose, iron oxide red, iron oxide yellow, microcrystalline cellulose, polyethylene glycol, polysorbate 80, stearic acid, titanium dioxide</p> <p>Questions or comments? 1-800-616-2471</p>	<p>NDC 0904-6747-80</p> <p>MAJOR[®]</p> <p>SEE REVISED WARNING</p> <p>Ibuprofen Tablets, 200mg</p> <p>PAIN RELIEVER FEVER REDUCER (NSAID)</p> <p>COMPARE TO the active ingredient of ADVIL[®]</p> <p>1000 COATED TABLETS</p> <p>Actual Size</p>	<p>Drug Facts</p> <p>Active ingredient (in each tablet) Purposes Ibuprofen 200 mg (NSAID)*.....Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug</p> <p>Uses</p> <ul style="list-style-type: none"> temporarily relieves minor aches and pains due to: headache, muscular aches, minor pain of arthritis, toothache, backache, the common cold, menstrual cramps temporarily reduces fever <p>Warnings</p> <p>Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin redness, rash, blisters</p> <p>If an allergic reaction occurs, stop use and seek medical help right away.</p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 60 or older, have had stomach ulcers or bleeding problems, take a blood thinning (anticoagulant) or steroid drug, take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), have 3 or more alcoholic drinks every day while using this product, take more or for a longer time than directed</p>	<p>Drug Facts (continued)</p> <p>Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.</p> <p>Do not use if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer Do not use if you have a history of stomach problems, such as heartburn, liver disease, asthma, or had a stroke Do not use if you are taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin Do not use if you are under a doctor's care for any serious condition Do not use if you are taking any other drug <p>When using this product</p> <ul style="list-style-type: none"> Take with food or milk if stomach upset occurs Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better </p>
---	---	---	--	---

IBUPROFEN

ibuprofen tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0904-6747
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	

CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	BROWN	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	I2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0904-6747-51	1 in 1 CARTON	09/20/2018	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:0904-6747-59	1 in 1 CARTON	09/20/2018	
2		100 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:0904-6747-70	1 in 1 CARTON	09/20/2018	
3		250 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:0904-6747-40	500 in 1 BOTTLE; Type 0: Not a Combination Product	09/20/2018	
5	NDC:0904-6747-80	1000 in 1 BOTTLE; Type 0: Not a Combination Product	09/20/2018	
6	NDC:0904-6747-24	1 in 1 CARTON	09/20/2018	
6		24 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA072096	09/20/2018	

Labeler - Major Pharmaceuticals (191427277)

