

CHILDRENS MOTRIN- ibuprofen tablet, chewable
Johnson & Johnson Consumer Inc.

Children's Motrin

Drug Facts

Active ingredient (in each chewable tablet)

Ibuprofen 100 mg (NSAID) ¹

¹ nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

Warnings

Allergy alert

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin,

ibuprofen, naproxen, or others)

- takes more or for a longer time than directed

Heart attack and stroke warning

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Sore throat warning

Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly.

Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- mouth or throat burning may occur; give with food or water
- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
 - feels faint
 - vomits blood
 - has bloody or black stools
 - has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing

- weakness in one part or side of body
- slurred speech
- leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- **chew or crush tablets completely before swallowing**
- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- find right dose on chart below. If possible, use weight to dose; otherwise use age.
- if needed, repeat dose every **6-8 hours**
- do not use more than **4 times a day**

Dosing Chart

| Weight (lb) | Age (yr) | Tablets |
|--------------------|-----------------|----------------|
| under 24 | under 2 | ask a doctor |
| 24-35 | 2-3 | 1 |
| 36-47 | 4-5 | 1 ½ |
| 48-59 | 6-8 | 2 |
| 60-71 | 9-10 | 2 ½ |
| 72-95 | 11 | 3 |

Other information

- phenylketonurics: contains phenylalanine 6 mg per tablet
- store between 20-25°C (68-77°F)

Inactive ingredients

acesulfame potassium, ammonium glycyrrhizin, aspartame, carnauba wax, croscarmellose sodium, hypromellose, magnesium stearate, mannitol, natural and artificial flavors, silicon dioxide, sodium lauryl sulfate, soybean oil, succinic acid

Questions?

Call **1-877-895-3665** (toll-free) or **215-273-8755** (collect) or visit www.motrin.com

PRINCIPAL DISPLAY PANEL

NDC 50580-932-01

For Ages 2 to 11 Years

**Children's
Motrin®**

**Ibuprofen Chewable Tablets, 100 mg
Pain Reliever / Fever Reducer (NSAID)
Chewables**

DYE-FREE

Lasts up to 8 hours

Actual Size

Chew or crush tablets
completely before swallowing

**24
Grape-Flavored Chewable Tablets**



NDC 50580-932-01
For Ages 2 to 11 Years
**Children's
Motrin®**
Ibuprofen Chewable Tablets, 100 mg
Pain Reliever / Fever Reducer (NSAID)
Chewables
DYE-FREE
Lasts up to 8 hours
Actual Size
Chew or crush tablets
completely before swallowing
**24
Grape-Flavored Chewable Tablets**

Important: Read all product information before using. Keep this box for important information. This product is for children only.

Drug Facts
Active ingredient *Purpose*
(in each chewable tablet)
Ibuprofen 100 mg (NSAID).....Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses temporarily: ■ reduces fever ■ relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child: ■ has had stomach ulcers or bleeding problems ■ takes a blood thinning (anticoagulant) or steroid drug ■ takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ takes more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. ➔

Drug Facts (continued)
Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.
Do not use ■ if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery
Ask a doctor before use if ■ stomach bleeding warning applies to your child ■ child has a history of stomach problems, such as heartburn ■ child has problems or serious side effects from taking pain relievers or fever reducers ■ child has not been drinking fluids ■ child has lost a lot of fluid due to vomiting or diarrhea ■ child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke ■ child has asthma ■ child is taking a diuretic
Ask a doctor or pharmacist before use if the child is ■ under a doctor's care for any serious condition ■ taking any other drug
When using this product ■ mouth or throat burning may occur; give with food or water ■ take with food or milk if stomach upset occurs
Stop use and ask a doctor if ■ child experiences any of the following signs of stomach bleeding: ■ feels faint ■ vomits blood ■ has bloody or black stools ■ has stomach pain that does not get better ■ child has symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling ■ the child does not get any relief within first day (24 hours) of treatment ■ fever or pain gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear
Keep out of reach of children. In case of overdose, get medical ➔

Drug Facts (continued)
help or contact a Poison Control Center right away. (1-800-222-1222)

Directions ■ chew or crush tablets completely before swallowing ■ this product does not contain directions or complete warnings for adult use ■ do not give more than directed ■ find right dose on chart below. If possible, use weight to dose, otherwise use age. ■ if needed, repeat dose every 6-8 hours ■ do not use more than 4 times a day

Dosing Chart

| Weight (lb) | Age (yr) | Tablets |
|-------------|----------|--------------|
| under 24 | under 2 | ask a doctor |
| 24-35 | 2-3 | 1 |
| 36-47 | 4-5 | 1 ½ |
| 48-59 | 6-8 | 2 |
| 60-71 | 9-10 | 2 ½ |
| 72-95 | 11 | 3 |

Other information ■ phenylketonurics: contains phenylalanine 6 mg per tablet ■ store between 20-25°C (68-77°F)

Inactive ingredients: acesulfame potassium, ammonium glycyrrhizin, aspartame, carnauba wax, croscarmellose sodium, hypromellose, magnesium stearate, mannitol, natural and artificial flavors, silicon dioxide, sodium lauryl sulfate, soybean oil, succinic acid

Questions? Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect) or visit www.motrin.com

LOT
EXP
← OPEN HERE
3 0045-0932-26 6
2U262 MNC3

CHILDRENS MOTRIN

ibuprofen tablet, chewable

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:50580-932 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|--------------------------|-----------------|
| IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM) | IBUPROFEN | 100 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|---|-----------------|
| ACESULFAME POTASSIUM (UNII: 23OV73Q5G9) | |
| AMMONIUM GLYCYRRHIZATE (UNII: 3VRD35U26C) | |
| ASPARTAME (UNII: Z0H242BBR1) | |
| CARNAUBA WAX (UNII: R12CBM0EIZ) | |
| CROSCARMELOSE SODIUM (UNII: M28OL1HH48) | |
| HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO) | |
| MAGNESIUM STEARATE (UNII: 70097M6I30) | |
| MANNITOL (UNII: 3OWL53L36A) | |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4) | |
| SODIUM LAURYL SULFATE (UNII: 368GB5141J) | |
| SOYBEAN OIL (UNII: 241ATL177A) | |
| SUCCINIC ACID (UNII: AB6MNQ6J6L) | |

Product Characteristics

| | | | |
|-----------------|-------------------|---------------------|----------|
| Color | white (Off White) | Score | 2 pieces |
| Shape | ROUND | Size | 13mm |
| Flavor | GRAPE | Imprint Code | M;100 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|----------|------------------|---|-----------------------------|---------------------------|
| 1 | NDC:50580-932-01 | 1 in 1 CARTON | 06/17/2019 | |
| 1 | | 24 in 1 BOTTLE; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------------|---|-----------------------------|---------------------------|
| ANDA | ANDA076359 | 06/17/2019 | |

Labeler - Johnson & Johnson Consumer Inc. (878046358)

Revised: 3/2023

Johnson & Johnson Consumer Inc.