

**STOMACH RELIEF - MAXIMUM STRENGTH- bismuth subsalicylate liquid**  
**AptaPharma Inc.**

-----  
**Stomach Relief - Maximum Strength**

**Drug Facts**

**Active ingredient**  
**(in each 30 mL dose cup or 2 tablespoons)**

Bismuth subsalicylate 1050 mg

**Purposes**

Bismuth subsalicylate ..... Upset stomach reliever  
and antidiarrheal

**Uses** relieves ■ travelers' diarrhea ■ diarrhea  
■ upset stomach due to overindulgence of food and  
drink including: ■ heart burn ■ indigestion  
■ nausea ■ gas ■ belching

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are  
recovering from chicken pox or flu-like symptoms should not  
use this product. When using this product, if changes in  
behavior with nausea and vomiting occur, consult a doctor  
because these symptoms could be an early sign of Reye's  
syndrome, a rare but serious illness.

**Allergy alert: Contains salicylate. Do not take if you are**

■ allergic to salicylates (including aspirin)  
■ taking other salicylate products

**Do not use** if you have ■ an ulcer  
■ a bleeding problem ■ black or bloody stool

**Ask a doctor before use if you have**

■ fever ■ mucus in stool

**Ask a doctor or pharmacist before use if you are**  
taking any drug for ■ anticoagulation (thinning of the blood)  
■ diabetes ■ gout ■ arthritis

**When using this product** a temporary, but harmless,  
darkening of the stool and/or tongue may occur

**Stop use and ask a doctor if**

■ diarrhea lasts more than 2 days  
■ symptoms get worse or last more than 2 days  
■ ringing in the ears or loss of hearing occurs

**If pregnant or breast feeding,** ask health professional  
before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

- Directions** ■ Shake well before use
- use dose cup or tablespoon (TBSP)
  - adults and children 12 years and over: 1 dose (30 mL or 2 TBSP) every 1 hour as needed
  - do not exceed 4 doses (120 mL or 8 TBSP) in 24 hours
  - use until diarrhea stops but not more than 2 days
  - children under 12 years: ask a doctor
  - drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Other information**

- **each 30 ml dose cup contains:**  
sodium 13 mg, salicylate 455 mg
- protect from freezing ■ avoid excessive heat (over 104°F or 40°C)
  - low sodium

**Inactive ingredients** benzoic acid, D&C red #22, D&C red # 28, flavor, purified water, saccharin sodium, salicylic acid, sodium salicylate, xanthan gum

**Questions?** 1-877-798-5944

**Principal Display Panel**

**AP SAFE** NDC 76281-538-26

**\*COMPARE TO  
the active ingredient in  
PEPTO-BISMOL™  
MAXIMUM STRENGTH**

**Stomach  
Relief  
Bismuth Subsalicylate  
Antidiarrheal / Upset Stomach Reliever**

**Maximum Strength**

**5 Symptom Relief of:**  
● Nausea ● Heartburn ● Indigestion  
● Upset stomach ● Diarrhea

**8 FL OZ (237 mL)**

**TAMPER EVIDENT: Do not use if imprinted  
shrinkband is missing or broken**

\*This product is not manufactured or distributed by Procter & Gamble, Inc., the distributor of Pepto-Bismol™.

**Manufactured by: AptaPharma Inc.,**

**Made in USA**

1533 Union Ave.  
Pennsauken, NJ 08110

3 76281 53826 7

AP-LR-12 REV01

PEEL  
HERE

<b>TAMPER EVIDENT: Do not use if imprinted shrinkband is missing or broken</b>	
<b>Drug Facts</b> <b>Active ingredient</b> <i>(In each 30 mL dose cup or 2 tablespoons)</i> Bismuth subsalicylate 1050 mg...Upset stomach reliever and antidiarrheal	<b>Purposes</b> Upset stomach reliever and antidiarrheal
<b>Uses</b> relieves ■ travelers' diarrhea ■ diarrhea ■ upset stomach due to overindulgence of food and drink including: ■ heart burn ■ indigestion ■ nausea ■ gas ■ belching	
<b>Warnings</b> <b>Reye's syndrome:</b> Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. <b>Allergy alert: Contains salicylate. Do not take if you are</b> ■ allergic to salicylates (including aspirin) ■ taking other salicylate products <b>Do not use if you have</b> ■ an ulcer ■ a bleeding problem ■ black or bloody stool <b>Ask a doctor before use if you have</b> ■ fever ■ mucus in stool	



NDC 76281-538-26

**\*COMPARE TO the active ingredient in PEPTO-BISMOL™ MAXIMUM STRENGTH**

# Stomach Relief

Bismuth Subsalicylate  
Antidiarrheal / Upset Stomach Reliever

**Maximum Strength**

**5 Symptom Relief of:**  
• Nausea • Heartburn • Indigestion  
• Upset stomach • Diarrhea

8 FL OZ (237 mL)

<b>Drug Facts (continued)</b> <b>Ask a doctor or pharmacist before use if you are taking any drug for</b> ■ anticoagulation (thinning of the blood) ■ diabetes ■ gout ■ arthritis <b>When using this product</b> a temporary, but harmless, darkening of the stool and/or tongue may occur <b>Stop use and ask a doctor if</b> ■ diarrhea lasts more than 2 days ■ symptoms get worse or last more than 2 days ■ ringing in the ears or loss of hearing occurs <b>If pregnant or breast feeding, ask health professional before use.</b> <b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.
--

\* This product is not manufactured or distributed by Procter & Gamble, Inc., the distributor of Pepto-Bismol™.

Manufactured by: AptaPharma Inc.,  
1533 Union Ave.  
Pennsauken, NJ 08110

Made in USA  
AP-LR-12 REV01



3 76281 53826 7 PEEL HERE

<b>Drug Facts (continued)</b> <b>Directions</b> ■ Shake well before use ■ use dose cup or tablespoon (TBSP) ■ adults and children 12 years and over: 1 dose (30 mL or 2 TBSP) every 1 hour as needed ■ do not exceed 4 doses (120 mL or 8 TBSP) in 24 hours ■ use until diarrhea stops but not more than 2 days ■ children under 12 years: ask a doctor ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea
<b>Other information</b> ■ each 30 mL dose cup contains: sodium 13 mg, salicylate 455 mg ■ protect from freezing ■ avoid excessive heat (over 104°F or 40°C) ■ low sodium
<b>Inactive ingredients</b> benzoic acid, D&C red # 22, D&C red # 28, flavor, purified water, saccharin sodium, salicylic acid, sodium salicylate, xanthan gum
<b>Questions?</b> 1-877-798-5944

Stomach Relief - Maximum Strength by AptaPharma Inc.

res

## STOMACH RELIEF - MAXIMUM STRENGTH

bismuth subsalicylate liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:76281-538
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BISMUTH SUBSALICYLATE</b> (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	1050 mg in 30 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>BENZOIC ACID</b> (UNII: 8SKN0B0MIM)	
<b>D&amp;C RED NO. 22</b> (UNII: 1678RKX8RT)	
<b>D&amp;C RED NO. 28</b> (UNII: 7671P0Y5NH)	
<b>WATER</b> (UNII: 059QF0KOOR)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SALICYLIC ACID</b> (UNII: O414PZ4LPZ)	
<b>SODIUM SALICYLATE</b> (UNII: WQ1H85SYP)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

### Product Characteristics

<b>Color</b>	pink (color suspension)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	MINT	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:76281-538-26	237 mL in 1 BOTTLE; Type 0: Not a Combination Product	10/01/2018	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	10/01/2018	

**Labeler** - AptaPharma Inc. (790523323)

**Registrant** - AptaPharma Inc. (790523323)

### Establishment

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

AptaPharma Inc.		790523323	manufacture(76281-538)
-----------------	--	-----------	------------------------

Revised: 12/2023

AptaPharma Inc.