

**BISMUTH SUBSALICYLATE- bismuth subsalicylate capsule, liquid filled  
MEIJER DISTRIBUTION, INC.**

-----

**Stomach Relief**

**Drug Facts**

**Active ingredient (in each softgel)**  
**Bismuth subsalicylate 262 mg**

**Purpose**  
**Upset stomach reliever and anti-diarrheal**

**Uses**  
**relieves**

- **travelers' diarrhea • diarrhea**
- **upset stomach due to overindulgence in food and drink, including:**
- **heartburn • indigestion • nausea • gas • belching • fullness**

**Warnings**  
**Reye's syndrome:**Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product.  
When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
**Allergy alert:** Contains salicylate. Do not take if you are

- **allergic to salicylates (including aspirin)**
- **taking other salicylate products**

**Do not Use if you have**

- **an ulcer**
- **a bleeding problem**
- **bloody or black stool**

**Ask a doctor before use if you have**

- **fever • mucus in the stool**

**Ask a doctor or pharmacist before use if you are taking any drug for**

- **anticoagulation (thinning the blood) • diabetes • gout • arthritis**

**When using this product a temporary, but harmless, darkening of the stool**

**and/or tongue may occur**

**Stop use and ask a doctor if**

- **symptoms get worse or last more than 2 days**
- **ringing in the ears or loss of hearing occurs**
- **diarrhea lasts more than 2 days**

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.**

#### **Directions**

- **swallow with water, do not chew**
- **adults and children 12 years and over:**
- **2 softgels every ½ hour or 4 softgels every hour as needed for diarrhea**
- **2 softgels every ½ hour to 1 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)**
- **do not exceed 8 doses (16 softgels) in 24 hours**
- **use until diarrhea stops but not more than 2 days**
- **children under 12 years: ask a doctor**
- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**

#### **Other Information**

**each softgel contains: salicylate 102 mg**

- **very low sodium**
- **do not exceed 25°C**

#### **Inactive ingredients**

**aerosil, beeswax, D&C Red#33, FD&C Yellow-6, gelatin, glycerin, light liquid paraffin, povidone, purified water, sodium carboxy methyl cellulose, sorbitol sorbitan solution, soya lecithin, titanium dioxide**

**Questions or comments? 1-888-333-9792**

**Principle Display Panel**



DIST. BY MEIJER  
6 RAND APDS, MI 49644  
www.meijer.com  
Mazinda  
ORIS 1/24

**Drug Facts**

**Active ingredient (in each softgel)**  
Bismuth subsalicylate 262 mg

**Purpose**  
Upset stomach relief and anti-diarrhea

**Uses**  
Relieves heartburn, indigestion, upset stomach, and diarrhea.

**Warnings**  
Do not use if you have a bleeding problem or black stools. Do not use if you have a fever or chills. Do not use if you have a sore throat or difficulty swallowing. Do not use if you have a rash or hives. Do not use if you have a history of stomach ulcers or bleeding. Do not use if you have a history of kidney disease. Do not use if you have a history of liver disease. Do not use if you have a history of heart disease. Do not use if you have a history of high blood pressure. Do not use if you have a history of diabetes. Do not use if you have a history of asthma. Do not use if you have a history of epilepsy. Do not use if you have a history of alcoholism. Do not use if you have a history of drug abuse. Do not use if you have a history of mental illness. Do not use if you have a history of pregnancy or breastfeeding. Do not use if you are taking other medications, especially aspirin, ibuprofen, or other NSAIDs. Do not use if you are taking other medications, especially aspirin, ibuprofen, or other NSAIDs. Do not use if you are taking other medications, especially aspirin, ibuprofen, or other NSAIDs.

**Directions**  
Swallow with water, 1 softgel every 2 hours as needed for upset stomach or diarrhea. Do not exceed 12 softgels in 24 hours.

**Other information**  
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Drug Facts (continued)**  
Contains 24 softgels. Each softgel contains 262 mg of bismuth subsalicylate. Do not use if you have a history of stomach ulcers or bleeding. Do not use if you have a history of kidney disease. Do not use if you have a history of liver disease. Do not use if you have a history of heart disease. Do not use if you have a history of high blood pressure. Do not use if you have a history of diabetes. Do not use if you have a history of asthma. Do not use if you have a history of epilepsy. Do not use if you have a history of alcoholism. Do not use if you have a history of drug abuse. Do not use if you have a history of mental illness. Do not use if you have a history of pregnancy or breastfeeding. Do not use if you are taking other medications, especially aspirin, ibuprofen, or other NSAIDs. Do not use if you are taking other medications, especially aspirin, ibuprofen, or other NSAIDs. Do not use if you are taking other medications, especially aspirin, ibuprofen, or other NSAIDs.

**Questions or comments?** 1-800-333-9922

**Caution:** This product is not manufactured or distributed by Procter & Gamble, owner of the trademarks Pepto-Bismol and Liquid Caps.

# meijer Stomach Relief

BISMUTH SUBSALICYLATE 262 mg / UPSET STOMACH RELIEVER • ANTIDIARRHEAL

# meijer Stomach Relief

5 SYMPTOM DIGESTIVE RELIEF

- HEARTBURN • INDIGESTION • UPSET STOMACH
- NAUSEA • DIARRHEA

BISMUTH SUBSALICYLATE 262 mg  
UPSET STOMACH RELIEVER • ANTIDIARRHEAL

COMPARE TO  
PEPTO-BISMOL®  
LIQUICAPS™  
ACTIVE INGREDIENT\*

24  
SOFTGELS



Actual Size

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BUSTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

LOT NO:  
EXP DATE

# meijer Stomach Relief

BISMUTH SUBSALICYLATE 262 mg / UPSET STOMACH RELIEVER • ANTIDIARRHEAL

BISMUTH SUBSALICYLATE

bismuth subsalicylate capsule, liquid filled

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:79481-0057
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name		Basis of Strength Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ)		BISMUTH SUBSALICYLATE 262 mg

Inactive Ingredients	
Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SYNTHETIC BEESWAX (UNII: 08MNR5YE2R)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
PARAFFIN (UNII: I9O0E3H2ZE)	
POVIDONE (UNII: FZ989GH94E)	
WATER (UNII: 059QF0KO0R)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics			
Color	pink	Score	no score
Shape	OVAL	Size	21mm
Flavor		Imprint Code	262
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79481-0057-2	2 in 1 CARTON	02/01/2024	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	02/01/2024	

**Labeler -** MEIJER DISTRIBUTION, INC. (006959555)

Revised: 4/2024

MEIJER DISTRIBUTION, INC.