# DG HEALTH MILK OF MAGNESIA- magnesium hydroxide suspension Dolgencorp, LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

## **Dolgencorp, LLC Milk of Magnesia Drug Facts**

# Active ingredient (in each 15 mL)

Magnesium hydroxide 1200 mg

# Purpose

Saline laxative

#### Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in ½ to 6 hours

# Warnings

# Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

# Ask a doctor or pharmacist before use if you are

taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

# Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

# If pregnant or breast-feeding,

ask a health professional before use.

# Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

#### Directions

• do not exceed the maximum recommended daily dose in a 24 hour period

- shake well before use
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided
- mL = milliliter

adults and children 12 years and older	30 mL to 60 mL
children 6 to 11 years	15 mL to 30 mL
children under 6 years	ask a doctor

#### Other information

- each 15 mL contains: calcium 20 mg and magnesium 595 mg
- does not meet USP requirements for preservative effectiveness
- store at 20°-25°C (68°-77°F)
- close cap tightly after use
- avoid freezing

# **Inactive ingredient**

purified water

## Questions or comments?

1-888-309-9030

# **Principal Display Panel**

Compare to the active ingredient of Phillips'® Milk of Magnesia

Milk of Magnesia

Magnesium Hydroxide 1200 mg per 15 mL

Saline Laxative

Comfortable, cramp free relief

Sugar free

Stimulant free

Original Flavor

12 FL OZ (355 mL)

1200 mg per 15 mL



Compare to the active ingredient of Phillips'® Milk of Magnesia\*

# Milk of Magnesia

Magnesium Hydroxide 1200 mg per 15 mL Saline Laxative

- Comfortable, cramp free relief
- Sugar free
- Stimulant free

Original Flavor

12 FL OZ (355 mL)

: 39640 VT F3 1

mg per 15 mL

# Drug Facts

#### DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING.

Active ingredient (in each 15 mL)

Magnesium hydroxide 1200 mg.....

Purpose Saline laxative

Uses ■ relieves occasional constipation (irregularity)
■ generally produces bowel movement in ½ to 6 hours

#### Warnings

Ask a doctor before use if you have

■ kidney disease
 ■ a magnesium-restricted diet
 ■ stomach pain, nausea, or vomiting
 ■ a sudden change in bowel habits that lasts more than 2 weeks

Ask a doctor or pharmacist before use if you are taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.

■ you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a
Poison Control Center right away (1-800-222-1222).

#### Directions

- do not exceed the maximum recommended daily dose in a 24 hour period
- shake well before use
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided
   mL = milliliter

adults and children 12 years and older	30 mL to 60 mL
children 6 to 11 years	15 mL to 30 mL
children under 6 years	ask a doctor

#### Other information

- each 15 mL contains: calcium 20 mg and magnesium 595 mg
- does not meet USP requirements for preservative effectiveness
- store at 20°-25°C (68°-77°F) close cap tightly after use avoid freezing

Inactive ingredient purified water

Questions or comments? 1-888-309-9030

DISTRIBUTED BY DOLGENCORP, LLC

TN 37072 USA

100 MISSION RIDGE GOODLETTSVILLE,

\*This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Phillips'® Milk of Magnesia.

: 39640 VT B3

100% Satisfaction Guaranteed! (888) 309-9030



#### DG HEALTH MILK OF MAGNESIA

magnesium hydroxide suspension

#### **Product Information**

Product Type HUMAN OTC DRUG Item Code (Source) NDC:55910-396

Route of Administration ORAL

#### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>MAGNESIUM HYDRO XIDE</b> (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6 V3LHY838, HYDRO XIDE ION - UNII:9159 UV381P)	MAGNESIUM HYDRO XIDE	1200 mg in 15 mL

Inactive Ingredients	
Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Product Characteristics			
Color	WHITE (opaque)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

ı	Packaging				
ı	#	Item Code	Package Description	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ı	1 N	DC:55910-396-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	05/07/2014	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	05/07/2014	

# Labeler - Dolgencorp, LLC (068331990)

Revised: 11/2020 Dolgencorp, LLC