# MILK OF MAGNESIA- magnesium hydroxide suspension Meijer Distribution Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

# Meijer Distribution, Inc. Milk of Magnesia Drug Facts

# Active ingredient (in each 15 mL)

Magnesium hydroxide 1200 mg

# Purpose

Saline laxative

### Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in ½ to 6 hours

# Warnings

# Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- · stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

# Ask a doctor or pharmacist before use if you are

taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

# Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

# If pregnant or breast-feeding,

ask a health professional before use.

# Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### Directions

• do not exceed the maximum recommended daily dose in a 24 hour period

- shake well before use
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided
- mL = milliliter

adults and children 12 years and older	30 mL to 60 mL
children 6 to 11 years	15 mL to 30 mL
children under 6 years	ask a doctor

## Other information

- each 15 mL contains: calcium 20 mg, magnesium 595 mg, potassium 2 mg and sodium 3 mg
- does not meet USP requirements for preservative effectiveness
- store at 20-25°C (68-77°F)
- close cap tightly after use
- avoid freezing

# **Inactive ingredients**

anhydrous citric acid, D&C red #28, flavor, glycerin, purified water, sorbitol solution, sucrose, xanthan gum

# Questions or comments?

1-800-719-9260

# **Principal Display Panel**

Compare to Phillips'® active ingredient

#1 DOCTOR RECOMMENDED saline laxative active ingredient

**CRAMP FREE** 

milk of magnesia

Magnesium Hydroxide, 1200 mg per 15 mL

Saline Laxative

Gentle, Dependable Overnight Relief

Wild Cherry Flavor

ALCOHOL 0.4% | Stimulant Free

26 FL OZ (1 pt 10 OZ) 769 mL

NDC 41250-949-35

meijer

Compare to Phillips'® active ingredient\*



**CRAMP FREE** 

# milk of magnesia

Magnesium Hydroxide, 1200 mg per 15 mL Saline Laxative

Gentle, Dependable Overnight Relief



# Wild Cherry Flavor

**ALCOHOL 0.4% | Stimulant Free** 

26 FL 0Z (1 pt 10 0Z) 769 mL

: 94935 bE F10

# **Drug Facts**

### DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING.

# Active ingredient (in each 15 mL)

Purpose

- **Uses** relieves occasional constipation (irregularity)
- generally produces bowel movement in ½ to 6 hours

# Warnings

### Ask a doctor before use if you have

- kidney disease a magnesium-restricted diet stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

# Ask a doctor or pharmacist before use if you are taking any other drug.

Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

# Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

# If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

## Directions

- do not exceed the maximum recommended daily dose in a 24 hour period
- shake well before use
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided
- mL = milliliter

adults and children 12 years and older	30 mL to 60 mL
children 6 to 11 years	15 mL to 30 mL
children under 6 years	ask a doctor

## Other information

- each 15 mL contains: calcium 20 mg, magnesium 595 mg, potassium 2 mg and sodium 3 mg
- does not meet USP requirements for preservative effectiveness
- store at 20-25°C (68-77°F)
   close cap tightly after use
   avoid freezing

**Inactive ingredients** anhydrous citric acid, D&C red #28, flavor, glycerin, purified water, sorbitol solution, sucrose, xanthan gum

## Questions or comments? 1-800-719-9260

\*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Phillips'®.

\*\*In the U.S. Source: AlphalmpactRx

DIST. BY MEIJER DISTRIBUTION, INC. GRAND RAPIDS, MI 49544 www.meijer.com





PID 455162

# MILK OF MAGNESIA

magnesium hydroxide suspension

Pro	duct	Infor	mation
110	uuci	TIHUI	mauvn

Product Type HUMAN OTC DRUG Item Code (Source) NDC:41250-949

Route of Administration ORAL

# **Active Ingredient/Active Moiety**

Ingredient Name

Basis of Strength

Strength

BASIS OF STRENGTH ST

**MAGNESIUM HYDRO XIDE** (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6 V3LHY838, HYDRO XIDE ION - UNII:9159 UV381P)

MAGNESIUM HYDRO XIDE 1200 mg in 15 mL

# **Inactive Ingredients**

Ingredient Name	Strength	
ANHYDRO US CITRIC ACID (UNII: XF417D3PSL)		
GLYCERIN (UNII: PDC6A3C0OX)		
WATER (UNII: 059QF0KO0R)		
SUCROSE (UNII: C151H8M554)		
SORBITOL (UNII: 506T60A25R)		
XANTHAN GUM (UNII: TTV12P4NEE)		
<b>D&amp;C RED NO. 28</b> (UNII: 767IP0 Y5NH)		

## **Product Characteristics**

1 Todaet Characteristics		
Color	PINK (Opaque Light pink)	Score
Shape		Size
Flavor	CHERRY (wild cherry)	Imprint Code
Contains		

Packaging

П				
	# Item Code	Package Description	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
l	1 NDC:41250-949-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	11/13/2004	12/13/2016
ı	2 NDC:41250-949-35	769 mL in 1 BOTTLE; Type 0: Not a Combination Product	10/18/2004	

# **Marketing Information**

Warketing information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	10/18/2004	

Revised: 11/2018 Meijer Distribution Inc