

MILK OF MAGNESIA- magnesium hydroxide liquid
Meijer Distribution Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Meijer Distribution, Inc. Milk of Magnesia Drug Facts

Active ingredient (in each 15 mL Tablespoon)

Magnesium hydroxide 1200 mg

Purpose

Saline laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in ½ to 6 hours

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

Ask a doctor or pharmacist before use if you are

taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- do not exceed the maximum recommended daily dose in a 24 hour period

- shake well before use
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided
- mL = milliliter; TBSP = Tablespoon

adults and children 12 years and older	30 mL (2 TBSP) to 60 mL (4 TBSP)
children 6 to 11 years	15 mL (1 TBSP) to 30 mL (2 TBSP)
children under 6 years	ask a doctor

Other information

- each tablespoon (15 mL) contains: calcium 20 mg, magnesium 595 mg, potassium 2 mg and sodium 3 mg
- does not meet USP requirements for preservative effectiveness
- store at 20-25°C (68-77°F)
- close cap tightly after use
- avoid freezing

Inactive ingredients

anhydrous citric acid, D&C red #28, flavor, glycerin, purified water, sorbitol solution, sucrose, xanthan gum

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to Phillips'® active ingredient

Genuine

Saline Laxative

Milk of Magnesia

Magnesium Hydroxide 1200mg

Wild Cherry Flavor

Comfortable, Cramp Free Relief

Stimulant Free

#1 DOCTOR RECOMMENDED saline laxative active ingredient

ALCOHOL 0.4%

12 FL OZ (355 mL)

NDC 41250-949-40

meijerTM

Compare to
Phillips'[®]
active ingredient*

Genuine
Saline Laxative

Milkof Magnesia

Magnesium Hydroxide 1200 mg

Wild Cherry Flavor

Comfortable, Cramp Free Relief
Stimulant Free



ALCOHOL 0.4%



12 FL OZ (355 mL)

: 94940 6E F8

Drug Facts

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING.

Active ingredient (in each 15 mL Tablespoon)

Magnesium hydroxide 1200 mg.....Saline laxative

Purpose

- Uses**
- relieves occasional constipation (irregularity)
 - generally produces bowel movement in ½ to 6 hours

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

Ask a doctor or pharmacist before use if you are taking any other drug.

Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.

- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not exceed the maximum recommended daily dose in a 24 hour period
- shake well before use
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided
- mL = milliliter; TBSP = Tablespoon

adults and children 12 years and older	30 mL (2 TBSP) to 60 mL (4 TBSP)
children 6 to 11 years	15 mL (1 TBSP) to 30 mL (2 TBSP)
children under 6 years	ask a doctor

Other information

- each tablespoon (15 mL) contains: calcium 20 mg, magnesium 595 mg, potassium 2 mg and sodium 3 mg
- does not meet USP requirements for preservative effectiveness
- store at 20-25°C (68-77°F)
- close cap tightly after use
- avoid freezing

Inactive ingredients anhydrous citric acid, D&C red #28, flavor, glycerin, purified water, sorbitol solution, sucrose, xanthan gum

Questions or comments? 1-800-719-9260

0h6hb :

*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Phillips'®.

**In the U.S. Source: Symphony Health Solutions.

LE 89

DIST. BY MEIJER
DISTRIBUTION, INC.
GRAND RAPIDS, MI 49544
www.meijer.com



MILK OF MAGNESIA

magnesium hydroxide liquid

Product Information

Product Type	HUMAN OTC DRUG LABEL	Item Code (Source)	NDC:41250-949
Route of Administration	ORAL	DEA Schedule	

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MAGNESIUM HYDRO XIDE (MAGNESIUM CATION and HYDROXIDE ION)	MAGNESIUM HYDROXIDE	1200 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID	
GLYCERIN	
WATER	
SORBITOL	
SUCROSE	
XANTHAN GUM	
D&C RED NO. 28	

Product Characteristics

Color	PINK (Opaque Light pink)	Score	
Shape		Size	
Flavor	CHERRY (wild cherry)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-949-40	355 mL in 1 BOTTLE		
2	NDC:41250-949-35	769 mL in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	10/18/2004	

Labeler - Meijer Distribution Inc (006959555)

Revised: 2/2015

Meijer Distribution Inc