

GAVILAX - polyethylene glycol 3350 powder, for solution
Lupin Pharmaceuticals, Inc.

Drug Facts

Active Ingredient

Polyethylene Glycol 3350, 17 g (cap filled to line).....Osmotic Laxative

Purpose

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

WARNINGS

Allergy alert: Do not use if you are allergic to polyethylene glycol

Do not use if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are taking a prescription drug

When using this product you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of the reach of children

In case of overdose, get medical help or contact a POISON CONTROL CENTER right away. (1-800-222-1222)

Directions

- **do not take more than directed unless advised by your doctor**
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line.
- adults and children 17 years of age and older:
 - use once a day
 - fill to top of line in cap which is marked to indicate the correct dose (17 g)
 - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
 - do not combine with starch-based thickeners used for difficulty swallowing
 - ensure that the powder is fully dissolved before drinking
 - do not drink if there are any clumps
 - do not use more than 7 days
- children 16 years of age or under: ask a doctor

Other Information

- store at 20°- 25°C (68°- 77°F)
- tamper-evident: do not use if printed foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken

Inactive Ingredient

none

Questions or comments?

1-866-403-7592

- Dissolves in any beverages
- Sugar Free

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

NDC 43386-312-08

Original Prescription Strength

GaviLax

Polyethylene Glycol 3350

Powder for Oral Solution, Osmotic Laxative

14 ONCE-DAILY DOSES

NET WT 8.3 OZ (238 g)

Compare to
MiraLAX™

NDC 43386-312-08

Original Prescription Strength

GaviLAX

**Polyethylene Glycol 3350
Powder for Oral Solution,
Osmotic Laxative**

- Relieves Occasional Constipation (irregularity)
- Softens Stool



LUPIN

14 ONCE-DAILY DOSES
NET WT 8.3 OZ (238 g)

3/8" White Space Area

Unvarnished area

*MiraLAX™ is a registered trademark of Bayer Healthcare LLC

Manufactured for:
Lupin Pharmaceuticals, Inc.
Baltimore, MD 21202

SAP Code: 268477 07/2021



3/8" White Space Area

Drug Facts

Active ingredient (in each dose) Purpose

Polyethylene Glycol 3350,
17 g (cap filled to line).....Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol
Do not use if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting, or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are taking a prescription drug

When using this product you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition
- you get diarrhea
- you need to use a laxative for longer than 1 week

Polyethylene Glycol 3350 increases frequency of bowel movements and softens the stool.

Dissolves in any beverage
Sugar Free

Peel Here

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a POISON CONTROL CENTER right away (1-800-222-1222).

Directions

- do not take more than directed unless advised by your doctor
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line
- adults and children 17 years of age and older:
 - use once a day
 - fill to top of the line in cap which is marked to indicate the correct dose (17 g)
 - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
 - do not combine with starch-based thickeners used for difficulty swallowing
 - ensure that the powder is fully dissolved before drinking
 - do not drink if there are any clumps
 - do not use more than 7 days
- Children 16 years of age or under: ask a doctor

Other information

- store at 20°-25°C (68°-77°F)
- tamper-evident: do not use if foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken

Inactive ingredients none

Questions or comments? 1-866-403-7592

NDC 43386-312-85

Original Prescription Strength

GaviLax

Polyethylene Glycol 3350

Powder for Oral Solution, Osmotic Laxative

50 ONCE-DAILY DOSES

NET WT 29.9 OZ (850 g)

Compare to MiraLAX®

NDC 43386-312-85

Original Prescription Strength

GaviLAX

Polyethylene Glycol 3350 Powder for Oral Solution, Osmotic Laxative

- Relieves Occasional Constipation (irregularity)
- Softens Stool

LUPIN 50 ONCE-DAILY DOSES
NET WT 29.9 OZ (850 g)

Unvarnished Area

*MiraLAX® is a registered trademark of Bayer Healthcare LLC

Manufactured for:
Lupin Pharmaceuticals, Inc.
Baltimore, MD 21202

SAP Code: 268479
Rev. 07/2021

6 White Space Area

6 White Space Area

Drug Facts	
Active ingredient (in each dose) Polyethylene Glycol 3350, 17 g (cap filled to line)	Purpose Osmotic Laxative
Use <ul style="list-style-type: none"> relieves occasional constipation (irregularity) generally produces a bowel movement in 1 to 3 days 	
Warnings Allergy alert: Do not use if you are allergic to polyethylene glycol. Do not use if you have kidney disease, except under the advice and supervision of a doctor. Ask a doctor before use if you have <ul style="list-style-type: none"> nausea, vomiting, or abdominal pain a sudden change in bowel habits that lasts over 2 weeks irritable bowel syndrome Ask a doctor or pharmacist before use if you are taking a prescription drug. When using this product you may have loose, watery, more frequent stools. Stop use and ask a doctor if <ul style="list-style-type: none"> you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition. you get diarrhea you need to use a laxative for longer than 1 week If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a POISON CONTROL CENTER right away (1-800-222-1222).	
Directions <ul style="list-style-type: none"> do not take more than directed unless advised by your doctor the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line adults and children 17 years of age and older: <ul style="list-style-type: none"> use once a day fill to top of the line in cap which is marked to indicate the correct dose (17 g) stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink do not combine with starch-based thickeners used for difficulty swallowing ensure that the powder is fully dissolved before drinking do not drink if there are any clumps do not use more than 7 days children 16 years of age or under: ask a doctor 	
Other information <ul style="list-style-type: none"> store at 20°-25°C (68°-77°F) tamper-evident: do not use if foil seal under cap, printed with "SEALED FOR YOUR PROTECTION" is missing, open or broken 	
Inactive ingredients none	
Questions or comments? 1-866-403-7592	Disolves in any beverage Sugar Free
Polyethylene Glycol 3350 increases frequency of bowel movements and softens the stool.	

NDC 43386-312-07

Original Prescription Strength

GaviLax

Polyethylene Glycol 3350

Powder for Oral Solution, Osmotic Laxative

7 ONCE-DAILY DOSES

NET WT 4.1 OZ (119 g)

Compare to MiraLAX®

NDC 43386-312-07

Original Prescription Strength

GaviLAX

Polyethylene Glycol 3350 Powder for Oral Solution, Osmotic Laxative

- Relieves Occasional Constipation (irregularity)
- Softens Stool

LUPIN 7 ONCE-DAILY DOSES
NET WT 4.1 OZ (119 g)

UNVARNISHED AREA

*MiraLAX® is a registered trademark of Bayer Healthcare LLC

Manufactured for:
Lupin Pharmaceuticals, Inc.
Baltimore, MD 21202

SAP Code: 268476 07/2021

White Space Area .453"

White Space Area .453"

Drug Facts	Drug Facts (continued)
Active ingredient (in each dose) Polyethylene Glycol 3350, 17 g (cap filled to line)	Purpose Osmotic Laxative
Use <ul style="list-style-type: none"> relieves occasional constipation (irregularity) generally produces a bowel movement in 1 to 3 days 	
Warnings Allergy alert: Do not use if you are allergic to polyethylene glycol. Do not use if you have kidney disease, except under the advice and supervision of a doctor. Ask a doctor before use if you have <ul style="list-style-type: none"> nausea, vomiting, or abdominal pain a sudden change in bowel habits that lasts over 2 weeks irritable bowel syndrome 	
Polyethylene Glycol 3350 increases frequency of bowel movements and softens the stool.	
	Ask a doctor or pharmacist before use if you are taking a prescription drug When using this product you may have loose, watery, more frequent stools Stop use and ask a doctor if <ul style="list-style-type: none"> you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition. you get diarrhea you need to use a laxative for longer than 1 week
	Peel Here →

NDC 43386-312-14

Original Prescription Strength

GaviLax

Polyethylene Glycol 3350

Powder for Oral Solution, Osmotic Laxative

30 ONCE-DAILY DOSES

NET WT 17.9 OZ (510 g)

Compare to MiraLAX®

NDC 43386-312-14

Original Prescription Strength

GaviLAX

Polyethylene Glycol 3350 Powder for Oral Solution, Osmotic Laxative

- Relieves Occasional Constipation (irregularity)
- Softens Stool

LUPIN 30 ONCE-DAILY DOSES
NET WT 17.9 OZ (510 g)


.5625" White Space Area

Unvarnished area

*MiraLAX® is a registered trademark of Bayer Healthcare LLC

Manufactured by:
Lupin Pharmaceuticals, Inc.
Baltimore, MD 21202

SAP Code: 268478
Rev. 07/2021



N 3 43386-312-14 6

.5625" White Space Area

Drug Facts	Drug Facts (continued)
<p>Active ingredient (in each dose).....Purpose Polyethylene Glycol 3350, 17 g (cap filled to line).....Osmotic Laxative</p> <p>Use • relieves occasional constipation (irregularity) • generally produces a bowel movement in 1 to 3 days</p> <p>Warnings Allergy alert: Do not use if you are allergic to polyethylene glycol Do not use if you have kidney disease, except under the advice and supervision of a doctor Ask a doctor before use if you have • nausea, vomiting, or abdominal pain • a sudden change in bowel habits that lasts over 2 weeks • irritable bowel syndrome Ask a doctor or pharmacist before use if you are taking a prescription drug When using this product you may have loose, watery, more frequent stools Stop use and ask a doctor if • you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition. • you get diarrhea • you need to use a laxative for longer than 1 week</p>	<p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a POISON CONTROL CENTER right away. (1-800-222-1222)</p> <p>Directions • do not take more than directed unless advised by your doctor • the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line • adults and children 17 years of age and older: • use once daily • fill to top of the line in cap which is marked to indicate the correct dose (17 g) • stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink • do not combine with starch-based thickeners used for difficulty swallowing • ensure that the powder is fully dissolved before drinking • do not drink if there are any clumps • do not use more than 7 days • Children 16 years of age or under: ask a doctor</p> <p>Other information • store at 20°-25°C (68°-77°F) • tamper-evident: do not use if foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken</p> <p>Inactive ingredients none</p> <p>Questions or comments? 1-866-403-7592</p> <p>Polyethylene Glycol 3350 increases frequency of bowel movements and softens the stool</p> <p style="text-align: right;"><i>Dissolves in Any Beverage Sugar free</i></p>

GAVILAX

polyethylene glycol 3350 powder, for solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:43386-312
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P) (POLYETHYLENE GLYCOL 3350 - UNII:G2M7P15E5P)	POLYETHYLENE GLYCOL 3350	17 g in 17 g

Inactive Ingredients

Ingredient Name	Strength
SODIUM HYDROXIDE (UNII: 55X04QC321)	

Product Characteristics

Color	WHITE	Score	
Shape		Size	
Flavor		Imprint Code	

Contains**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:43386-312-14	510 g in 1 BOTTLE; Type 0: Not a Combination Product	10/09/2009	
2	NDC:43386-312-08	238 g in 1 BOTTLE; Type 0: Not a Combination Product	10/09/2009	
3	NDC:43386-312-07	119 g in 1 BOTTLE; Type 0: Not a Combination Product	01/31/2040	
4	NDC:43386-312-85	850 g in 1 BOTTLE; Type 0: Not a Combination Product	01/31/2040	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091077	10/09/2009	

Labeler - Lupin Pharmaceuticals, Inc. (089153071)**Registrant** - Novel Laboratories, Inc. (793518643)**Establishment**

Name	Address	ID/FEI	Business Operations
Novel Laboratories, Inc.		793518643	ANALYSIS(43386-312) , MANUFACTURE(43386-312) , PACK(43386-312)

Revised: 1/2024

Lupin Pharmaceuticals, Inc.