

**MUCINEX FAST-MAX SEVERE CONGESTION AND COUGH- guaifenesin, phenylephrine hydrochloride, and dextromethorphan hydrobromide tablet, film coated**  
**RB Health (US) LLC**

-----

**Mucinex® Fast-Max®**  
**Severe Congestion & Cough**

***Drug Facts***

<b><i>Active ingredients (in each caplet)</i></b>	<b><i>Purposes</i></b>
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

**Uses**

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
  - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
  - the intensity of coughing
  - the impulse to cough to help you get to sleep
  - nasal congestion due to a cold

**Warnings**

**Do not use**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

**When using this product do not use more than directed**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

**Other information**

- store between 20-25°C (68-77°F)

**Inactive ingredients**

corn starch, FD&C blue no. 2 aluminum lake, FD&C red no. 40 aluminum lake, magnesium stearate, maltodextrin, mica, microcrystalline cellulose, polyethylene glycol, polysorbate 80, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

**Questions?****1-866-MUCINEX (1-866-682-4639)**

You may also report side effects to this phone number.

Dist. by: Reckitt Benckiser, Parsippany, NJ 07054-0224

**PRINCIPAL DISPLAY PANEL - 20 Caplet Blister Pack Carton**

MAXIMUM STRENGTH

NDC 63824-515-01

Mucinex®

FAST-MAX®

SEVERE CONGESTION

& COUGH

Dextromethorphan HBr – Cough Suppressant

Guaifenesin – Expectorant

Phenylephrine HCl – Nasal Decongestant

- Controls Cough
- Relieves Nasal & Chest Congestion

- Thins & Loosens Mucus

20 CAPLETS  
FOR AGES 12+



<p><b>Drug Facts</b></p> <p><b>Active ingredients (in each caplet)</b></p> <p>Dextromethorphan HBr 10 mg.....Cough suppressant  Guaifenesin 200 mg.....Expectorant  Phenylephrine HCl 5 mg.....Nasal decongestant</p>	<p><b>Drug Facts (continued)</b></p> <p><b>When using this product do not use more than directed</b></p> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>nervousness, dizziness, or sleeplessness occur</li> <li>symptoms do not get better within 7 days or occur with fever</li> <li>cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.</li> </ul> <p><b>If pregnant or breast-feeding, ask a health professional before use.</b>  <b>Keep out of reach of children.</b>  In case of overdose, get medical help or contact a Poison Control Center right away.</p>
<p><b>Uses</b></p> <ul style="list-style-type: none"> <li>helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive</li> <li>temporarily relieves: <ul style="list-style-type: none"> <li>cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants</li> <li>the intensity of coughing</li> <li>the impulse to cough to help you get to sleep</li> <li>nasal congestion due to a cold</li> </ul> </li> </ul>	<p><b>Directions</b></p> <ul style="list-style-type: none"> <li>do not take more than 12 caplets in any 24-hour period</li> <li>adults and children 12 years of age and over: take 2 caplets every 4 hours</li> <li>children under 12 years of age: do not use</li> </ul>
<p><b>Warnings</b></p> <p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</li> </ul> <p><b>Ask a doctor before use if you have</b></p> <ul style="list-style-type: none"> <li>heart disease</li> <li>diabetes</li> <li>high blood pressure</li> <li>thyroid disease</li> <li>trouble urinating due to an enlarged prostate gland</li> <li>persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema</li> <li>cough that occurs with too much phlegm (mucus)</li> </ul>	<p><b>Other information</b></p> <ul style="list-style-type: none"> <li>store at 20-25°C (68-77°F)</li> </ul> <p><b>Inactive ingredients</b> colloidal silicon dioxide, FD&amp;C blue no. 2 aluminum lake, FD&amp;C red no. 40 aluminum lake, magnesium stearate, maltodextrin, mica, microcrystalline cellulose, polyethylene glycol, polysorbate 80, polyvinyl alcohol, povidone, pregelatinized starch, sodium starch glycolate, stearic acid, talc, titanium dioxide</p> <p><b>Questions? 1-866-MUCINEX (1-866-682-4639)</b></p>

<b>MUCINEX FAST-MAX SEVERE CONGESTION AND COUGH</b>			
guaifenesin, phenylephrine hydrochloride, and dextromethorphan hydrobromide tablet, film coated			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:63824-515
<b>Route of Administration</b>	ORAL		
<b>Active Ingredient/Active Moiety</b>			
<b>Ingredient Name</b>		<b>Basis of Strength</b>	<b>Strength</b>
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)		GUAIFENESIN	200 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)		PHENYLEPHRINE HYDROCHLORIDE	5 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)		DEXTROMETHORPHAN HYDROBROMIDE	10 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>FD&amp;C BLUE NO. 2</b> (UNII: L06K8R7DQK)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)	
<b>MICA</b> (UNII: V8A1AW0880)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	red	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	SRS
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63824-515-01	2 in 1 CARTON	08/01/2018	02/16/2025
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:63824-515-20	4 in 1 CARTON	01/01/2024	
2		5 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/01/2018	

**Labeler** - RB Health (US) LLC (081049410)

