

**FAMOTIDINE- famotidine tablet, film coated**  
**AMERISOURCE BERGEN**

-----  
**Famotidine**

***Drug Facts***

**Active ingredient (in each tablet)**

Famotidine, USP 20 mg

**Purpose**

Acid reducer

**Uses**

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

**Warnings**

**Allergy alert**

Do not use if you are allergic to famotidine or other acid reducers

**Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

**Ask a doctor before use if you have**

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

**Ask a doctor or pharmacist before use if you are** taking a prescription drug. Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if**

- your heartburn continues or worsens
- you need to take this product for more than 14 days

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### **Directions**

- adults and children 12 years and over:
  - to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
  - to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **10 to 60 minutes before** eating food or drinking beverages that cause heartburn
  - do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

### **Other information**

- read the directions and warnings before use
- store at 20° to 25° C (68° to 77° F)
- protect from moisture

### **Inactive ingredients**

colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, pregelatinized starch, talc, titanium dioxide

### **Questions?**

Call toll-free Monday-Friday 8:30 am to 5 pm EST at **1800-406-7984**.

Distributed By: AmerisourceBergen,  
1300 Morris Drive, Chesterbrook, PA 19087

### **PRINCIPAL DISPLAY PANEL - 20 mg Tablet Bottle Label**

NDC 46122-168-78

GOOD  
NEIGHBOR  
PHARMACY®

†Compare to the active ingredient of  
Maximum Strength Pepcid AC®

famotidine tablets, USP 20 mg

Acid Reducer

Maximum Strength

Just One Tablet

Prevents & Relieves Heartburn

Due to Acid Indigestion

100 tablets



NDC 46122-168-78

Compare to the active ingredient of Maximum Strength Pepcid AC®

famotidine tablets, USP 20 mg

# Acid Reducer

**Maximum Strength**Just One Tablet  
Prevents & Relieves Heartburn  
Due to Acid Indigestion**100 tablets**

## Drug Facts

**Active ingredient (in each tablet)**  
Famotidine, USP 20 mg**Purpose**  
Acid reducer**Uses** ■ relieves heartburn associated with acid indigestion and sour stomach  
■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

### Warnings

**Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers**TAMPER EVIDENT: DO NOT USE IF PRINTED FOIL UNDER CAP IS OPEN OR TORN.**

JUST ONE TABLET prevents and relieves heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.

R0719

ABC# 10101179

5197522

Expiration Date  
Batch No.open here  
Non Varnish Area

### Drug Facts (continued)

**Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

**Ask a doctor before use if you have**

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating, or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss

### Drug Facts (continued)

- nausea or vomiting
- stomach pain
- kidney disease

**Ask a doctor or pharmacist before use if you are taking a prescription drug.** Acid reducers may interact with certain prescription drugs.**Stop use and ask a doctor if**

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### Drug Facts (continued)

**Directions** ■ adults and children 12 years and over:

- to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

**Other information** ■ read the directions and warnings before use  
■ store at 20° to 25° C (68° to 77° F) ■ protect from moistureDistributed By: AmerisourceBergen,  
1300 Morris Drive, Chesterbrook, PA 19087  
Questions or Concerns?  
Visit [www.MyGNP.com](http://www.MyGNP.com)

### Drug Facts (continued)

**Inactive ingredients** colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, pregelatinized starch, talc, titanium dioxide**Questions?** Call toll-free Monday-Friday 8:30 am to 5 pm EST at 1800-406-7984.**Tips for Managing Heartburn**

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

\*All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Pepcid AC®.

## FAMOTIDINE

famotidine tablet, film coated

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:46 122-168
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	20 mg

## Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYDROXYPROPYL CELLULOSE (160000 WAMW) (UNII: RFW2ET671P)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
STARCH, CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

## Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	036
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46122-168-78	100 in 1 BOTTLE; Type 0: Not a Combination Product	05/01/2013	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090283	05/01/2013	

**Labeler** - AMERISOURCE BERGEN (007914906)

**Registrant** - Ranbaxy Pharmaceuticals Inc. (937890044)

## Establishment

Name	Address	ID/FEI	Business Operations
Ohm Laboratories Inc.		184769029	MANUFACTURE(46122-168)

Revised: 7/2019

AMERISOURCE BERGEN