

ASPIRIN- aspirin tablet
United Exchange Corp.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Aspirin 100 Tablets 60507, 2019

Active ingredient (in each tablet)	Purpose
Aspirin 325 mg (NSAID)*.....	Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug

Uses

temporarily relieves minor aches and pains due to:

- headache
- muscular aches
- minor pain of arthritis
- backache
- the common cold
- toothache
- premenstrual and menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- facial swelling
- shock
- hives
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if

- you have asthma
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease

- you are taking a diuretic

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away.
- ringing in the ears or loss of hearing occurs
- pain gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 to 2 tablets every 4 hours while symptoms last. Do not take more than 12 tablets in 24 hours unless directed by a doctor.
- children under 12 years: do not use unless directed by a doctor

Other information

- Use by expiration date on package
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredient corn starch

Distributed by:

United Exchange Corp.

17211 Valley View Ave.

Cerritos, CA 90703

Made in USA

†Compare to the Active Ingredient in Genuine Bayer®

ASPIRIN (NSAID)

Pain Reliever • Fever Reducer

For the temporary relief from the pain of:

- headache • muscle pain • toothache
- menstrual pain • pain and fever of colds
- minor pain of arthritis



100 TABLETS

325 mg each



7 8070760501 3

This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark, Genuine Bayer® Aspirin. 50844 ORG011324912

Distributed By:

UNITED EXCHANGE CORP
172111 Valley View Ave. Cerritos, CA 90703

Drug Facts

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Active ingredient (in each tablet)
Aspirin 325 mg (NSAID)* Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses temporarily relieves minor aches and pains due to:
 ■ headache ■ muscular aches ■ minor pain of arthritis
 ■ backache ■ the common cold ■ toothache
 ■ premenstrual and menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:
 ■ facial swelling
 ■ shock ■ hives ■ asthma (wheezing)
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
 ■ are age 60 or older
 ■ take a blood thinning (anticoagulant) or steroid drug
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

PEEL HERE FOR MORE DRUG FACTS

No print/No varnish
Lot & Exp date

Adhesive Only

STOP PEELING

Questions or comments? 1-800-426-9391

Inactive ingredient corn starch

Other information ■ use by expiration date on package
 ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Directions ■ drink a full glass of water with each dose
 ■ adults and children 12 years and over: take 1 to 2 tablets every 4 hours while symptoms last. Do not take more than 12 tablets in 24 hours unless directed by a doctor.
 ■ children under 12 years: do not use unless directed by a doctor

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222), right away.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding:
 ■ vomit blood ■ have bloody or black stools
 ■ have stomach pain that does not get better
 ■ an allergic reaction occurs. Seek medical help right away.
 ■ ringing in the ears or loss of hearing occurs
 ■ pain gets worse or lasts more than 10 days
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present ■ new symptoms occur

Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer.

Ask a doctor before use if ■ you have asthma
 ■ stomach bleeding warning applies to you
 ■ you have a history of stomach problems, such as heartburn
 ■ you have high blood pressure, heart disease, liver cirrhosis or kidney disease ■ you are taking a diuretic

Drug Facts (continued)

■ have had stomach ulcers or bleeding problems
 ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

■ you are allergic to aspirin or any other pain reliever/fever reducer.

■ you have asthma

■ stomach bleeding warning applies to you

■ you have a history of stomach problems, such as heartburn

■ you have high blood pressure, heart disease, liver cirrhosis or kidney disease

■ you are taking a diuretic

ASPIRIN

aspirin tablet

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:65923-507

Route of Administration

ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	ASPIRIN;A1
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65923-507-00	100 in 1 BOTTLE; Type 0: Not a Combination Product	03/02/2016	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part343	03/02/2016	

Labeler - United Exchange Corp. (840130579)

Revised: 12/2019

United Exchange Corp.