# GUMNUMB TOPICAL ANESTHETIC- benzocaine gel Crosstex International Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

### Active Ingredient (in each gram)

Benzocaine 200mg

#### **Purpose**

Oral Anesthetic

#### Use

For oral mucosa use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.

#### Warnings

**Methemoglobinemia warning:** Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

**Allergy Alert:** Do not use on patients with a history of allergies to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

#### Do not use

- for more than 7 days unless directed by a physician. If sore mouth symptoms do not improve in 7 days; irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your physician promptly.
- for teething
- in children under 2 years of age

**When using this product** Avoid contact with eyes. If contact occurs, flush with water.

**Do not exceed recommended dosage.** If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

If pregnant or breast feeding, ask a physician before use.

Keep out of reach of children.

#### **Directions**

- Apply only amount needed to the oral mucosa to prevent or relieve pain.
- children under 2 years of age: do not use

#### Other Information

Store at  $59^{\circ}$  -  $86^{\circ}$ F ( $15^{\circ}$ - $30^{\circ}$ C).

Protect from freezing.

## **Inactive Ingredients**

flavoring, PEG 3350, PEG 400, sodium saccharin. May contain blue #1, green #3, green #5, red #3, red #28, red #40, yellow #5, (tartrazine), yellow #6, as a color additive.

## **Questions or comments?**

crosstex@crosstex.com or www.crosstex.com



REF UTM





UTM-BTL Rev. 7 - 7/18

Manufactured for: Crosstex International, Inc. 10 Ranick Road Hauppauge, NY 11788-4209



from CROSSIEX

tartrazine), yellow #6, as a color

additive.

olue #1, green #3, green #5, red

sodium saccharin. May contair

nactive ingredients lavoring, PEG 3350, PEG 400 #3, red #28, red #40, yellow #5,

in 7 days; irritation, pain, or redness mouth symptoms do not improve persists or worsens; or if swelling, rash or fever develops, see your

in children under 2 years of age When using this product physician promptly. · for teething

Do not exceed recommended Avoid contact with eyes. If contact occurs, flush with water

Poison Control Center right away dosage. If more than used for oain is accidentally swallowed get medical help or contact a

warning: Use of this product may

Methemoglobinemia

cause methemoglobinemia, a serious condition that must be if pregnant or breast feeding, (eep out of reach of children. ask a physician before use.

treated promptly because it reduces

the amount of oxygen carried in

 Apply only amount needed to children under 2 years of age: the oral mucosa to prevent or **Directions** relieve pain. the blood. This can occur even if you have used this product before. Stop

use and seek immediate medical

attention if you or a child in your pale, gray or blue colored skin care develops: cvanosis)

headache

 dizziness or lightheadedness fatigue or lack of energy shortness of breath

rapid heart rate

Store at 59° - 86°F (15 - 30°C).

Protect from freezing.

Other information

do not use

procaine, butacaine, benzocaine or patients with a history of allergies Allergy Alert: Do not use on to local anesthetics such as

other "caine" anesthetics.

directed by a physician. If sore for more than 7 days unless Do not use

Questions or comments?

crosstex@crosstex.com or

www.crosstex.com

99190003 Rev. 7 - 7/18

**Drug Facts** 

Oral Ánesthetic Purpose

**Active ingredients** *(in each gram)* Benzocaine 200 mg

## **GUMNUMB TOPICAL ANESTHETIC**

For oral mucosal use only, as

directed by dentist. For the

temporary relief of pain due to

minor dental procedures.

benzocaine gel

**Product Information** 

**Inactive Ingredients** 

NDC:24794-221 **Product Type HUMAN OTC DRUG** Item Code (Source)

DENTAL **Route of Administration** 

Active Ingredient/Active Moiety

**Ingredient Name Basis of Strength** Strength BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5) **BENZ OCAINE** 200 mg in 1 g

**Ingredient Name** Strength POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P) POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)

SACCHARIN SODIUM (UNII: SB8ZUX40TY) FD&C YELLOW NO. 5 (UNII: I753WB2F1M)

FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
FD&C RED NO. 3 (UNII: PN2ZH5LOQY)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
<b>D&amp;C RED NO. 28</b> (UNII: 767IP0Y5NH)	
D&C GREEN NO. 5 (UNII: 8J6RDU8L9X)	

Product Characteristics				
Color	green	Score		
Shape		Size		
Flavor	MINT	Imprint Code		
Contains				

Packaging				
# Item Code	Package Description	Marketing Start Date	Marketing End Date	
1 NDC:24794-221-	30 g in 1 JAR; Type 0: Not a Combination Product	07/01/2018		

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph not final	part356	07/01/2013		

## **Labeler -** Crosstex International Inc. (057728685)

Revised: 1/2022 Crosstex International Inc.