STONA- dextromethorphan hydrobromide lozenge Sato Pharmaceutical Co., Ltd.

Stona Troche

Active ingredients (in each lozenge) Dextromethorphan HBr 10mg

Purpose

Cough suppressant

Uses temporarily relieves cough due to minor throat and bronchial irritation associated with

■a cold ■inhaled irritants

Enter section text here

Do not use

■this product for persistent or chronic cough such as occurs with smoking, asthema, or emphysem, or if cough is accompanied by excessive phlegm (mucus) unless directed by a doctor.

Ask a doctor or pharmacist before use if

■you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. A persistent cough may be a sign of a serious condition. If cough persists for more than 1 week, tends to recur, or is accompanied by fever, rash, or persistent headache, consult a doctor.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away.

Directions

Adults and children 12 years and older - Let 1 or 2 lozenges dissolve slowly in the mouth every 4 hours, not more than 12 lozenges in 24 hours.

Children 6 years to under 12 years - Let 1 lozenge dissolve slowly in the mouth every 4 hours, not more than 6 lozenges in 24 hours.

Children under 6 years - Ask a doctor

- ■once or twice daily, preferably morning and evening, when needed or as directed by a doctor
- ■start initial dosage with minimum dose, then adjust it to suit to bowel condition

Other information

■Phenylketonurics: Contains Phenylalanine (18.3mg) per lozenge.

Inactive ingredients aspartame, cetylpyridinium chloride, flavor, glycyrrhiza, hydroxypropyl cellulose, silicon dioxide, magnesium stearate, sucrose.

stonatrochecart.jpg

COUGH SUPPRESSANT STONA TROCHE

NDC 49873-301-01

EXP.DATE 使用期限 딞 ない。

0 86

sato

★ FOR YOUR SAFETY

ONA TROCHE

COUGH SUPPRESSANT

かぜなどによる咳に ストナトローチ

Manufactured by

SATO PHARMACEUTICAL CO., LTD.

1-5-27 MOTOAKASAKA MINATO-KU TOKYO, JAPAN

Reports of serious side effects associated with use of this product can be sent to: SATO PHARMACEUTICALINC 20501 Earl Street, Suite 3, Torrance, CA 90503



12 Lozenges(12錠)

かぜなどによる咳に ストナトローチ

12錠

よってこういつんもかな。ムーテルンスへ 会覧の参の学 表質は、スーロハセルンプログッキのシュ、よキエクシッセ、評書 をかってってっている。 のかのかのでは、ステアールでは、またのかのでは、 はなっていて、またのかのでは、 はないできた。

"ままい2.13を計算器 8.81かくニモベルニュてコ中級1:非常遊覧とすかか 事業の命の分

景小の萬未下3	パイセス/アノ制御に前題
%√r@.£11~9	き、J-セスリンと中で作り、第1回! なく、J-セスリンエリを扱いよこ類の J-角種で大陸を加るコロに対面を フィスンファンタ
表小の主幕を引む個人大	かるか)でかず内の、285~1回1 となりてしま聞いるに関係をして である第2にの内は関係的ない。 ではいるいのでは、2000年間

- 意託の土田御

は個のおくれるもあの

支管液や3のより生すでも:1重体の修設率や雰囲 譲 歳 (中盤1)量代・役類成序 8m01ペヤベルイスロイスキャ類素末か具 示表品藥因

Anactive ingvedients asparteme, cetylpyridinium chlorde, flavor, glycymhza, flydroxypropyl ceilulose, silicon dioside, magnesium stearabe, sucrose.

Other Information Contains Phenylalanine (18.3mg)

Prenylalanine (18.3mg)

6 years to Let 1 lozenge dissolve slowly in the mouth every 4 house, not a more than 6 lozenges in 24 hours, not Directions

Steels and chicken Let 1 or 2 losenges in 24

Adults and chicken Let 0 or 3 losenges in 24

Adults and chicken Let 0 or 3 losenges in 24

Directions

Directions

Vester and the control of children. In case of overdose, get medical help or contact a Polson Control Center night away.

Uses temporarily relieves cough due to minor threat and bronchist imitation associated with a soold a inhalted imitation Dextromethorphan HBr 10mg... Active ingredient(in each lozenge) Purpose Drug Facts

Use only if carton is sealed.
Lozenges individually blister sealed.
If seal is broken, do not use.
技会のため外籍の對が聞いている場合は服用しないでください。
微剤を包装しているシールが吸れている場合は服用しないでください。

STONA

dextromethorphan hydrobromide lozenge

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49873-301
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg

Inactive Ingredients		
Ingredient Name	Strength	
CETYLPYRIDINIUM CHLORIDE (UNII: D9OM4SK49P)		
LICORICE (UNII: 61ZBX54883)		
HYDROXYPROPYL CELLULOSE (UNII: RFW2ET671P)		
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)		
MAGNESIUM STEARATE (UNII: 70097M6I30)		
SUCROSE (UNII: C151H8M554)		
ASPARTAME (UNII: Z0H242BBR1)		

Product Characteristics			
Color	brown (pale brown)	Score	no score
Shape	ROUND	Size	14mm
Flavor	BLUEBERRY, GRAPE	Imprint Code	SATO;SATO
Contains			

F	Packaging				
#	tem Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:49873- 301-01	1 in 1 CARTON	11/19/1995		
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	11/19/1995	

Labeler - Sato Pharmaceutical Co., Ltd. (690575642)

Establishment			
Name	Address	ID/FEI	Business Operations
Sato Pharmaceutical Co., Ltd.		715699133	manufacture(49873-301) , label(49873-301) , pack(49873-301)

Revised: 11/2023 Sato Pharmaceutical Co., Ltd.