

**TARGET MAXIMUM STRENGTH COLD FLU AND SORE THROAT- acetaminophen,
dextromethorphan hbr, guaifenesin, phenylephrine hcl solution
TARGET CORPORATION**

Target Maximum Strength Cold, Flu & Sore Throat

Active ingredients (in each 20 mL)

Acetaminophen 650 mg

Dextromethorphan HBr 20 mg

Guaifenesin 400 mg

Phenylephrine HCl 10 mg

Purposes

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - nasal congestion
 - minor aches and pain
 - sore throat
 - headache
 - stuffy nose
 - sinus congestion and pressure
- temporarily reduces fever
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

- 3 or more alcoholic drinks everyday while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

When using this product,

do not use more than directed.

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present

- new symptoms occur
- cough comes back, or occurs with fever, rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided.
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- mL = milliliter
- **adults and children 12 years of age and older:** 20 mL in dosing cup provided every 4 hours
- **children under 12 years of age:** Do not use

Other information

- **each 20 mL contains:** sodium 8 mg
- store at room temperature
- do not refrigerate.

Inactive ingredients

anhydrous citric acid, edetate disodium, FD&C Blue No. 1, FD&C Red No. 40, flavor, potassium citrate, propylene glycol, propyl gallate, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum

Questions or comments?

1-866-467-2748

Principal Display Panel

Compare to active ingredients in Maximum Strength[‡] Mucinex[®] Fast-Max[®] Cold, Flu & Sore Throat*

NDC# 11673-773-06

Maximum Strength[‡]

Cold, Flu & Sore Throat

Acetaminophen - Pain Reliever/Fever Reducer

Dextromethorphan HBr - Cough Suppressant

Guaifenesin - Expectorant

Phenylephrine HCl - Nasal Decongestant

- Controls Cough, This & Loosens Mucus
- Nasal & Chest Congestion
- Sinus Pressure & Congestion
- Body Pain, Headache, Fever & Sore Throat

For Ages 12+

6 FL OZ (180 mL)

Tamper evident: Do not use if printed seal under cap is broken or missing.

[‡]Per 4-hour dose.

Up & up

AGES 12+ YEARS

Distributed by:

Target Corp.

Mpls., MN 5403

***This product is not manufactured or distributed by Reckitt Benckiser, the distributor of Maximum Strength Mucinex® Fast-Max™ Cold, Flu & Sore Throat.**

PEEL CORNER TO READ COMPLETE DRUG FACTS AND INFORMATION

MULTI-SYMPOM



Top Layer Art

DIE LINE - DOES NOT PRINT

3.5"

Drug Facts

Active ingredients (in each 20 mL) Purposes

Acetaminophen 650 mg Pain reliever/fever reducer
 Dextromethorphan HBr 20 mg Cough suppressant
 Guaifenesin 400 mg Expectorant
 Phenylephrine HCl 10 mg Nasal decongestant

Uses

temporarily relieves these common cold and flu symptoms: cough
 nasal congestion minor aches and pains sore throat
 headache stuffy nose sinus congestion and pressure
 temporarily reduces fever temporarily promotes nasal and/or
 sinus drainage helps loosen phlegm (mucus) and thin bronchial
 secretions to rid the bronchial passageways of bothersome mucus
 and make coughs more productive

**Tamper evident: Do not use if printed inner seal under cap
 is broken or missing.**

094 14 8715 R00 C-001227-01-055
 Dist. by Target Corp., Mpls., MN 55403
 TM & ©2020 Target Brands, Inc.

Lot: 73706T1B
 Exp:

*This product is not manufactured or
 distributed by Reckitt Benckiser; the
 distributor of Maximum Strength Mucinex®
 Fast-Max® Cold & Sore Throat.

PEEL CORNER TO READ COMPLETE
 DRUG FACTS AND INFORMATION

Top layer Back Art

hinge

Drug Facts (continued)

Warnings

Liver warnings: This product contains acetaminophen. Severe liver
 damage may occur if you take: more than 6 doses in 24 hours,
 which is the maximum daily amount with other drugs containing
 acetaminophen 3 or more alcoholic drinks daily while using this
 product.

Allergy alert: Acetaminophen may cause severe skin reactions.
 Symptoms may include: skin reddening blisters rash. If
 a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more
 than 2 days, is accompanied or followed by fever, headache, rash,
 rashes, or vomiting, consult a doctor promptly.

Do not use with any other drug containing acetaminophen
 (prescription or nonprescription). If you are not sure whether
 a drug contains acetaminophen, ask a doctor or pharmacist.
 If you are now taking a prescription monoamine oxidase
 inhibitor (MAOI) (certain drugs for depression, psychiatric, or
 emotional conditions, or Parkinson's disease), or for 2 weeks after
 stopping the MAOI drug, if you do not know if your prescription
 drug contains an MAOI, ask a doctor or pharmacist before taking
 this product.

Ask a doctor before use if you have liver disease heart disease
 diabetes high blood pressure thyroid disease trouble
 urinating due to an enlarged prostate gland persistent or chronic
 cough such as occurs with smoking, asthma, chronic bronchitis,
 or emphysema cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the
 blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if nervousness, dizziness, or

Middle Layer Face Art

hinge

Drug Facts (continued)

sleeplessness occur pain, nasal congestion, or cough gets
 worse, or lasts more than 7 days fever gets worse, or lasts
 more than 3 days redness or swelling is present new
 symptoms occur cough comes back, or occurs with fever, rash,
 or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children.**

Overdose warning: Taking more than the recommended dose
 (overdose) may cause liver damage. In case of overdose, get
 medical help or contact a Poison Control Center right away at
 1-800-222-1222. Quick medical attention is critical for adults as
 well as for children even if you don't notice any signs or
 symptoms.

Directions

do not take more than directed (see Overdose warning)
 do not take more than 6 doses in any 24-hour period
 measure only with dosing cup provided do not use dosing cup
 with other products dose as follows or as directed by a doctor
 mL = milliliter adults and children 12 years of age and
 older: 20 mL in dosing cup provided every 4 hours. children
 under 12 years of age: Do not use.

Other information

each 20 mL contains: sodium 8 mg store at room
 temperature do not refrigerate

Inactive ingredients anhydrous citric acid, edetate
 disodium, FD&C Blue No. 1, FD&C Red No. 40, flavors, potassium
 citrate, propylene glycol, propyl gallate, purified water, sodium
 benzoate, sorbitol, sucralose, xanthan gum

Questions? call 1-866-467-2748

TARGET MAXIMUM STRENGTH COLD FLU AND SORE THROAT

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11673-773
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 20 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
POTASSIUM CITRATE (UNII: EE90ONI6FF)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYL GALLATE (UNII: 8D4SNN7V92)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	

SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	BLUE	Score	
Shape		Size	
Flavor	FRUIT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-773-06	180 mL in 1 BOTTLE, PLASTIC; Type 9: Other Type of Part 3 Combination Product (e.g., Drug/Device/Biological Product)	04/02/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/02/2020	

Labeler - TARGET CORPORATION (006961700)