

**DAY AND NIGHT COLD AND FLU FORMULA- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride**  
**Wal-Mart Stores Inc**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**Day and Night Cold and Flu Formula**

**Non-Drowsy Day Cold and Flu**  
**Active ingredients (in each softgel)**

Acetaminophen 325 mg  
Dextromethorphan hydrobromide 10 mg  
Phenylephrine hydrochloride 5 mg

**Night Cold and Flu**  
**Active ingredients (in each softgel)**

Acetaminophen 325 mg  
Dextromethorphan hydrobromide 10 mg  
Doxylamine succinate 6.25 mg  
Phenylephrine hydrochloride 5 mg

**Purposes**

**DAY COLD & FLU FORMULA**

Pain reliever/fever reducer  
Cough suppressant  
Nasal decongestant

**Purposes**

**NIGHT COLD & FLU FORMULA**

Pain reliever/fever reducer  
Cough suppressant  
Antihistamine  
Nasal decongestant

**Uses**

**DAY COLD & FLU FORMULA**

- temporarily relieves these symptoms due to a cold or flu:
  - minor aches and pains
  - headache
  - cough
  - sore throat
  - nasal and sinus congestion

- temporarily reduces fever

## Uses

### **NIGHT COLD & FLU FORMULA**

- temporarily relieves these symptoms due to a cold or flu:
  - minor aches and pains
  - headache
  - nasal and sinus congestion
  - cough
  - sore throat
  - runny nose
  - sneezing
- temporarily reduces fever

### **DAY COLD & FLU FORMULA**

#### **Warnings**

##### **Liver warning**

These products contain acetaminophen. Severe liver damage may occur if you take

- more than 10 softgels in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

##### **Sore throat warning**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

##### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

##### **Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- cough with excessive phlegm (mucus)

- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin.

**When using this product do not exceed recommended dosage**

**Stop use and ask a doctor if**

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

- nervousness, dizziness, or sleeplessness occurs

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## **NIGHTTIME COLD & FLU FORMULA**

### **Warnings**

#### **Liver warning**

These products contain acetaminophen. Severe liver damage may occur if you take

- more than 10 softgels in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

#### **Sore throat warning**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use to sedate children.**

#### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

- in children under 12 years of age

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough with excessive phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

### **Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

### **When using this product**

- **do not exceed recommended dosage**
- may cause marked drowsiness
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

### **Stop use and ask a doctor if**

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

- nervousness, dizziness, or sleeplessness occurs

**If pregnant or breast-feeding**, ask a health professional before use.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## **DAY COLD & FLU FORMULA**

### **Directions**

- do not take more than the recommended dose
- adults and children 12 years and over: take 2 softgels with water every 4 hours. Do not exceed 10 softgels in 24 hours or as directed by a doctor.
- children under 12 years: do not use

## **NIGHTTIME COLD & FLU FORMULA**

### **Directions**

- do not take more than the recommended dose
- adults and children 12 years and over: take 2 softgels with water every 4 hours. Do not exceed 10 softgels in 24 hours or as directed by a doctor.
- children under 12 years: do not use

#### **DAY COLD & FLU FORMULA**

##### **Other information**

- store at room temperature 15°-30°C (59°-86°F).

#### **NIGHTTIME COLD & FLU FORMULA**

##### **Other information**

- store at room temperature 15°-30°C (59°-86°F) and avoid excessive heat.

#### **DAY COLD & FLU FORMULA**

##### **Inactive ingredients**

FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, and white edible ink

#### **NIGHTTIME COLD & FLU FORMULA**

##### **Inactive ingredients**

D&C yellow #10, FD&C blue #1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, and white edible ink

#### **DAY COLD & FLU FORMULA**

##### **Questions or comments?**

Call toll free: **1-888-287-1915**

#### **NIGHTTIME COLD & FLU FORMULA**

##### **Questions or comments?**

Call toll free: **1-888-287-1915**

#### **Principal Display Panel - Carton Label**

Equate Day and Night Cold and Flu Formula 20ct

NDC 49035-315-08

\*Compare to the active ingredients in Alka-Seltzer PLUS<sup>®</sup> Day and Night Cold and Flu Formula



**DAY COLD & FLU FORMULA**

**Drug Facts (continued)**

See these warnings if you develop severe, painful sores that last 7 days, are accompanied by fever, headache, rash, nausea, or vomiting, or if you develop:

- severe allergic reactions that cause trouble with breathing, difficulty swallowing, or hives; or
- severe dizziness, fainting, or loss of consciousness.

Do not use:

- with any other drug containing acetaminophen, aspirin, or ibuprofen. If you are not sure, ask your doctor or pharmacist.
- if you are now taking or planning to take an MAO inhibitor (MAOI) such as selegiline, rasagiline, or saquinone, or a monoamine oxidase inhibitor (MAOI) such as phenelzine, tranylcypromine, or a related medicine, or if you have used one of these drugs in the last 14 days.
- if you have used another MAOI, such as a doctor or pharmacist, within the last 14 days.
- if you have used another MAOI, such as a doctor or pharmacist, within the last 14 days.
- if you have used another MAOI, such as a doctor or pharmacist, within the last 14 days.

Ask a doctor or pharmacist before use if you are taking the following medicines:

- alcohol, or any other drug containing acetaminophen, aspirin, or ibuprofen.
- any other drug containing acetaminophen, aspirin, or ibuprofen.
- any other drug containing acetaminophen, aspirin, or ibuprofen.
- any other drug containing acetaminophen, aspirin, or ibuprofen.

Some warning information may be omitted for brevity.

**NIGHT COLD & FLU FORMULA**

**Drug Facts (continued)**

Do not use:

- with any other drug containing acetaminophen, aspirin, or ibuprofen. If you are not sure, ask your doctor or pharmacist.
- if you are now taking or planning to take an MAO inhibitor (MAOI) such as selegiline, rasagiline, or saquinone, or a monoamine oxidase inhibitor (MAOI) such as phenelzine, tranylcypromine, or a related medicine, or if you have used one of these drugs in the last 14 days.
- if you have used another MAOI, such as a doctor or pharmacist, within the last 14 days.
- if you have used another MAOI, such as a doctor or pharmacist, within the last 14 days.
- if you have used another MAOI, such as a doctor or pharmacist, within the last 14 days.

Ask a doctor or pharmacist before use if you are taking the following medicines:

- alcohol, or any other drug containing acetaminophen, aspirin, or ibuprofen.
- any other drug containing acetaminophen, aspirin, or ibuprofen.
- any other drug containing acetaminophen, aspirin, or ibuprofen.
- any other drug containing acetaminophen, aspirin, or ibuprofen.

Some warning information may be omitted for brevity.

<b>DAY AND NIGHT COLD AND FLU FORMULA</b>			
acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride kit			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:49035-315

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49035-315-08	1 in 1 CARTON; Type 0: Not a Combination Product	08/23/2017	

## Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BLISTER PACK	12
Part 2	1 BLISTER PACK	8

## Part 1 of 2

### DAYTIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid filled

## Product Information

Route of Administration ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

## Inactive Ingredients

Ingredient Name	Strength
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	

## Product Characteristics

Color orange (clear) Score no score

<b>Shape</b>	capsule (oblong)	<b>Size</b>	20 mm
<b>Flavor</b>		<b>Imprint Code</b>	PC9
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		1 in 1 CARTON		
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	08/23/2017	

### Part 2 of 2

#### NIGHT COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride capsule, liquid filled

### Product Information

<b>Route of Administration</b>	ORAL
--------------------------------	------

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

### Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	

SORBITAN (UNII: 6O92ICV9RU)

### Product Characteristics

<b>Color</b>	green (clear)	<b>Score</b>	no score
<b>Shape</b>	capsule (oblong)	<b>Size</b>	21mm
<b>Flavor</b>		<b>Imprint Code</b>	PC22
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		1 in 1 CARTON		
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	08/23/2017	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	08/23/2017	

**Labeler** - Wal-Mart Stores Inc (051957769)

### Establishment

Name	Address	ID/FEI	Business Operations
Humanwell PuraCap Pharmaceuticals (Wuhan) Co., Ltd		421293287	manufacture(49035-315) , analysis(49035-315)

Revised: 11/2019

Wal-Mart Stores Inc