

MAGNESIUM CITRATE - magnesium citrate liquid

Cardinal Health

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredient

Magnesium citrate 1.745 g per fl oz

Purpose

Saline Laxative

Use for relief of occasional constipation (irregularity). Generally produces bowel movement in 1/2 to 6 hours

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium or potassium-restricted diet
- noticed a sudden change in bowel habits that persists over a period of 2 weeks
- already used a laxative for a period longer than 1 week

Ask a doctor or a pharmacist before use if you are taking any other drug. Take this product 2 or more

more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions •drink a full glass (8 ounces) of liquid with each dose •may be taken as a single daily dose or in divided doses

adults and children 12 years of age and over - 6.5 to 10 fl oz maximum 10 fl oz in 24 hours

children 6 to under 12 years of age - 3 to 7 fl oz maximum 7 fl oz in 24 hours

children 2 to under 6 years of age - 2 to 3 fl oz maximum 3 fl oz in 24 hours

children under 2 years of age- ask a doctor

discard unused product within 24 hours of opening bottle

Other information • magnesium content 290 mg per 1 fl oz (30 mL)

•potassium content 80 mg per 1 fl oz (30mL)

•store at temperatures between 46° and 86° F (8° and 30° C)

Inactive ingredients citric acid, lemon oil, potassium bicarbonate, sodium saccharin, water purified

DO NOT USE IF TAMPER EVIDENT TWIST OFF-OFF CAP IS MISSING, BROKEN OR SEPARATED FROM NECKRING

NDC37205-110-38

LEADER

Magnesium

Citrate

Oral Solution

Saline Laxative
 Lemon flavor
 Pasteurized
 The Sparkling Laxative
 10 FL OZ (296 mL)

DO NOT USE IF TAMPER EVIDENT TWIST-OFF CAP IS MISSING, BROKEN OR SEPARATED FROM NECKING
 NDC 37205-110-38

LEADER

Magnesium Citrate Oral Solution
 Saline Laxative
 Lemon Flavor
 Pasteurized
 The Sparkling Laxative

10 FL OZ (296 mL)

Drug Facts
Active ingredient Magnesium citrate 1.745 g per fl oz
Purpose Saline laxative

Use for relief of a casual constipation (irregularity). Generally produces bowel movement in 12 to 6 hours.

Warnings
 Do not use if you are on a low salt diet.
 Ask a doctor before use if you have: kidney disease; a magnesium- or sodium-restricted diet; abdominal pain, nausea, or vomiting; noticed a sudden change in bowel habits that persists over a period of 2 weeks; already used a laxative for a period longer than 1 week.
 Ask a doctor or pharmacist before use if you are taking any other drug. Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.
 Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.
 If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions Drink a full glass (8 ounces) of liquid with each dose. May be taken as a single daily dose or in divided doses.
 Adults and children 12 years of age and over: 6.5 to 10 fl oz maximum 10 fl oz in 24 hours
 Children 6 to under 12 years of age: 3 to 7 fl oz maximum 7 fl oz in 24 hours
 Children 2 to under 6 years of age: 2 to 3 fl oz maximum 3 fl oz in 24 hours
 Children under 2 years of age: ask a doctor
 Discard unused product within 24 hours of opening bottle.

Other information Magnesium content 290 mg per 1 fl oz (30 mL). Sodium content 45 mg per 1 fl oz (30 mL). Store at temperatures between 49° and 69° F (5° and 30° C).

Inactive ingredients citric acid, lemon oil, polyethylene glycol, sodium bicarbonate, sodium saccharin, sucrose, water purified.

DISTRIBUTED BY CARDINAL HEALTH, DUBLIN, OHIO 43017
 CRI 1332-031 www.mylaxative.com 1-800-294-6318
 L00019635C
 56296 3827

MAGNESIUM CITRATE

magnesium citrate liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37205-110
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Magnesium citrate (UNII: RHO26O1T9V) (magnesium cation - UNII:T6V3LHY838)	Magnesium citrate	1.745 g in 29.6 mL

Inactive Ingredients

Ingredient Name	Strength
LEMON OIL (UNII: I9GRO824LL)	
POTASSIUM BICARBONATE (UNII: HM5Z15LEBN)	
WATER (UNII: 059QF0K00R)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	LEMON (LEMON FLAVOR)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37205-110-38	.296 mL in 1 BOTTLE, GLASS		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	06/21/1989	

Labeler - Cardinal Health (097537435)

Registrant - Vi-Jon, Inc (790752542)

Establishment

Name	Address	ID/FEI	Business Operations
Vi-Jon, Inc		790752542	manufacture

Revised: 11/2011

Cardinal Health