

**ZYRTEC-D ALLERGY PLUS CONGESTION- cetirizine hydrochloride and pseudoephedrine hydrochloride tablet, film coated, extended release  
Johnson & Johnson Consumer Inc.**

-----  
**ZYRTEC-D**

**Allergy Plus Congestion**

***Drug Facts***

<b><i>Active ingredients (in each extended release tablet)</i></b>	<b><i>Purpose</i></b>
Cetirizine HCl 5 mg	Antihistamine
Pseudoephedrine HCl 120 mg	Nasal decongestant

**Uses**

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
  - runny nose
  - sneezing
  - itchy, watery eyes
  - itching of the nose or throat
  - nasal congestion
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily restores freer breathing through the nose

**Warnings**

**Do not use**

- if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- heart disease
- thyroid disease
- diabetes
- glaucoma
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- liver or kidney disease. Your doctor should determine if you need a different dose.

**Ask a doctor or pharmacist before use if you are** taking tranquilizers or sedatives.

### **When using this product**

- **do not use more than directed**
- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

### **Stop use and ask a doctor if**

- an allergic reaction to this product occurs. Seek medical help right away.
- you get nervous, dizzy, or sleepless
- symptoms do not improve within 7 days or are accompanied by fever

### **If pregnant or breast-feeding:**

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

### **Directions**

- do not break or chew tablet; swallow tablet whole

adults and children 12 years and over	take 1 tablet every 12 hours; do not take more than 2 tablets in 24 hours.
adults 65 years and over	ask a doctor
children under 12 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

### **Other information**

- store between 20° to 25°C (68° to 77°F)
- **do not use if carton is opened or if blister unit is broken**

### **Inactive ingredients**

colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide

### **Questions?**

call **1-800-343-7805** (toll-free) or **215-273-8755** (collect)

**PRINCIPAL DISPLAY PANEL**

***Original Prescription Strength***

NDC 50580-728-24

**ZYRTEC-D**®

*Cetirizine HCl* **5 mg**/antihistamine

*Pseudoephedrine HCl* **120 mg**/nasal decongestant

*Extended Release Tablets*

**ALLERGY + CONGESTION**

**INDOOR + OUTDOOR**

**ALLERGIES**

**12**

**HOUR**

**RELIEF OF**

- *Sneezing*
- *Itchy, Watery Eyes*
- *Runny Nose*
- *Itchy Throat or Nose*
- *Sinus Pressure*
- *Nasal Congestion*

(Actual Size)

**24**

**EXTENDED**

**RELEASE TABLETS**

3 0045-0204-24 0



Made in Belgium  
D. Tribu by:  
JOHNSON & JOHNSON CONSUMER INC.  
Fort Washington, PA 19084 USA  
© J&J 2017  
www.zyrtec.com

**Questions?** call 1-800-343-7805 (toll-free) or 215-273-8755 (collect)

**Inactive ingredients** (continued)  
magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide

The trade dress of this ZYRTEC-D package is subject to trademark protection.  
U.S. Patent Nos. 6468009, 7014867, 7226614  
30038756

**Drug Facts (continued)**

**Other information**

- do not use if carton is opened or blister unit is broken
- store between 20° to 25°C (68° to 77°F)
- ask a doctor if you have:
  - heart disease
  - thyroid disease
  - diabetes
  - glaucoma
  - high blood pressure
- ask a doctor before use if you have:
  - heart disease
  - thyroid disease
  - diabetes
  - glaucoma
  - high blood pressure
- ask a doctor before use if you have:
  - heart disease
  - thyroid disease
  - diabetes
  - glaucoma
  - high blood pressure

**Directions**

- do not break or chew tablet; swallow tablet whole
- take 1 tablet every 12 hours, do not take more than 2 tablets in 24 hours.
- ask a doctor if you have:
  - heart disease
  - thyroid disease
  - diabetes
  - glaucoma
  - high blood pressure

**Warnings**

- do not use if you have had an allergic reaction to this product or any of its ingredients
- do not use if you are taking a prescription monoamine oxidase inhibitor (MAOI) (permanently for depression, psychiatric, or emotional conditions, or Parkinson's disease), or an MAOI within 14 days of the last dose of this product.
- ask a doctor before use if you have:
  - heart disease
  - thyroid disease
  - diabetes
  - glaucoma
  - high blood pressure

**Uses**

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
  - sneezing
  - runny nose
  - itchy, watery eyes
  - itching of the nose or throat
  - nasal congestion
  - reduces swelling of nasal passages
  - temporarily relieves sinus congestion and pressure
  - temporarily relieves sinus congestion and pressure

**Drug Facts**

**Active ingredients (in each extended release tablet)** Purpose  
Cetirizine HCl 5 mg Antihistamine  
Pseudoephedrine HCl 120 mg Nasal decongestant

**Drug Facts**

**Warnings**

- do not use more than directed
- do not use more than directed
- do not use more than directed

**Directions**

- do not use more than directed
- do not use more than directed
- do not use more than directed

**Other information**

- do not use if carton is opened or blister unit is broken
- store between 20° to 25°C (68° to 77°F)
- ask a doctor if you have:
  - heart disease
  - thyroid disease
  - diabetes
  - glaucoma
  - high blood pressure



**ALLERGY + CONGESTION**

# ZYRTEC-D

**ALLERGY + CONGESTION**

Cetirizine HCl 5 mg/antihistamine  
Pseudoephedrine HCl 120 mg/nasal decongestant  
Extended Release Tablets

**24**  
EXTENDED  
RELEASE TABLETS

Original Prescription Strength

NDC 50580-728-24

# ZYRTEC-D

Cetirizine HCl 5 mg/antihistamine  
Pseudoephedrine HCl 120 mg/nasal decongestant  
Extended Release Tablets

**ALLERGY + CONGESTION**

**INDOOR + OUTDOOR  
ALLERGIES**

**12 HOUR  
RELIEF OF**

- Sneezing
- Runny Nose
- Sinus Pressure
- Itchy, Watery Eyes
- Itchy Throat or Nose
- Nasal Congestion


(Actual Size)



**24**  
EXTENDED  
RELEASE TABLETS



30038756



30038756

cetirizine hydrochloride and pseudoephedrine hydrochloride tablet, film coated, extended release

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:50580-728
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>CETIRIZINE HYDROCHLORIDE</b> (UNII: 64O047KTOA) (CETIRIZINE - UNII:YO7261ME24)	CETIRIZINE HYDROCHLORIDE	5 mg
<b>PSEUDOEPHEDRINE HYDROCHLORIDE</b> (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	120 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	white (White to off white)	<b>Score</b>	no score
<b>Shape</b>	ROUND (Biconvex)	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	Zyrtec;D
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50580-728-12	2 in 1 CARTON	01/01/2008	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:50580-728-24	4 in 1 CARTON	01/01/2008	
2		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:50580-728-25	4 in 1 CARTON	01/01/2008	
3		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA021150	01/01/2008	

**Labeler** - Johnson & Johnson Consumer Inc. (878046358)

Revised: 3/2023

Johnson & Johnson Consumer Inc.