

SEVERE NIGHTTIME SEVERE DAYTIME- acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl

CVS Pharmacy

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

CVS Pharmacy, Inc. Daytime Nighttime Cold & Flu Relief Drug Facts

Nighttime Severe Cold & Flu

Active ingredients (in each 30 mL)

Acetaminophen 650 mg

Dextromethorphan HBr 20 mg

Doxylamine succinate 12.5 mg

Phenylephrine HCl 10 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion and pressure
- minor aches and pains
- headache
- fever
- sore throat
- runny nose and sneezing
- cough due to minor throat and bronchial irritation
- cough to help you sleep
- reduces swelling of nasal passages
- promotes nasal and/or sinus drainage
- temporarily restores freer breathing through the nose

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen

- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks

- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed - see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- **each 30 mL contains:** sodium 41 mg
- store at 20-25°C (68-77°F)

Inactive ingredients

anhydrous citric acid, edetate disodium, FD&C blue #1, FD&C red #40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose, xanthan gum

Questions or comments?

1-800-719-9260

Daytime Severe Cold & Flu

Active ingredients (in each 15 mL)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Guaifenesin 200 mg

Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion and pressure
- cough due to minor throat and bronchial irritation
- minor aches and pains
- headache
- fever
- sore throat
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- thyroid disease
- diabetes
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

When using this product

do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In care of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed - see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- **each 15 mL contains:** sodium 6 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

butylated hydroxyanisole, edetate disodium, FD&C yellow #6, flavor, glycerin, menthol, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum

Questions or comments?

1-800-719-9260

Principal Display Panel - Daytime

TWIN PACK

Compare to the active ingredients in Vicks® DayQuil® Severe

Severe

Non-Drowsy

Daytime

COLD & FLU RELIEF

ACETAMINOPHEN – Pain reliever; Fever reducer

DEXTROMETHORPHAN HBr – Cough suppressant

GUAIFENESIN – Expectorant

PHENYLEPHRINE HCl – Nasal decongestant

MAXIMUM STRENGTH

Relieves:

Aches; Fever; Cough; Nasal congestion; Sore throat; Chest congestion

Alcohol free

Antihistamine free

Compare to the active ingredients in Vicks® NyQuil® Severe

Severe

Nighttime

COLD & FLU RELIEF

ACETAMINOPHEN – Pain reliever; Fever reducer

DEXTROMETHORPHAN HBr – Cough suppressant

DOXYLAMINE SUCCINATE – Antihistamine

PHENYLEPHRINE HCl – Nasal decongestant

MAXIMUM STRENGTH

Relieves:

Aches; Fever; Sore throat; Cough; Sneezing; Runny nose; Nasal/sinus congestion & Sinus pressure

Berry Flavor

Alcohol free

TWO 12 FL OZ (355 mL) UNITS TOTAL 24 FL OZ (1.5 PT) (710 mL)

TWIN PACK



Compare to the active ingredients
in Vicks® DayQuil® Severe*

Severe



Compare to the active ingredients
in Vicks® NyQuil® Severe*

Severe

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

*These products are not manufactured or distributed
by Procter & Gamble, distributor of Vicks® DayQuil®
Severe and Vicks® NyQuil® Severe.

Non-Drowsy

Daytime

COLD & FLU RELIEF

ACETAMINOPHEN - Pain reliever; Fever reducer
DEXTROMETHORPHAN HBr - Cough suppressant
GUAIFENESIN - Expectorant
PHENYLEPHRINE HCl - Nasal decongestant

MAXIMUM STRENGTH

Relieves:

Aches; Fever; Cough;
Nasal congestion;
Sore throat;
Chest congestion



Alcohol free
Antihistamine free

TWO 12 FL OZ (355 mL) UNITS

NDC 59779-597-02

Nighttime

COLD & FLU RELIEF

ACETAMINOPHEN - Pain reliever; Fever reducer
DEXTROMETHORPHAN HBr - Cough suppressant
DOXYLAMINE SUCCINATE - Antihistamine
PHENYLEPHRINE HCl - Nasal decongestant

MAXIMUM STRENGTH

Relieves:

Aches; Fever; Sore throat;
Cough; Sneezing;
Runny nose; Nasal/sinus
congestion &
Sinus pressure



Berry Flavor
Alcohol free

TOTAL 24 FL OZ (1.5 PT) (710 mL)



Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2018 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
V-16430

**DO NOT USE IF PRINTED NECKBAND IS
BROKEN OR MISSING**

Drug Facts (continued)
Inactive ingredients butylated hydroxyanisole, edetate disodium, FD&C yellow #6, flavor, glycerin, menthol, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum.
Questions or comments? 1-800-719-9260

Drug Facts (continued)
Inactive ingredients antihydroxy citric acid, edetate disodium, FD&C blue #1, FD&C red #40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose, xanthan gum
Questions or comments? 1-800-719-9260

Drug Facts	Nighttime Severe Cold & Flu	Drug Facts	Daytime Severe Cold & Flu
Active ingredients (in each 30 mL)	Purpose Acetaminophen 650 mg.....Pain reliever/fever reducer Dextromethorphan HBr 2.0 mg.....Coughs suppressant Doxylamine succinate 1.25 mg.....Antihistamine Phenylephrine HCl 10 mg.....Nasal decongestant	Active ingredients (in each 15 mL)	Purpose Acetaminophen 325 mg.....Pain reliever/fever reducer Dextromethorphan HBr 10 mg.....Coughs suppressant Guaifenesin 2.00 mg.....Expectorant Phenylephrine HCl 5 mg.....Nasal decongestant
Uses temporarily relieves common cold/flu symptoms: ■ nasal congestion ■ sinus congestion and pressure ■ minor aches and pains ■ headache ■ fever ■ sore throat ■ runny nose and sneezing ■ cough due to minor throat and bronchial irritation ■ cough to help you sleep ■ reduces swelling of nasal passages ■ promotes nasal and/or sinus drainage ■ temporarily restores freer breathing through the nose	Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours. ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product. Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.	Uses temporarily relieves common cold/flu symptoms: ■ nasal congestion ■ sinus congestion and pressure ■ cough due to minor throat and bronchial irritation ■ minor aches and pains ■ headache ■ fever ■ sore throat ■ reduces swelling of nasal passages ■ temporarily restores freer breathing through the nose ■ promotes nasal and/or sinus drainage ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive	Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if ■ adult takes more than 4,000 mg of acetaminophen in 24 hours ■ adult takes more than 5 doses in 24 hours ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product. Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease, or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients	Ask a doctor before use if you have ■ liver disease ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ glaucoma ■ cough that occurs with too much phlegm (mucus) ■ a breathing problem such as emphysema or chronic bronchitis ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema ■ a sodium-restricted diet	Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease, or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients	Ask a doctor before use if you have ■ liver disease ■ heart disease ■ thyroid disease ■ diabetes ■ high blood pressure ■ trouble urinating due to an enlarged prostate gland ■ cough that occurs with too much phlegm (mucus) ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
Ask a doctor or pharmacist before use if you are ■ taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin	When using this product ■ do not use more than directed ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery ■ alcohol, sedatives, and tranquilizers may increase drowsiness	Ask a doctor or pharmacist before use if you are ■ taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin	When using this product ■ do not use more than directed
Stop use and ask a doctor if ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.	If pregnant or breast-feeding , ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	Stop use and ask a doctor if ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults) ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.	If pregnant or breast-feeding , ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Directions ■ take only as directed - see Overdose warning ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs adults & children 12 yrs & over 30 mL every 4 hrs children 4 to under 12 yrs ask a doctor children under 4 yrs do not use	Other information ■ each 30 mL contains: sodium 41 mg ■ store at 20-25°C (68-77°F)	Directions ■ take only as directed - see Overdose warning ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs adults & children 12 yrs & over 30 mL every 4 hrs children 6 to under 12 yrs 15 mL every 4 hrs children 4 to under 6 yrs ask a doctor children under 4 yrs do not use	Other information ■ each 15 mL contains: sodium 6 mg ■ store at 20-25°C (68-77°F). Do not refrigerate.

h 0 2 T 20265 :



SEVERE NIGHTTIME SEVERE DAYTIME

acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:59779-597
---------------------	----------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59779-597-02	1 in 1 CARTON; Type 0: Not a Combination Product	12/27/2013	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE	355 mL
Part 2	1 BOTTLE	355 mL

SEVERE NIGHTTIME

acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl solution

Product Information

Item Code (Source)	NDC:59779-763
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	RED (clear, dark)	Score	
Shape		Size	
Flavor	BERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59779-763-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	12/27/2013	

Part 2 of 2

SEVERE DAYTIME

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl solution

Product Information

Item Code (Source)	NDC:59779-603
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 15 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GLYCERIN (UNII: PDC6A3C0OX)	
MENTHOL (UNII: L7T10EIP3A)	
SODIUM PHOSPHATE, MONOBASIC (UNII: 3980JH2SW)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SUCROSE (UNII: C151H8M554)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	ORANGE (clear)	Score	
-------	----------------	-------	--

Shape		Size	
Flavor	FRUIT, MENTHOL	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59779-603-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	12/27/2013	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	12/27/2013	

Labeler - CVS Pharmacy (062312574)

Revised: 1/2019

CVS Pharmacy