

FEXOFENADINE HYDROCHLORIDE AND PSEUDOEPHEDRINE HYDROCHLORIDE-
fexofenadine hydrochloride and pseudoephedrine hydrochloride tablet, film coated, extended
release
walgreens

Fexofenadine Hydrochloride and Pseudoephedrine Hydrochloride Extended-Release Tablets, 60
mg/120 mg

Active ingredients

(in each extended-release tablet)

Fexofenadine HCl, USP 60 mg

Pseudoephedrine HCl, USP 120 mg

Purpose

Antihistamine

Nasal Decongestant

Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 - runny nose
 - sneezing
 - itchy, watery eyes
 - itching of the nose or throat
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily restores freer breathing through the nose

Warnings

Do not use

- if you have ever had an allergic reaction to this product or any of its ingredients
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have difficulty swallowing

Ask a doctor before use if you have

- heart disease
- thyroid disease
- glaucoma
- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland
- kidney disease. Your doctor should determine if you need a different dose.

When using this product

- **do not take more than directed**
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

Stop use and ask doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- you get nervous, dizzy, or sleepless

If pregnant or breast-feeding

ask a health professional before use.

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not divide, crush, chew or dissolve the tablet; swallow tablet whole

adults and children 12 years of age and over	take 1 tablet with a glass of water every 12 hours on an empty stomach; do not take more than 2 tablets in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

Other information

- do not use if carton is opened or if individual blister units are torn or opened.
- store between 20° to 25°C (68° to 77°F)
- USP dissolution test is pending.

Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, dibasic calcium phosphate dihydrate, ethyl cellulose, ferric oxide yellow, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, stearic acid.

Questions or comments?

call toll free 1-800-818-4555 weekdays

PRINCIPAL DISPLAY PANEL - 30 Tablet Blister Pack Carton

NON-DROWSY
ORIGINAL PRESCRIPTION STRENGTH

Walgreens

Wal-Fex[®] D

ALLERGY & CONGESTION

FEXOFENADINE HCl 60 mg / ANTIHISTAMINE
PSEUDOEPHEDRINE HCl 120 mg / NASAL DECONGESTANT

12 HOUR
EXTENDED-RELEASE TABLETS, USP

INDOOR & OUTDOOR ALLERGIES

- Nasal and Sinus Congestion Due to Colds or Allergies
- Sneezing; Runny Nose; Itchy, Watery Eyes and Itchy Nose or Throat Due to Allergies

30
EXTENDED-RELEASE
TABLETS

DO NOT USE IF INDIVIDUAL BLISTER UNIT IS OPEN OR TORN

Compare to Allegra-D[®] 12 Hour Allergy &
Congestion Tablets active ingredients^{††}

NDC 0363-2110-30

12
HOUR

ACTUAL SIZE

NO COATING

Drug Facts (continued)

Ask a doctor before use if you have

- heart disease
- thyroid disease
- glaucoma
- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland
- kidney disease. Your doctor should determine if you need a different dose.

When using this product

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- you get nervous, dizzy, or sleepless

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1800-222-1222).

Drug Facts (continued)

Directions

- do not divide, crush, chew or dissolve the tablet; swallow tablet whole

adults and children 12 years of age and over	take 1 tablet with a glass of water every 12 hours on an empty stomach; do not take more than 2 tablets in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

Other Information

- do not use if carton is opened or if individual blister units are torn or opened
- store between 20° to 25°C (68° to 77°F)
- USP dissolution test is pending.

Inactive Ingredients

colloidal silicon dioxide, croscarmellose sodium, dibasic calcium phosphate dihydrate, ethyl cellulose, ferric oxide yellow, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, stearic acid.

Questions or comments?
Call toll-free 1-800-818-4555 weekdays

NO COATING

[†]Walgreens Pharmacist Survey.
[‡]All trademarks are property of their respective owners.
This product is not affiliated with the makers of Allegra-D®.

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015

Walgreens
100% SATISFACTION GUARANTEED

walgreens.com ©2018 Walgreen Co.
MADE IN INDIA

ITEM 995657 W00000-0000-0



3 11917 19454 7

ORIG111&F
REV-0219

NON VARNISH

PHARMACIST
Walgreens
Trusted since 1901™
RECOMMENDED!

Health expertise you rely on.™

NON-DROWSY ORIGINAL PRESCRIPTION STRENGTH

Walgreens

Wal-Fex® D

ALLERGY & CONGESTION

Compare to Allegra-D® 12 Hour Allergy & Congestion Tablets active ingredients^{††}

NDC 0363-2110-30

FEXOFENADINE HCl 60 mg / ANTIHISTAMINE
PSEUDOEPHEDRINE HCl 120 mg / NASAL DECONGESTANT

12 HOUR EXTENDED-RELEASE TABLETS, USP

INDOOR & OUTDOOR ALLERGIES

- Nasal and Sinus Congestion Due to Colds or Allergies
- Sneezing; Runny Nose; Itchy, Watery Eyes and Itchy Nose or Throat Due to Allergies

30 EXTENDED-RELEASE TABLETS

DO NOT USE IF INDIVIDUAL BLISTER UNIT IS OPEN OR TORN

12 HOUR ACTUAL SIZE

NON VARNISH

6 x 5 Tablets

Drug Facts

Active ingredients (in each extended-release tablet)	Purpose
Fexofenadine HCl, USP 60 mg.....	Antihistamine
Pseudoephedrine HCl, USP 120 mg.....	Nasal Decongestant

Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 - runny nose
 - sneezing
 - itchy, watery eyes
 - itching of the nose or throat
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- reduces swelling of nasal passages

Drug Facts (continued)

Warnings
Do not use

- if you have ever had an allergic reaction to this product or any of its ingredients
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have difficulty swallowing

P68D0903A

FEXOFENADINE HYDROCHLORIDE AND PSEUDOEPHEDRINE

HYDROCHLORIDE

fexofenadine hydrochloride and pseudoephedrine hydrochloride tablet, film coated, extended release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-2110
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FEXO FENADINE HYDRO CHLORIDE (UNII: 2S068B75ZU) (FEXOFENADINE - UNII:E6582LOH6V)	FEXOFENADINE HYDROCHLORIDE	60 mg
PSEUDOEPHEDRINE HYDRO CHLORIDE (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	120 mg

Inactive Ingredients

Ingredient Name	Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
POVIDONE K30 (UNII: U725QWY32X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
ETHYLCELLULOSE, UNSPECIFIED (UNII: 7Z8S9VYZ4B)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	

Product Characteristics

Color	WHITE, YELLOW	Score	no score
Shape	CAPSULE (bilayer)	Size	17mm
Flavor		Imprint Code	724
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-2110-20	20 in 1 BLISTER PACK; Type 0: Not a Combination Product	04/17/2018	
2	NDC:0363-2110-30	30 in 1 BLISTER PACK; Type 0: Not a Combination Product	04/17/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

ANDA	ANDA090818	04/17/2018	
------	------------	------------	--

Labeler - walgreens (008965063)

Establishment

Name	Address	ID/FEI	Business Operations
Sun Pharmaceutical Industries Limited		650445203	ANALYSIS(0363-2110) , MANUFACTURE(0363-2110)

Revised: 2/2019

walgreens